

Symplectic Frieze Patterns

Sophie MORIER-GENOUD

Sorbonne Université, Université Paris Diderot, CNRS, Institut de Mathématiques de Jussieu-Paris Rive Gauche, IMJ-PRG, F-75005, Paris, France

E-mail: sophie.morier-genoud@imj-prg.fr

URL: <https://webusers.imj-prg.fr/~sophie.morier-genoud/>

Received June 18, 2019, in final form November 07, 2019; Published online November 14, 2019

<https://doi.org/10.3842/SIGMA.2019.089>

Abstract. We introduce a new class of friezes which is related to symplectic geometry. On the algebraic and combinatorics sides, this variant of friezes is related to the cluster algebras involving the Dynkin diagrams of type C_2 and A_m . On the geometric side, they are related to the moduli space of Lagrangian configurations of points in the 4-dimensional symplectic space introduced in [Conley C.H., Ovsienko V., *Math. Ann.* **375** (2019), 1105–1145]. Symplectic friezes share similar combinatorial properties to those of Coxeter friezes and SL-friezes.

Key words: frieze; cluster algebra; moduli space; difference equation; Lagrangian configuration

2010 Mathematics Subject Classification: 13F60; 05E10; 14N20; 53D30

To Dmitry Borisovich Fuchs on his 80th birthday

1 Introduction

The notion of friezes goes back to Coxeter in the early 70's [9]. Friezes are arrays of numbers where neighboring values are related by a local arithmetic rule modeled on the group SL_2 . Coxeter's friezes are surprisingly connected to many classical areas of mathematics such as projective geometry, number theory, enumerative combinatorics [7, 9]. Many variants of friezes have been recently studied in connection with more fields: cluster algebras, quiver representations, moduli spaces, integrable systems, algebraic combinatorics, see, e.g., [1, 2, 3, 5, 24, 25].

In [24], we establish a property called the “trianality” that identifies three different objects: SL_k -friezes, configurations of points in the projective spaces and periodic difference equations. A main application of the triality is that the combinatorics related to friezes allows to describe nice coordinate systems on the configurations spaces and in particular to exhibit a structure of cluster variety.

In [6], Lagrangian configurations of lines in the symplectic spaces are identified with symmetric periodic linear difference equations. Valentin Ovsienko suggested to complete the triality in the case of Lagrangian configurations with a combinatorial notion of “symplectic friezes”. This is what we do in the present paper in the case of 4-dimensional symplectic spaces by introducing a new family of friezes that we call “symplectic 2-friezes”.

Lagrangian configurations in dimension 4 are interpreted as a discrete version of Legendrian knots. Symplectic 2-friezes are a combinatorial interpretation of the space of Lagrangian configurations modulo Sp_4 -transformations. In particular they give special coordinate systems that provide a cluster variety structure on the moduli space of Lagrangian configurations.

The symplectic Lie group Sp_4 appears in two ways. From an algebraic point of view, the local rule for the symplectic 2-friezes is interpreted as cluster mutations involving the Dynkin diagram

This paper is a contribution to the Special Issue on Algebra, Topology, and Dynamics in Interaction in honor of Dmitry Fuchs. The full collection is available at <https://www.emis.de/journals/SIGMA/Fuchs.html>

of type C_2 associated to the Lie group Sp_4 . From a geometric point of view, the symplectic 2-friezes parametrize particular configurations of lines in the 4-dimensional symplectic space modulo the action of the Lie group Sp_4 .

Another particularity of the 4-dimensional case is that the “local frieze rule” used to define the symplectic 2-friezes involves (2×2) -determinants instead of 4×4 expected ones. This rule mixes the rule of the 2-friezes of [25] and the rule of wall numbers [8].

Symplectic 2-friezes enjoy nice properties which are very similar to those of the classical Coxeter friezes. In particular, they are invariant under the glide symmetry and therefore periodic. The periodicity of the friezes is a case of Zamolodchikov periodicity, see, e.g., [20].

In Section 2 we expose the main results of the paper: the combinatorial properties of the symplectic 2-friezes and the links with difference equations, symplectic geometry and cluster algebras. Section 3 contains examples illustrating the main results.

In Section 4 we study the combinatorial structure of the symplectic 2-friezes and characterize them in terms of SL-friezes. We explain the correspondence between symplectic 2-friezes and symmetric difference equations. We also give explicit determinantal formulas for the entries in the friezes.

In Section 5 we explain the link between the 2-friezes and symplectic geometry. The results in this section are mainly based on [6]. We identify the symplectic 2-friezes with Legendrian configurations in $\mathbb{C}P^3$ and discuss the relationship with symplectic forms.

Section 6 makes the link between symplectic 2-friezes and cluster algebras. The corresponding cluster algebras are generated from a product of Dynkin diagrams of type C_2 and A_w .

In Section 7, we discuss open questions about the combinatorics of symplectic 2-friezes with positive integer entries and about variants and more general symplectic friezes.

Appendixes A and B recollect important formulas and results on classical SL-friezes from [4] and [24] which are extensively used throughout the paper.

2 Main results

2.1 Combinatorial description and characterizations of the symplectic 2-friezes

Roughly speaking, friezes may be defined as arrangements of numbers in a planar strip such that neighboring entries forming a given pattern always satisfy the same arithmetic relationship. For instance, the original Coxeter’s friezes satisfy that every four adjacent entries a, b, c, d forming a square are related by $ad - bc = 1$, see [9]. Changing the arithmetic relation leads to different variants of friezes, see [23] for a survey on the subject.

We introduce a new family of friezes. We call *symplectic 2-frieze* an array of complex numbers (or polynomials, rational functions, etc.) in the plane satisfying the following conditions:

- the array has finitely many infinite rows, bounded top and bottom by a row of 1’s;
- the entries are alternatingly colored in black and white;
- the colored entries are subject to the local rules
 - every white entry is equal to the (2×2) -determinant of the matrix formed by the four adjacent black entries;
 - every square of a black entry is equal to the (2×2) -determinant of the matrix formed by the four adjacent white entries.

Symplectic 2-friezes are represented as follows

$$\begin{array}{cccccccccccc}
 \cdots & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \cdots \\
 & \vdots & & \vdots & & \vdots & & \vdots & & \vdots & & \\
 \cdots & \bullet & \circ & \bullet & \circ & \mathbf{B} & f & \bullet & \circ & \bullet & \circ & \cdots \\
 & & \circ & \bullet & \circ & \mathbf{A} & e & \mathbf{D} & h & \bullet & \circ & \bullet \\
 \cdots & \bullet & \circ & \bullet & \circ & \mathbf{C} & g & \bullet & \circ & \bullet & \circ & \cdots \\
 & \vdots & & \vdots & & \vdots & & \vdots & & \vdots & & \\
 \cdots & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \cdots
 \end{array} \tag{2.1}$$

in which the local rules read

$$AD - BC = e, \quad eh - fg = D^2, \quad \dots$$

The *width* of the frieze is the number of rows strictly between the top and bottom rows of 1's.

It is sometimes convenient to extend the array with additional rows of 0's above and below the bounding rows of 1's.

We will consider “tame” friezes which are friezes satisfying an extra condition of genericity (see Definition 4.2 for the details). For instance, friezes with no zero entries are all tame.

Symplectic 2-friezes can be compared to the 2-friezes of [25]. Recall that the latter are arrays as (2.1) with no color and in which the local rule reads with no squared values, i.e., for an ordinary 2-frieze the rule would be the same everywhere $AD - BC = e$, $eh - fg = D$, \dots . Symplectic 2-friezes can also be compared to the so called “number walls” for which the local rule in the array (2.1) with no color would be $AD + BC = e^2$, $eh + fg = D^2$, see [8] and J. Propp’s webpage for discussions on the subject.

The tame symplectic 2-friezes share the same properties of symmetry as Coxeter friezes [9] and as the 2-friezes of [25].

Theorem 2.1. *All tame symplectic 2-friezes of width w are $2(w + 5)$ -periodic. Moreover, the arrays are all invariant under a glide reflection with respect to the median line.*

It turns out that the subarray consisting of the black entries of a tame symplectic 2-frieze form a classical SL_4 -frieze. This allows the following characterization in terms of SL_k -friezes.

Theorem 2.2. *The set of tame symplectic 2-friezes of width w is in one-to-one correspondence with each of the following sets:*

- 1) *the tame SL_4 -friezes of width w in which every adjacent (3×3) -minors are equal to their central elements,*
- 2) *the tame SL_4 -friezes of width w that are invariant under a glide reflection with respect to the median line,*
- 3) *the tame SL_{w+1} -friezes of width 3 that are symmetric with respect to the middle row.*

Theorems 2.1 and 2.2 are proved in Section 4.5.

2.2 Symplectic 2-friezes and difference equations

Consider the system of linear difference equations of the form

$$V_i = a_i V_{i-1} - b_i V_{i-2} + a_{i-1} V_{i-3} - V_{i-4}, \tag{2.2}$$

for $i \in \mathbb{Z}$, where $a_i, b_i \in \mathbb{C}$ are given coefficients, and $(V_i)_i$ is a sequence of indeterminates or a “solution”.

Following [21, 24], we call such difference equation *n-superperiodic* if

$$\begin{array}{cccccccccccccccccccc}
\dots & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \dots \\
\dots & \mathbf{1} & \mathbf{2} & \mathbf{5} & \mathbf{4} & \mathbf{6} & \mathbf{4} & \mathbf{6} & \mathbf{3} & \mathbf{2} & \mathbf{1} & \mathbf{1} & \mathbf{4} & \mathbf{30} & \mathbf{10} & \mathbf{4} & \mathbf{1} & \mathbf{1} & \mathbf{2} & \dots & \dots \\
\dots & \mathbf{1} & \mathbf{1} & \mathbf{3} & \mathbf{14} & \mathbf{10} & \mathbf{20} & \mathbf{6} & \mathbf{3} & \mathbf{1} & \mathbf{1} & \mathbf{3} & \mathbf{14} & \mathbf{10} & \mathbf{20} & \mathbf{6} & \mathbf{3} & \mathbf{1} & \mathbf{1} & \dots & \dots \\
\dots & \mathbf{2} & \mathbf{1} & \mathbf{1} & \mathbf{4} & \mathbf{30} & \mathbf{10} & \mathbf{4} & \mathbf{1} & \mathbf{1} & \mathbf{2} & \mathbf{5} & \mathbf{4} & \mathbf{6} & \mathbf{4} & \mathbf{6} & \mathbf{3} & \mathbf{2} & \mathbf{1} & \dots & \dots \\
\dots & \mathbf{1} & \dots
\end{array} \tag{3.4}$$

Example 3.2 (glide reflection). Recall that a glide reflection is the composition of a reflection about a line and a translation along that line. The invariance mentioned in Theorem 2.1 can be easily observed in the above examples. For instance, we can display a fundamental domain in the array (3.4) that repeats under a glide reflection. Note that the glide symmetry implies the periodicity of the arrays.




Figure 1. Glide symmetry in the frieze (3.4).

Example 3.3 (corresponding SL-friezes). We illustrate Theorem 2.2. The SL_4 -frieze corresponding to a symplectic 2-frieze is simply given by the subarray of black entries. Applying the combinatorial Gale duality (see Appendix B.5) one obtains the corresponding SL_{w+1} -frieze of width 3.




Figure 2. The SL_3 -frieze of width 3 (left) and the SL_4 -frieze of width 2 (right) corresponding to the symplectic 2-frieze (3.3).

The SL-friezes corresponding to the symplectic 2-frieze (3.3) are given in Fig. 2. The friezes are related by Gale duality. One can check that $WD({}^tV) = 0$ where W and V are respectively the 3×7 and 4×7 matrices defined on Fig. 2, and D the 7×7 diagonal matrix with diagonal coefficients $(1, -1, 1, -1, 1, -1, 1)$.

Example 3.4 (corresponding difference equation). We illustrate Theorem 2.3. The array (3.3) provides the following 7-periodic sequences of coefficients:

$$\begin{aligned}
(\dots, a_0, a_1, \dots, a_6, \dots) &= (\dots, 6, 3, 1, 3, 4, 2, 1, \dots), \\
(\dots, b_0, b_1, \dots, b_6, \dots) &= (\dots, 3, 14, 1, 2, 6, 5, 1, \dots).
\end{aligned}$$

Let us check that the associated recurrence (2.2) is indeed 7-antiperiodic. Choose the initial values $(V_{-3}, V_{-2}, V_{-1}, V_0) = (x, y, z, t)$ and compute

$$\begin{aligned} V_1 &= 3V_0 - 14V_{-1} + 6V_{-2} - V_{-3} = 3t - 14z + 6y - x, \\ V_2 &= V_1 - V_0 + 3V_{-1} - V_{-2} = 2t - 11z + 5y - x, \\ V_3 &= 3V_2 - 2V_1 + V_0 - V_{-1} = t - 6z + 3y - x, \\ V_4 &= 4V_3 - 6V_2 + 3V_1 - V_0 = -x = -V_{-3}, \\ V_5 &= 2V_4 - 5V_3 + 4V_2 - V_1 = -y = -V_{-2}, \\ V_6 &= V_5 - V_4 + 2V_3 - V_2 = -z = -V_{-1}, \\ V_7 &= 6V_6 - 3V_5 + V_4 - V_3 = -t = -V_0. \end{aligned}$$

Example 3.5 (equations of the variety). For $w = 1$ the system of equations (2.3) is

$$\begin{aligned} a_3 &= a_6, & (1) \quad a_1 - a_3b_2 + a_2 &= 0, & (4) \quad b_1 - a_1a_3 + 1 &= 0, & (7) \\ a_3a_2 - b_3 &= 1, & (2) \quad a_2 - a_4b_3 + a_3 &= 0, & (5) \quad b_2 - a_2a_4 + 1 &= 0, & (8) \\ a_4a_3 - b_4 &= 1, & (3) \quad a_5 - a_3b_5 + a_4 &= 0, & (6) \quad b_5 - a_4a_2 + 1 &= 0, & (9) \\ & & & & & & b_6 - a_5a_3 + 1 &= 0. & (10) \end{aligned}$$

Note that using the first row expansion or last column expansion in $\Delta_{i,j}$ the subsets of equations in the rows of the system (2.3) can be immediately simplified using the previous subsets, in order to decrease the degrees of the equations.

The dimension of the variety is 2. The above system can be solved generically using two parameters $(a, b) \neq (0, 0)$. E.g., choosing $(a, b) = (a_3, b_3)$ we deduce b_4 from b_3b_4 using a combination of equations number 3, 5 and 2:

$$b_3b_4 = b_3(a_4a_3 - 1) = b_3a_4a_3 - b_3 = (a_2 + a_3)a_3 - b_3 = a_3^2 + (a_2a_3 - b_3) = a_3^2 + 1.$$

So that $b_4 = \frac{1+a_3^2}{b_3}$. Then all the variables are deduced one after each other using only one equation of the system, in the following order

$$\begin{aligned} a_2 &= \frac{1+b_3}{a_3}, & a_4 &= \frac{1+b_4}{a_3}, \\ b_2 &= a_2a_4 - 1, & a_1 &= a_3b_2 - a_2, & b_1 &= a_1a_3 - 1, \\ b_5 &= a_2a_4 - 1, & a_5 &= a_3b_5 - a_4, & b_6 &= a_5a_3 - 1, \\ a_6 &= a_3. \end{aligned}$$

Expressing everything in terms of $(a, b) = (a_3, b_3)$, one gets

$$\begin{aligned} b_1 &= b_4 = \frac{1+a^2}{b}, & a_1 &= a_4 = \frac{1+b+a^2}{ab}, \\ b_2 &= b_5 = \frac{(1+b)^2+a^2}{a^2b}, & a_2 &= a_5 = \frac{1+b}{a}, \\ b_3 &= b_6 = b, & a_3 &= a_6 = a. \end{aligned}$$

Example 3.6 (corresponding Legendrian n -gon). We illustrate Theorem 2.7. We can associate a Legendrian heptagon to the 2-frieze of width 2 given in (3.3). We choose a 4×7 -subarray of black entries in the frieze, e.g., the block displayed in Fig. 2, and denote by V_i its columns, so

that

$$\begin{array}{ccccccc} V_1 & V_2 & V_3 & V_4 & V_5 & V_6 & V_7 \\ \left(\begin{array}{ccccccc} 1 & 4 & 3 & 1 & 0 & 0 & 0 \\ 0 & 1 & 2 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 3 & 1 & 0 \\ 0 & 0 & 0 & 1 & 6 & 4 & 1 \end{array} \right) \end{array}$$

We extend the sequence by periodicity $V_{i+7} = V_i$.

Consider the symplectic form on \mathbb{C}^4 given by

$$\omega = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & -4 & 1 \\ -1 & 4 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}.$$

Attach to each V_i the hyperplane $H_i := \{V_i\}^{\perp\omega}$ formed by the orthogonal vectors. One can easily check that V_{i-1}, V_{i+1} belong to H_i for all i . The 7-periodic sequence $v = (CV_i)_i$ forms a Legendrian heptagon in $\mathbb{C}\mathbb{P}^3$.

From the Legendrian n -gon, one can compute the entries in the corresponding frieze using the values of

$$\omega(V_i, V_j).$$

E.g., in this example the black entries of the first row (which is the same as the second row by glide symmetry) of (3.3) is $\omega(V_2, V_5), \omega(V_3, V_6), \omega(V_4, V_7), \dots$

If we move the initial (4×7) -subarray along the north-east diagonal, the sequence changes by a shift of indices. If we move the initial (4×7) -subarray along the north-west diagonal, the sequence changes under the action of Sp_4 . Therefore any choice of (4×7) -subarrays leads to the same Legendrian n -gon, modulo a shift of indices of the vertices and modulo Sp_4 -action.

Example 3.7 (cluster variables in the frieze). Every symplectic 2-frieze is generically determined by the entries in 2 consecutive columns. All the other entries can be expressed as rational fractions of the initial entries using the frieze rule. These expressions can be recognized as cluster variables of type $C_2 \times A_w$, where w is the width of the frieze. For instance, a frieze of width $w = 1$ will produce the 6 cluster variables of type C_2 :

$$\begin{array}{cccccccccc} \cdots & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \cdots \\ \cdots & x_1 & \mathbf{x}_2 & \frac{1+x_2^2}{x_1} & \frac{1+\mathbf{x}_1+\mathbf{x}_2^2}{\mathbf{x}_1\mathbf{x}_2} & \frac{(1+x_1)^2+x_2^2}{x_1x_2^2} & \frac{1+\mathbf{x}_1}{\mathbf{x}_2} & x_1 & \mathbf{x}_2 & \cdots \\ \cdots & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \mathbf{1} & \cdots \end{array} \quad (3.5)$$

4 Symplectic 2-friezes, SL-friezes and difference equations

4.1 Notation for the symplectic 2-friezes

We complete the description of the 2-friezes given in Section 2.1 by introducing a labelling of the entries and more convention.

The entries in a symplectic 2-frieze of width w are denoted by $(d_{i,j})$, $i, j \in \mathbb{Z}$ for the ‘‘black entries’’ and $(d_{i+\frac{1}{2},j+\frac{1}{2}})$, $i, j \in \mathbb{Z}$ for the ‘‘white entries’’. The local rules around a white entry is

$$d_{i+\frac{1}{2},j+\frac{1}{2}} = d_{i,j}d_{i+1,j+1} - d_{i+1,j}d_{i,j+1},$$




Figure 3. Coordinate labelling in a symplectic 2-frieze. Black entries are connected with plain lines and the white entries with dashed lines.

and the local rule around a black entry is

$$d_{i,j}^2 = d_{i-\frac{1}{2},j-\frac{1}{2}} d_{i+\frac{1}{2},j+\frac{1}{2}} - d_{i+\frac{1}{2},j-\frac{1}{2}} d_{i-\frac{1}{2},j+\frac{1}{2}}.$$

The boundary conditions are

- $d_{i,i-1} = d_{i+\frac{1}{2},i-1+\frac{1}{2}} = 1,$
- $d_{i,i+w} = d_{i+\frac{1}{2},i+w+\frac{1}{2}} = 1.$

Note that the index set for the entries is

$$\{(i, j) \mid i \in \mathbb{Z}, i-1 \leq j \leq i+w\}.$$

However, we will often consider the 2-friezes as infinite arrays by adding the extra conventional conditions:

- $d_{i,i-\ell} = d_{i+\frac{1}{2},i-\ell+\frac{1}{2}} = 0,$ for $\ell = 2, 3, 4,$
- $d_{i,j+w+5} = -d_{i,j}$ for all $i, j,$

which means that we add three rows of zeros at the top and the bottom of the frieze and extend it to an antiperiodic infinite array.

Fig. 3 shows how the entries are organized in the plane. Note that a row in the frieze consists in the entries $d_{i,j}$ where $j-i$ is a fixed constant whereas i fixed or j fixed form the diagonals of the frieze.

4.2 The tameness condition

We define the notion of “tame” 2-friezes. This notion is an analog of the tameness condition for the classical SL-friezes, see Appendix B.1. It is understood as a genericity condition extending the properties satisfied by the friezes with no zero entries, see Proposition 4.1 and Remark 4.3.

Proposition 4.1. *A symplectic 2-frieze containing no zero entries satisfies the following properties.*

- (i) *All (3×3) -minors of adjacent black entries are equal to their central elements, i.e., for all $i, j \in \mathbb{Z}$*

$$\begin{vmatrix} d_{i-1,j-1} & d_{i-1,j} & d_{i-1,j+1} \\ d_{i,j-1} & d_{i,j} & d_{i,j+1} \\ d_{i+1,j-1} & d_{i+1,j} & d_{i+1,j+1} \end{vmatrix} = d_{i,j}. \quad (4.1)$$

- (ii) *All (4×4) -minors of adjacent black entries are equal to 1, i.e., for all $i, j \in \mathbb{Z}$*

$$\begin{vmatrix} d_{i-1,j-1} & d_{i-1,j} & d_{i-1,j+1} & d_{i-1,j+2} \\ d_{i,j-1} & d_{i,j} & d_{i,j+1} & d_{i,j+2} \\ d_{i+1,j-1} & d_{i+1,j} & d_{i+1,j+1} & d_{i+1,j+2} \\ d_{i+2,j-1} & d_{i+2,j} & d_{i+2,j+1} & d_{i+2,j+2} \end{vmatrix} = 1. \quad (4.2)$$

- (iii) *All (5×5) -minors of adjacent black entries are equal to 0, i.e., for all $i, j \in \mathbb{Z}$*

$$\begin{vmatrix} d_{i-1,j-1} & d_{i-1,j} & \cdots & d_{i-1,j+3} \\ d_{i,j-1} & d_{i,j} & \cdots & \vdots \\ \vdots & \vdots & \ddots & \vdots \\ d_{i+3,j-1} & \cdots & \cdots & d_{i+3,j+3} \end{vmatrix} = 0. \quad (4.3)$$

Proof. The key formula is the Desnanot–Jacobi determinantal identity, see Appendix A. Consider the following piece of symplectic 2-frieze, where the upper case (resp. lower case) entries are considered black entries (resp. white entries):

$$\begin{array}{cccccc} & & D & & & \\ & & C & c & H & \\ & B & b & G & f & L \\ A & a & F & e & K & i & P \\ & E & d & J & h & O \\ & & I & g & N & \\ & & & M & & \end{array}$$

By Desnanot–Jacobi identity one has the following relation between the minors

$$\begin{vmatrix} A & B & C \\ E & F & G \\ I & J & K \end{vmatrix} F = \begin{vmatrix} A & B \\ E & F \end{vmatrix} \begin{vmatrix} F & G \\ J & K \end{vmatrix} - \begin{vmatrix} E & F \\ I & J \end{vmatrix} \begin{vmatrix} B & C \\ F & G \end{vmatrix}. \quad (4.4)$$

Using the frieze rule in the above equation one obtains the relation

$$\begin{vmatrix} A & B & C \\ E & F & G \\ I & J & K \end{vmatrix} F = ae - bd = F^2,$$

from which we deduce (4.1) by cancelling out $F \neq 0$:

$$\begin{vmatrix} A & B & C \\ E & F & G \\ I & J & K \end{vmatrix} = F.$$

Using again Desnanot–Jacobi identity, and then (4.1) one obtains

$$\begin{aligned} \begin{vmatrix} A & B & C & D \\ E & F & G & H \\ I & J & K & L \\ M & N & O & P \end{vmatrix} \cdot \begin{vmatrix} F & G \\ J & K \end{vmatrix} &= \begin{vmatrix} A & B & C \\ E & F & G \\ I & J & K \end{vmatrix} \cdot \begin{vmatrix} F & G & H \\ J & K & L \\ N & O & P \end{vmatrix} \\ &- \begin{vmatrix} E & F & G \\ I & J & K \\ M & N & O \end{vmatrix} \cdot \begin{vmatrix} B & C & D \\ F & G & H \\ J & K & L \end{vmatrix} = FK - JG, \end{aligned}$$

from which we deduce (4.2) by cancelling out $FK - JG = e \neq 0$:

$$\begin{vmatrix} A & B & C & D \\ E & F & G & H \\ I & J & K & L \\ M & N & O & P \end{vmatrix} = 1.$$

Applying again Desnanot–Jacobi identity to compute the (5×5) -minors and using (4.1) and (4.2) already established one gets that all (5×5) -minors vanish. Hence (4.3) holds. \blacksquare

Definition 4.2. A symplectic 2-frieze is called *tame* if the three conditions (4.1), (4.2) and (4.3) are all satisfied.

By Proposition 4.1 the symplectic 2-friezes containing no zero entries are all tame. However the conditions of tame friezes allow to have zero entries, see Examples 4.4 and 4.5 below.

Remark 4.3. Note that generically the three conditions for the tameness property of a symplectic 2-frieze are automatically satisfied according to the arguments given in the proof of Proposition 4.1. More precisely,

- (4.1) is satisfied for a given (i, j) whenever $d_{i,j} \neq 0$;
- when (4.1) is satisfied, then (4.2) is automatically satisfied for a given (i, j) whenever $d_{i+\frac{1}{2}, j+\frac{1}{2}} \neq 0$;
- when (4.1) and (4.2) are satisfied, then (4.3) is automatically satisfied for a given (i, j) whenever $d_{i+1, j+1} \neq 0$.

Example 4.4. (a) In the following array of width one, containing a black zero entry, the frieze rule is satisfied for any values of x :

$$\begin{array}{cccccccccccc} \dots & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \dots \\ \dots & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & \dots \\ \dots & -1 & \mathbf{1} & -2 & \mathbf{1} & -1 & \mathbf{0} & -1 & \mathbf{x} & -1 - x^2 & \dots \\ \dots & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & \dots \\ \dots & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \dots \end{array}$$

The next two entries on the right can be computed uniquely whenever $x(1+x^2) \neq 0$. But if we want the array to be tame, the condition (4.1) centered at the black 0 imposes

$$\begin{vmatrix} -1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & x \end{vmatrix} = 0,$$

so that it determines $x = 1$. This will lead to the (6-periodic) tame frieze (3.1).

(b) In the following array, containing a white zero entry, the frieze rule is satisfied for any values of x (that will then uniquely determine y and z):

$$\begin{array}{cccccccc} \dots & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \dots \\ \dots & \sqrt{2} & 1 & \sqrt{2} & 0 & -\sqrt{2}+1 & x & y & \dots \\ \dots & -1 & \mathbf{1} & -2 & -\mathbf{2}+\sqrt{2} & -3+2\sqrt{2} & \mathbf{1} & z & \dots \\ \dots & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & \dots \end{array}$$

The tameness condition (4.1) is also satisfied for any values of x , since there are no black zeroes. Condition (4.2), centered at the white 0, imposes

$$\begin{vmatrix} \sqrt{2} & 1 & 0 & 0 \\ 1 & \sqrt{2} & 1 & 0 \\ 1 & -2+\sqrt{2} & -\sqrt{2}+1 & 1 \\ 0 & 1 & 1 & y \end{vmatrix} = 1,$$

so that one solves $y = \sqrt{2} - 1$, and then $x = -4 + 2\sqrt{2}$, $z = 1$. Then there is a unique way to extend the array to get a tame symplectic 2-frieze, which will be 7-periodic.

Example 4.5. (a) The following symplectic 2-frieze

$$\begin{array}{cccccccccccc} \dots & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \dots \\ \dots & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & \dots \\ \dots & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \dots \\ \dots & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & \dots \end{array}$$

is not tame because condition (4.1), centered at a black 0 of the second row, fails

$$\begin{vmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{vmatrix} \neq 0.$$

(b) The following symplectic 2-frieze

$$\begin{array}{cccccccccccc} \dots & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \dots \\ \dots & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & \dots \\ \dots & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \dots \\ \dots & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & \dots \\ \dots & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \dots \end{array}$$

is not tame because condition (4.2), centered at a white 0 of the third row, fails

$$\begin{vmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{vmatrix} \neq 1.$$

(c) There is only one tame symplectic 2-frieze with constant first row equal to 0. It is the following frieze of width 7:

$$\begin{array}{cccccccc} \dots & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & \dots \\ \dots & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \dots \\ \dots & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & \dots \\ \dots & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \dots \\ \dots & 1 & -\mathbf{1} & 1 & -\mathbf{1} & 1 & -\mathbf{1} & 1 & -\mathbf{1} & 1 & -\mathbf{1} & \dots \\ \dots & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \dots \\ \dots & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & \dots \\ \dots & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \mathbf{0} & 0 & \dots \\ \dots & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & \dots \end{array} \tag{4.5}$$

Remark 4.6. The conventional choice to extend the array of a symplectic 2-frieze with three rows of zeros at the top and bottom and then by antiperiodicity $d_{i+w+5,j} = d_{i,j+w+5} = -d_{i,j}$ (cf. Section 4.1) is justified as the only choice that extends a 2-frieze with no zero entry to an infinite tame array.

4.3 SL-friezes and difference equations

By definition, the tameness of the symplectic frieze forces the subarray of the black entries to form a tame SL_4 -frieze. Hence, many properties of the symplectic friezes are immediate consequences of known results on SL-friezes, see Appendix B.

Proposition 4.7. *Let w be the width of the friezes.*

- (i) *The subarray of the black entries in a tame symplectic 2-frieze forms a tame SL_4 -frieze in which the adjacent (3×3) -minors are equal to the central element. Conversely, every such SL_4 -frieze determines a unique tame symplectic 2-frieze.*
- (ii) *The subarray of the black entries in a tame symplectic 2-frieze forms a tame SL_4 -frieze invariant under the glide reflection along the median line:*

$$d_{j-w-2,i-3} = d_{i,j} = d_{j+3,i+w+2}, \quad i \leq j \leq i + w - 1.$$

Conversely, every such SL_4 -frieze determines a unique tame symplectic 2-frieze.

- (iii) *In a tame symplectic 2-frieze the entries $a_i := d_{i,i}$, $b_i := d_{i-\frac{1}{2},i-\frac{1}{2}}$ define the coefficients of a $(w+5)$ -superperiodic equation of the form (2.2). Moreover, the sequences on the diagonals $(d_{i_0,i})_{i \in \mathbb{Z}}$, for every fixed $i_0 \in \mathbb{Z}$, are solutions of the equation (2.2), i.e., for all $i_0 \in \mathbb{Z}$*

$$d_{i_0,i} = a_i d_{i_0,i-1} - b_i d_{i_0,i-2} + a_{i-1} d_{i_0,i-3} - d_{i_0,i-4}, \quad \text{for all } i \in \mathbb{Z}.$$

Conversely, every n -superperiodic equation of the form (2.2) determines a unique tame symplectic 2-frieze.

Proof. In a tame symplectic 2-frieze, the subarray of the black entries forms a tame SL_4 -frieze according to conditions (4.2) and (4.3) required in Definition 4.2 for the tame symplectic friezes. From [24], we already know that this SL_4 -frieze corresponds to a superperiodic difference equation of order 4, see Proposition B.1 for $k = 3$. Let us simply denote by

$$V_i = a_i V_{i-1} - b_i V_{i-2} + c_i V_{i-3} - V_{i-4}, \tag{4.6}$$

the corresponding superperiodic equation, where we use the simpler notation a_i, b_i, c_i for the constant coefficients a_i^1, a_i^2, a_i^3 of (B.2). These coefficients can be explicitly computed from the entries of the frieze, see (B.3).

In general SL_k -friezes are not invariant under a glide reflection for $k \geq 3$, unlike Coxeter's friezes (i.e., SL_2 -friezes). We first show the following lemma giving equivalent conditions to the invariance under a glide reflection in the case of SL_4 -friezes.

Lemma 4.8. *Given a tame SL_4 -frieze, the following properties are equivalent:*

- (1) *all (3×3) -minors of adjacent entries are equal to their central elements,*
- (2) *the array is invariant under a glide reflection along the median line,*
- (3) *the corresponding superperiodic equation is of the form (2.2).*

Proposition 4.9. *Consider a tame symplectic 2-frieze, and let $a_i := d_{i,i}$, $b_i := d_{i-\frac{1}{2},i-\frac{1}{2}}$ be the entries in the non trivial bottom row. The entries of the frieze are given by*

$$d_{i,j} = \begin{vmatrix} a_i & b_{i+1} & a_{i+1} & 1 & & & \\ 1 & a_{i+1} & b_{i+2} & a_{i+2} & 1 & & \\ & \ddots & \ddots & \ddots & \ddots & \ddots & \\ & & 1 & a_{j-2} & b_{j-1} & a_{j-1} & \\ & & & 1 & a_{j-1} & b_j & \\ & & & & 1 & a_j & \end{vmatrix},$$

$$d_{i-\frac{1}{2},j-\frac{1}{2}} = \begin{vmatrix} b_i & a_i & 1 & & & & \\ a_i & b_{i+1} & a_{i+1} & 1 & & & \\ 1 & a_{i+1} & b_{i+2} & a_{i+2} & 1 & & \\ & \ddots & \ddots & \ddots & \ddots & \ddots & \\ & & 1 & a_{j-2} & b_{j-1} & a_{j-1} & \\ & & & 1 & a_{j-1} & b_j & \end{vmatrix},$$

for all $i, j \in \mathbb{Z}$.

Proof. We know by Proposition 4.7(iii) that (a_i, b_i) give the coefficients of the superperiodic equation associated with the SL_4 -frieze formed by the black entries of the symplectic frieze. Hence, the formula for $d_{i,j}$ is given by (B.4). Using the frieze rule $d_{i-\frac{1}{2},j-\frac{1}{2}} = d_{i,j}d_{i-1,j-1} - d_{i-1,j}d_{i,j-1}$ and Desnanot–Jacobi identity in the determinant of $d_{i-1,j}$ one deduces the above formula for $d_{i-\frac{1}{2},j-\frac{1}{2}}$. ■

4.5 Proofs of Theorems 2.1, 2.2 and 2.3

The theorems are mainly consequences and reformulations of Proposition 4.7. Theorem 2.1 is an immediate consequence of Proposition 4.7(ii). Theorem 2.2(1) and (2) are also immediate consequences of Proposition 4.7(i) and (ii). Theorem 2.2(3) is obtained by applying the Gale duality, see Appendix B.5. More precisely, if a_i, b_i are the coefficients of the associated difference equation (i.e., the entries in the first row of the symplectic frieze) then the corresponding SL_{w+1} -frieze of width 3 is given as the following array

$$\begin{array}{cccccc} 1 & 1 & 1 & 1 & 1 & \\ \cdots & a_0 & a_1 & a_2 & a_3 & \cdots \\ & b_0 & b_1 & b_2 & b_3 & \cdots \\ \cdots & a_0 & a_1 & a_2 & a_3 & \cdots \\ & 1 & 1 & 1 & 1 & 1 \end{array}$$

Theorem 2.3 is an immediate consequence of Proposition 4.7(iii).

5 Moduli space of Legendrian configurations

5.1 Lift of the projective n -gons in \mathbb{C}^4

Consider the symplectic space (\mathbb{C}^4, ω_0) where ω_0 is the symplectic form given by the matrix

$$\Omega_0 = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}.$$

Applying $\omega_0(V_{i-2}, -)$ in (5.3) and using the normalization conditions of (1), we get $d_i = 1$ for all $i \in \mathbb{Z}$. Applying $\omega_0(V_{i-3}, -)$ in (5.3), we obtain $a_i = \omega_0(V_{i-3}, V_i)$ for all $i \in \mathbb{Z}$, that gives by a shift $a_{i-1} = \omega_0(V_{i-4}, V_{i-1})$. And finally applying $\omega_0(-, V_{i-1})$ in (5.3), we obtain $c_i = \omega_0(V_{i-4}, V_{i-1}) = a_{i-1}$. Item (3) is proved.

Let v and v' be two Legendrian n -gons such that their normalized lifts V and V' satisfy the same recurrence relation (5.1). Define T in GL_4 such that $TV_i = V'_i$ for $i = 1, 2, 3, 4$. The recurrence relation implies $TV_i = V'_i$ for all i . From the normalization conditions of (1) and the fact that $a_4 = \omega_0(V_1, V_4) = \omega_0(V'_1, V'_4)$, one can see that the matrix T preserves ω_0 . Hence, T is an element of Sp_4 that transforms v to v' .

Conversely if v' and v are PSp_4 -equivalent then there exists $T \in \mathrm{Sp}_4$ that transforms the normalized lifts V to V' . Therefore the sequences V and V' satisfy the same recurrence relation. Item (4) is proved. \blacksquare

Remark 5.2 (proof of Theorem 2.7). From Proposition 5.1 one can identify the Legendrian n -gons, modulo PSp_4 -equivalence, with n -superperiodic equations of the form (5.1), provided n is odd. Therefore Theorem 2.7 is an immediate consequence of Theorem 2.3.

5.2 Symplectic 2-friezes from Legendrian n -gons

One can obtain a tame symplectic 2-frieze from the lift of a Legendrian n -gon using the canonical symplectic form ω_0 on \mathbb{C}^4 .

Proposition 5.3. *Let $v = (v_i)$ be a Legendrian n -gon in \mathbb{CP}^3 and $V = (V_i)$ be its normalized lift in (\mathbb{C}^4, ω_0) given by Proposition 5.1. The following assignment*

$$d_{i,j} := \omega_0(V_{i-3}, V_j), \quad \text{for all } i, j \in \mathbb{Z}$$

defines the black array of a tame symplectic 2-frieze of width $w = n - 5$.

Proof. Using the normalization properties of $V = (V_i)$ and the linearity of ω_0 one deduces that the sequences

$$d_{i,\bullet} = (0, \dots, 0, 1, d_{i,i}, d_{i,i+1}, \dots, d_{i,i+w-1}, 1)$$

are solutions of the same $(w + 5)$ -superperiodic equation of the form (5.1) with different initial conditions

$$(V_{i-4}, \dots, V_{i-1}) = (0, \dots, 0, 1).$$

By Proposition B.1 this implies that $(d_{i,j})$ form a tame SL_4 -frieze. And since the coefficients of the equation are symmetric it implies by Proposition 4.7 that this SL_4 -frieze is indeed the black array of a tame symplectic 2-frieze. \blacksquare

Remark 5.4. Alternatively, one can compute the symplectic 2-frieze using 4×4 -determinants involving four vertices of the n -gon:

$$d_{i,j} = \det(V_{i-4}, V_{i-3}, V_{i-2}, V_j), \quad d_{i-\frac{1}{2}, j-\frac{1}{2}} = \det(V_{i-4}, V_{i-3}, V_{j-1}, V_j).$$

This is obtained as a particular case of the general correspondence between n -gons and SL_k -friezes without considering the extra symplectic structure, see, e.g., formula (B.8).

5.3 Symplectic forms and 2-friezes

Consider the following family of matrices of symplectic forms

$$\Omega_a = \begin{pmatrix} 0 & 0 & 1 & a \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ -a & -1 & 0 & 0 \end{pmatrix}, \quad \check{\Omega}_a = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & -a & 1 \\ -1 & a & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix} = {}^t\Omega_a^{-1} = -\Omega_a^{-1},$$

where a is a complex parameter.

Proposition 5.5. *In a symplectic 2-frieze, every 4×4 submatrices of black entries*

$$D_{i,j} := \begin{pmatrix} d_{i,j-3} & \cdots & d_{i,j} \\ \vdots & & \vdots \\ d_{i+3,j-3} & \cdots & d_{i+3,j+3} \end{pmatrix} \quad (5.4)$$

satisfies

$${}^tD_{i,j}\check{\Omega}_{a_i}D_{i,j} = \Omega_{a_j},$$

where $a_i = d_{i,i}$ are the black entries in the first (and last) row of the 2-frieze.

Proof. It is easy to check that the property holds for the matrices $D_{i,i}$ since $D_{i,i} = \Omega_{a_i}$. By the recurrence relation in the frieze one has $D_{i,j+1} = D_{i,j}E_{j+1}$, where E_{j+1} is the matrix

$$E_{j+1} := \begin{pmatrix} 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & a_j \\ 0 & 1 & 0 & -b_j \\ 0 & 0 & 1 & a_j \end{pmatrix}. \quad (5.5)$$

Hence, the property for all $D_{i,j}$ follows by easy induction on j . ■

5.4 Proof of Proposition 2.4

Let M be a monodromy matrix associated to the equation (2.2). It can be defined as the product $M = E_1E_2 \cdots E_n$, where E_i are the matrices (5.5). The condition of superperiodicity of (2.2) translates as $M = -\text{Id}$. We want to prove that the algebraic conditions on the coefficients (a_i) , (b_i) given by $M = -\text{Id}$ are equivalent to the one given in system (2.3).

We apply the matrix M on the 4×4 black subarray of the symplectic 2-frieze written as the matrix (5.4) for $i = j = 0$, i.e., on the matrix

$$D_{0,0} = \begin{pmatrix} 0 & 0 & 1 & a_0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ -a_0 & -1 & 0 & 0 \end{pmatrix}.$$

The recurrence relations in the frieze imply that $D_{0,0}M = D_{0,n}$. Hence, $M = -\text{Id}$ is equivalent to $D_{0,n} = -D_{0,0}$. This gives 16 equations when identifying the entries of the matrices. However, since $D_{0,n}$ satisfies the relation ${}^tD_{0,n}\check{\Omega}_{a_0}D_{0,n} = \Omega_{a_0}$, according to Proposition 5.5, it is enough to identify the ten entries in the triangular lower parts of the matrices, i.e.,

$$\begin{pmatrix} d_{0,n-3} & \cdot & \cdot & \cdot \\ d_{1,n-3} & d_{1,n-2} & \cdot & \cdot \\ d_{2,n-3} & d_{2,n-2} & d_{2,n-1} & \cdot \\ d_{3,n-3} & d_{3,n-2} & d_{3,n-1} & d_{3,n} \end{pmatrix} = \begin{pmatrix} 0 & \cdot & \cdot & \cdot \\ 0 & 0 & \cdot & \cdot \\ 1 & 0 & 0 & \cdot \\ a_0 & 1 & 0 & 0 \end{pmatrix},$$

which lead to the system (2.3) when using the determinantal expressions for the entries $d_{i,j}$ given in Proposition 4.9.

5.5 More correspondences

We explain different ways to obtain symplectic 2-friezes, Legendrian n -gons, and difference equations one from another.

If one knows the (a_i, b_i) -parameters of the Legendrian n -gon as defined in Proposition 5.1, or equivalently the (a_i, b_i) -coefficients of the associated difference equation (2.2), then the corresponding 2-frieze can be computed in two alternative easy ways.

1. One can write down the coefficients in the first non-trivial row $(\dots, b_i, \mathbf{a}_i, b_{i+1}, \mathbf{a}_{i+1}, \dots)$ of the array, and compute the rest of the entries by applying the frieze rule.
2. One can recursively compute the diagonals of the black entries of the 2-frieze using the equations (2.2) and initial conditions $(0, \dots, 0, 1) = (V_{i-4}, \dots, V_{i-1}) = (d_{i,i-4}, \dots, d_{i,i-1})$.

The first procedure is limited to the case of generic elements, while the second one can be applied in all cases.

Now, we describe the correspondence between symplectic 2-friezes and Legendrian n -gons without using explicitly the coefficients of the recurrence relations. For the rest of the section, we assume n is odd.

Starting from a Legendrian n -gon $v = (v_i)_{i \in \mathbb{Z}}$, we first consider the normalized lift $V = (V_i)$ given by Proposition 5.1. We transform (V_i) to (W_i) where $W_i = TV_i$ for the matrix T of Sp_4 defined by

$$T(V_{-3}) = \begin{pmatrix} 0 \\ 0 \\ -1 \\ -a_0 \end{pmatrix}, \quad T(V_{-2}) = \begin{pmatrix} 0 \\ 0 \\ 0 \\ -1 \end{pmatrix}, \quad T(V_{-1}) = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \quad T(V_0) = \begin{pmatrix} a_0 \\ 1 \\ 0 \\ 0 \end{pmatrix}$$

with $a_0 = \omega_0(V_{-3}, V_0)$.

The sequence W will have the following form

$$\begin{pmatrix} W_{-3} & W_{-2} & W_{-1} & W_0 & W_1 & W_2 & W_3 & \cdots & \cdots & W_{n-3} & W_{n-2} & W_{n-1} & W_n \\ \begin{pmatrix} 0 & 0 & 1 & a_0 & \bullet & \cdots & \bullet & 1 & 0 & 0 & 0 & -1 & -a_0 \\ 0 & 0 & 0 & 1 & a_1 & \bullet & \cdots & \bullet & 1 & 0 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 & 1 & a_2 & \bullet & \cdots & \bullet & 1 & 0 & 0 & 0 \\ -a_0 & -1 & 0 & 0 & 0 & 1 & a_3 & \bullet & \cdots & \bullet & 1 & 0 & 0 \end{pmatrix} \end{pmatrix} \quad (5.6)$$

and can uniquely be extended in a tame SL_4 -frieze. Adding the 2×2 minors inside the SL_4 -frieze will lead to a symplectic 2-frieze.

Note that the sequence W does not satisfy the normalization given in Proposition 5.1(1) for ω_0 but for the form given by Ω_{a_0} of Section 5.3

Conversely, if one cuts a $4 \times n$ block of the form (5.6) in the black subarray of a symplectic 2-frieze it gives n points in \mathbb{C}^4 satisfying the condition of Proposition 5.1(1) for some symplectic form Ω_a . After renormalizing the points they will project to a Legendrian n -gons in $(\mathbb{C}\mathbb{P}^3, \omega_0)$.

5.6 The case when n is even

In the case when n is even the system of equations (5.2) splits into two subsystems: one for the λ_i with i odd and one for the λ_i with i even. If n is a multiple of 4 the set of solutions for the subsystems is in general empty, otherwise each subsystem has two opposite sets of solutions $\pm(\lambda_1, \lambda_3, \dots, \lambda_{n-1})$ and $\pm(\lambda_2, \lambda_4, \dots, \lambda_n)$. So that we get four sequences of lifted points $\pm(V_1, \pm V_2, \dots, V_{2i-1}, \pm V_{2i}, \dots)$ satisfying the normalisation conditions of Proposition 5.1, which

define the same parameters (a_i, b_i) up to a sign for the coefficients a_i . This will lead to two distinct 2-friezes that differ under the action by conjugation of the matrix

$$\begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

on the black subarray.

For instance, the 2-frieze of width 3 given in (3.4) and the following one

$$\begin{array}{cccccccccccccccccccc} \dots & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \dots \\ \dots & -\mathbf{1} & 1 & -\mathbf{2} & 5 & -\mathbf{4} & 6 & -\mathbf{4} & 6 & -\mathbf{3} & 2 & -\mathbf{1} & 1 & -\mathbf{4} & 30 & -\mathbf{10} & 4 & -\mathbf{1} & 1 & \dots \\ \dots & 3 & \mathbf{1} & 1 & \mathbf{3} & 14 & \mathbf{10} & 20 & \mathbf{6} & 3 & \mathbf{1} & 1 & \mathbf{3} & 14 & \mathbf{10} & 20 & \mathbf{6} & 3 & \mathbf{1} & \dots \\ \dots & -\mathbf{3} & 2 & -\mathbf{1} & 1 & -\mathbf{4} & 30 & -\mathbf{10} & 4 & -\mathbf{1} & 1 & -\mathbf{2} & 5 & -\mathbf{4} & 6 & -\mathbf{4} & 6 & -\mathbf{3} & 2 & \dots \\ \dots & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \dots \end{array}$$

correspond to the same class of symplectic 8-gons.

5.7 Legendrian n -gons and self-dual polygons

Let $v = (v_i)$ be a Legendrian n -gon in \mathbb{CP}^3 , for some odd integer n . The symmetry of the coefficients in the corresponding difference equation (5.1) (or equivalently the symmetry in the corresponding SL_4 -friezes) implies that the polygon v can be identified with its dual polygon v^* in $(\mathbb{CP}^3)^*$. More precisely there is a projective map that sends v_i to v_{i+n-2}^* . The shift can be computed in terms of SL_4 -frieze using the formula (B.7) and the glide symmetry.

Legendrian n -gons are $(n-2)$ -self-dual polygons, in the terminology of [17]. The space of Legendrian n -gons is of dimension $2(n-5)$ and provides a 3-dimensional analog of the space $\mathcal{M}_{m,n}$ of [17].

6 Cluster algebras

6.1 Preliminaries

Cluster algebras are due to Fomin and Zelevinsky [13, 14]. They are commutative associative algebras defined by generators and relations inside a field of fractions. The generators and relations of a cluster algebra are not given from the beginning but are produced recursively.

A *seed* in a cluster algebra is a couple $\Sigma = ((u_1, \dots, u_m), B)$, where (u_1, \dots, u_m) is a transcendence basis of $\mathbb{C}(x_1, \dots, x_m)$ over \mathbb{C} and B is a skew-symmetrizable matrix of integer entries (alternatively the matrices can be replaced by quivers). The initial seed $\Sigma_0 = ((x_1, \dots, x_m), B_0)$ is the initial data needed to generate the entire cluster algebra. All seeds are created by a process of mutation from Σ_0 . The variables created by mutations are called *cluster variables*, they come grouped in a m -tuple called *cluster*. The complex cluster algebra \mathcal{A}_{Σ_0} is the subalgebra of $\mathbb{C}(x_1, \dots, x_m)$ generated by all the cluster variables.

To fix notation and convention we recall briefly the main definitions related to cluster algebras but refer to the original papers [13, 14] or the introductory texts [12, 19] for details. We also use the approach of [11] and [20].

6.1.1 Mutations from skew-symmetrizable matrices

The mutation μ_k , $1 \leq k \leq m$, of a seed $\Sigma = ((u_1, \dots, u_m), B)$ where $B = (b_{i,j})_{1 \leq i, j \leq m}$, is a skew-symmetrizable matrix with integer entries, is a new seed $\Sigma' = \mu_k(\Sigma) = ((u'_1, \dots, u'_m), B')$

where the new variables are given by

$$u'_k = \frac{1}{x_k} \left(\prod_{i: b_{ik} > 0} u_i^{b_{ik}} + \prod_{i: b_{ik} < 0} u_i^{-b_{ik}} \right),$$

$$u'_\ell = u_\ell, \quad \ell \neq k,$$

and the new matrix $\mu_k(B) = B' = (b'_{i,j})_{1 \leq i, j \leq m}$ given by

$$b'_{ij} = \begin{cases} -b_{ij} & \text{if } i = k \text{ or } j = k, \\ b_{ij} + b_{ik}b_{kj} & \text{if } b_{ik} > 0 \text{ and } b_{kj} > 0, \\ b_{ij} - b_{ik}b_{kj} & \text{if } b_{ik} < 0 \text{ and } b_{kj} < 0, \\ b_{ij} & \text{otherwise.} \end{cases}$$

6.1.2 Valued quivers

The valued quiver $Q_v(B)$ associated to a skew-symmetrizable matrix $B = (b_{i,j})_{1 \leq i, j \leq m}$ with integer entries, is defined as follows:

- $\{1, \dots, m\}$ is the set of vertices,
- each entry $b_{i,j} > 0$ gives a weighted arrow $i \rightarrow j$ of weight $(|b_{i,j}|, |b_{j,i}|)$.

Recall that in a skew-symmetrizable matrix one has $b_{i,j} > 0$ if and only if $b_{j,i} \leq 0$. When there is a weighted arrow between the vertices i and j the weight $|b_{i,j}|$ is “attached” to the vertex i and the weight $|b_{j,i}|$ “attached” to the vertex j . On the graph of a valued quiver we write the weights on the top or bottom of the arrows. The weighted arrow $i \xrightarrow{(a,b)} j$ can be also represented by $j \xleftarrow{(b,a)} i$.


In the case when $(|b_{i,j}|, |b_{j,i}|) = (b, b)$, we replace the weighted arrow by b arrows. For consistence of formulas it may be convenient to adopt the following convention: $i \xrightarrow{(0,0)} j$ means no arrow between i and j , and $i \xrightarrow{(-a,-b)} j$ means $i \xleftarrow{(a,b)} j$.

Our notation differs from that of [20].

Example 6.1. For example the matrix

$$\begin{pmatrix} 0 & 1 & 0 & -1 \\ -2 & 0 & -1 & 4 \\ 0 & 1 & 0 & -2 \\ 1 & -2 & 1 & 0 \end{pmatrix}$$


gives the following valued quiver




6.1.3 Mutations of valued quivers

The mutation at vertex k of a valued quiver Q_v gives a valued quiver $\mu_k(Q_v)$ defined by the following transformations:

1. For all paths $i \rightarrow k \rightarrow j$ in Q , change the weighted arrows



and



where $a, b, c, d > 0$ and $e, f, g, h \geq 0$.

2. Reverse the weighted arrows incident with k , i.e., change all

$$i \xrightarrow{(a,b)} k \quad \text{to} \quad i \xleftarrow{(a,b)} k$$

and

$$k \xrightarrow{(c,d)} j \quad \text{to} \quad k \xleftarrow{(c,d)} j.$$

This rule gives the correspondence $\mu_k(Q_v(B)) = Q_v(\mu_k(B))$. Examples of mutations can be found in Examples 6.2 and 6.3.

6.1.4 Product of valued quivers

From the skew-symmetrizable, resp. skew-symmetric, companion matrices corresponding to the Dynkin type C_2 , resp. A_w , consider the corresponding valued quivers as follows

$$C_2: 1 \xrightarrow{(1,2)} 2 \qquad A_w: 1 \longrightarrow 2 \longrightarrow \dots \longrightarrow w-1 \longrightarrow w.$$

Following [20], we define the square product of the valued quivers:


$$C_2 \square A_w := \begin{array}{ccccccc}
 \bullet_{w+1} & \xrightarrow{\quad} & \circ_{w+2} & \xleftarrow{\quad} & \bullet_{w+3} & \xrightarrow{\quad} & \dots \\
 \uparrow (1,2) & & \downarrow (1,2) & & \uparrow (1,2) & & \dots \\
 \circ_1 & \xleftarrow{\quad} & \bullet_2 & \xrightarrow{\quad} & \circ_3 & \xleftarrow{\quad} & \dots \\
 & & & & & & \dots \xrightarrow{\quad} \circ_{w-1} \xleftarrow{\quad} \bullet_w \\
 & & & & & & \uparrow (1,2) \quad \downarrow (1,2)
 \end{array} \tag{6.1}$$

It is the valued quiver associated to the following $2w \times 2w$ skew-symmetrizable matrix


$$B_{C_2 \square A_w} = \left(\begin{array}{cccc|cccc}
 0 & -1 & 0 & 0 & \dots & 1 & 0 & 0 & 0 & \dots \\
 1 & 0 & 1 & 0 & \dots & 0 & -1 & 0 & 0 & \dots \\
 0 & -1 & 0 & -1 & \dots & 0 & 0 & 1 & 0 & \dots \\
 \vdots & & & & \vdots & \vdots & & & \vdots & \\
 -2 & 0 & 0 & 0 & \dots & 0 & 1 & 0 & 0 & \dots \\
 0 & 2 & 0 & 0 & \dots & -1 & 0 & -1 & 0 & \dots \\
 0 & 0 & -2 & 0 & \dots & 0 & 1 & 0 & 1 & \dots \\
 \vdots & & & & \vdots & \vdots & & & \vdots &
 \end{array} \right)$$

Example 6.2. (a) For $w = 1$, $C_2 \square A_1$ is just the valued quiver $C_2 : 1 \xrightarrow{(1,2)} 2$ associated to the matrix $\begin{pmatrix} 0 & 1 \\ -2 & 0 \end{pmatrix}$.

(b) For $w = 2$, one can show that $C_2 \square A_2$ is mutation equivalent to a valued quiver of Dynkin type F_4 . Indeed,




Example 6.3. Let $\Sigma = ((x_1, x_2, \dots, x_{2w}), C_2 \square A_w)$ be a seed. A mutation at vertex i for $1 \leq i \leq w$ in (6.1) locally changes the graph around the vertices i and $w + i$ into



if i is even (and a similar picture if i is odd but with reversed arrows), and changes the variable x_i to

$$x'_i = \frac{x_{i-1}x_{i+1} + x_{w+i}^2}{x_i}. \quad (6.2)$$

A mutation at vertex $w + i$ for $1 \leq i \leq w$ in (6.1) locally changes the graph around the vertices i and $w + i$ into



if i is even (and a similar picture if i is odd but with reversed arrows), and changes the variable x_{w+i} to

$$x'_{w+i} = \frac{x_{w+i-1}x_{w+i+1} + x_i}{x_{w+i}}. \quad (6.3)$$

6.1.5 Bipartite belt

A *bipartite* quiver is such that the vertices can be colored in two colors so that vertices of the same color are not connected by any arrow. For instance, the quiver (6.1) is bipartite.

Let \mathcal{Q} be a bipartite quiver. Following Fomin–Zelevinsky [15], consider the iterated mutations

$$\mu_+ = \prod_{\text{white } k} \mu_k, \quad \mu_- = \prod_{\text{black } k} \mu_k.$$

Note that the mutation μ_k 's with k of a fixed color commute with each other since the vertices are not connected.

The mutations μ_+ and μ_- on Q give the same quiver with reversed orientation:

$$\mu_+(\mathcal{Q}) = \mathcal{Q}^{\text{op}}, \quad \mu_-(\mathcal{Q}^{\text{op}}) = \mathcal{Q}.$$

Let $\Sigma_0 = (\underline{x}, \mathcal{Q})$ be an initial seed. The *bipartite belt*, see [15], of the cluster algebra \mathcal{A}_{Σ_0} is the collection of seeds obtained from Σ_0 by applying successively μ_+ or μ_- :

$$\dots, \mu_-\mu_+\mu_-(\Sigma_0), \mu_+\mu_-(\Sigma_0), \mu_-(\Sigma_0), \Sigma_0, \mu_+(\Sigma_0), \mu_-\mu_+(\Sigma_0), \mu_+\mu_-\mu_+(\Sigma_0), \dots$$

Remark 6.4. In general, the bipartite belt is infinite. It has been proved, see [19, 20], that for a bipartite (valued) quiver constructed as a product of two Dynkin quivers $\Delta \square \Delta'$ the mutation $\mu_+\mu_-$ has finite order, more precisely

$$(\mu_+\mu_-)^{2(h+h')} = (\mu_-\mu_+)^{2(h+h')} = \text{Id},$$

where h and h' are the Coxeter numbers of the Dynkin quivers Δ, Δ' respectively. This property is known as *Zamolodchikov periodicity*.

6.2 Cluster variables in the symplectic 2-friezes

Let F_w be the tame symplectic 2-frieze of width w of rational fractions in the free variables $(x_1, x_2, \dots, x_{2w})$ computed with the frieze rule from two consecutive columns containing the variables $(x_1, \dots, x_w, \mathbf{x}_{w+1}, \dots, \mathbf{x}_{2w})$ in a zig-zag as follows.

$$F_w: \begin{array}{cccccccc} \dots & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ & & x_1 & \mathbf{x}_{w+1} & x'_1 & \mathbf{x}'_{w+1} & & \\ & & \mathbf{x}_{w+2} & x_2 & \mathbf{x}'_{w+2} & x'_2 & & \\ & & x_3 & \mathbf{x}_{w+3} & x'_3 & \mathbf{x}'_{w+3} & & \\ & & \vdots & \vdots & \vdots & \vdots & & \\ & & \mathbf{x}_{2w} & x_w & \mathbf{x}'_{2w} & x'_w & & \\ \dots & 1 & 1 & 1 & 1 & 1 & 1 & \dots \end{array} \quad (6.4)$$

The frieze (3.5) is exactly the frieze F_w for $w = 1$.

Remark 6.5. The frieze F_w is well defined. The operations used to compute the entries in the symplectic 2-frieze in terms of the initial entries $(x_1, x_2, \dots, x_{2w})$ are subtraction free. So that the rational expressions that one computes for the next entries $(x'_1, x'_2, \dots, x'_{2w})$ do not vanish. This allows to compute recursively all the entries of the frieze as elements of $\mathbb{C}(x_1, \dots, x_{2w})$. Moreover, since the entries are non zero, the frieze F_w is tame by Proposition 4.1.

Proposition 6.6. *Let $\mathcal{A} = \mathcal{A}_{\Sigma_0}$ be the cluster algebra generated from the initial seed $\Sigma_0 = ((x_1, x_2, \dots, x_{2w}), \mathcal{Q})$, where $\mathcal{Q} = \text{C}_2 \square \text{A}_w$ is the valued quiver (6.1). Let F_w be the formal frieze (6.4) containing the same initial variables $(x_1, x_2, \dots, x_{2w})$.*

- (i) *The entries of F_w are all cluster variables of \mathcal{A} ;*
- (ii) *The pairs of consecutive columns of F_w give all the clusters of the bipartite belt of \mathcal{A} .*

Proof. (i) is a trivial consequence of (ii). Statement (ii) is established by direct computations. Indeed, let $(x'_1, x'_{w+2}, x'_3, \dots)$ and $(x'_{w+1}, x'_2, x'_{w+3}, \dots)$ be the third and fourth column in F_w . The entries are deduced using the frieze rule so that we have for i odd, $1 \leq i \leq w$,

$$x'_i = \frac{x_{i-1}x_{i+1} + x_{w+i}^2}{x_i},$$

and i even, $1 \leq i \leq w$,

$$x'_{w+i} = \frac{x_{w+i-1}x_{w+i+1} + x_i}{x_{w+i}}.$$

Comparing with the result of the mutation μ_+ , see (6.2), (6.3), one immediately gets

$$\begin{aligned} (x'_1, x_2, x'_3, \dots, x'_{2w-1}, x_{2w}, \mathcal{Q}^{\text{op}}) &= \mu_+ \Sigma_0, \\ (x'_1, x'_2, x'_3, \dots, x'_{2w-1}, x'_{2w}, \mathcal{Q}) &= \mu_- \mu_+ \Sigma_0, \end{aligned}$$

and so on. ■

Remark 6.7. From the cluster algebra theory Theorem 2.1 becomes an easy consequence of Proposition 6.6 and the Zamolodchikov periodicity, see Remark 6.4 (recall that the Coxeter number for C_2 is 4 and the one for A_w is $w + 1$). However, the periodicity of the 2-friezes or the periodicity of the SL-friezes can be established with elementary arguments using the identification with superperiodic difference equations, see [24].

6.3 More clusters in the 2-friezes

From F_w one can extract more clusters of \mathcal{A} than the ones of the bipartite belt. These clusters are given as “double zig-zags” in the array. We follow [25].

We call *double zig-zag* in the 2-frieze (6.4) a sequence $\zeta = (\tilde{x}_1, \dots, \tilde{x}_w, \tilde{y}_1, \dots, \tilde{y}_w)$ where \tilde{x}_i 's are white entries and \tilde{y}_i 's black entries that are locally in the following 6 configurations


$$\begin{array}{ccc} \begin{array}{c} \tilde{x}_i \quad \tilde{y}_i \\ \diagdown \quad \diagup \\ \tilde{x}_{i+1} \quad \tilde{y}_{i+1} \end{array} & \begin{array}{c} \tilde{x}_i \quad \tilde{y}_i \\ \diagup \quad \diagdown \\ \tilde{y}_{i+1} \quad \tilde{x}_{i+1} \end{array} & \begin{array}{c} \tilde{x}_i \quad \tilde{y}_i \\ \diagdown \quad \diagdown \\ \tilde{x}_{i+1} \quad \tilde{y}_{i+1} \end{array} \\ \begin{array}{c} \tilde{y}_i \quad \tilde{x}_i \\ \diagup \quad \diagdown \\ \tilde{y}_{i+1} \quad \tilde{x}_{i+1} \end{array} & \begin{array}{c} \tilde{y}_i \quad \tilde{x}_i \\ \diagdown \quad \diagup \\ \tilde{x}_{i+1} \quad \tilde{y}_{i+1} \end{array} & \begin{array}{c} \tilde{y}_i \quad \tilde{x}_i \\ \diagdown \quad \diagdown \\ \tilde{y}_{i+1} \quad \tilde{x}_{i+1} \end{array} \end{array} \tag{6.5}$$

Example 6.8. The following sequence forms a double zig-zag in a frieze of width 5:

$$\begin{array}{cccccccc} \dots & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ & & \tilde{x}_1 & \tilde{y}_1 & & & & \\ & & & \tilde{x}_2 & \tilde{y}_2 & & & \\ & & \tilde{x}_3 & \tilde{y}_3 & & & & \\ & & \tilde{y}_4 & \tilde{x}_4 & & & & \\ & & \tilde{y}_5 & \tilde{x}_5 & & & & \\ \dots & 1 & 1 & 1 & 1 & 1 & 1 & \dots \end{array} \tag{6.6}$$


To every double zig-zag ζ one associates a valued quiver \mathcal{Q}_ζ defined over the set of vertices $\{1, 2, \dots, 2w\}$ as follows. Each of the 6 configurations of (6.5) gives respectively the following

set of weighted arrows:



when i is odd, and same pictures with reversed arrows when i is even.

Example 6.9. The quiver associated to the double zig-zag of (6.6) is



One can extend Proposition 6.6 with the following statement.

Proposition 6.10. For every double zig-zag ζ in F_w , the couple $(\zeta, \mathcal{Q}_\zeta)$ is a seed of \mathcal{A}_{Σ_0} .

Proof. The statement follows from the mutation rule and the frieze rule. Every double zig-zag can be redressed in two consecutive columns by elementary moves that correspond to mutations. ■

6.4 The cluster variety of tame symplectic 2-friezes

Let $\mathcal{A} = \mathcal{A}_{\Sigma_0}$ be the cluster algebra generated from the initial seed $\Sigma_0 = ((x_1, x_2, \dots, x_{2w}), \mathbb{C}_2 \times A_w)$, and let $\chi = (u_1, \dots, u_{2w})$ be an arbitrary cluster of \mathcal{A} . We know from the cluster algebra theory that all the cluster variables of \mathcal{A} can be expressed as Laurent polynomials with positive integer coefficients in the variables u_i . Denote by $F_w(u_1, \dots, u_{2w})$ the frieze of Laurent polynomials obtained by expressing all the entries of F_w in terms of u_i . When specializing the variables u_i to non-zero complex numbers one gets a tame symplectic 2-frieze over complex numbers. This construction can be summarized with the following statement.

Corollary 6.11. Every cluster $\chi = (u_1, \dots, u_{2w})$ of \mathcal{A} , defines an injective map

$$\phi_\chi: (z_1, \dots, z_{2w}) \mapsto F_w(u_1, \dots, u_{2w})|_{u_i=z_i}$$

from $(\mathbb{C}^*)^{2w}$ to the variety of tame symplectic 2-friezes.

Obviously the maps ϕ_χ are not surjective. A natural question is the following.

Open problem 6.12. Is the variety of tame symplectic 2-friezes of width w equal to the union over all clusters of the images of the maps ϕ_χ ?

Let us mention that the similar question in the case of tame Coxeter's friezes has been addressed and answered by Cuntz and Holm [10] (in that case the answer is positive when the width is not equal to 3 modulo 4, otherwise there exists an extra tame frieze).

Definition 6.13. A tame symplectic 2-frieze is called *cluster-singular* if it is not in any of the images of the maps ϕ_χ , otherwise it is called *cluster-regular*

For $w = 1, 2$, one can give the following (partial) answers.

Proposition 6.14.

(i) *There is only one cluster-singular symplectic 2-frieze of width 1 given by*

$$\begin{array}{cccccccccccc} \dots & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & \dots \\ \dots & -1 & \mathbf{0} & -1 & \mathbf{0} & -1 & \mathbf{0} & -1 & \mathbf{0} & -1 & \dots \\ \dots & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & \dots \end{array} \quad (6.7)$$

(ii) *There is at most one cluster-singular tame symplectic 2-frieze of width 2.*

Proof. For $w = 1$, a tame symplectic 2-frieze is entirely determined by two consecutive non-zero entries. Indeed, a tame symplectic 2-frieze of width 1 is necessarily 6-periodic and two consecutive non-zero entries allow to obtain 6 consecutive entries using the frieze rule. The cluster algebra of type C_2 has exactly 6 clusters given as two consecutive entries in the single non trivial row of the array (3.5). There are only two ways to avoid two consecutive non zero entries in a symplectic 2-frieze of width 2, leading to (6.7) and to

$$\begin{array}{cccccccccccc} \dots & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & \dots \\ \dots & 0 & \mathbf{i} & 0 & -\mathbf{i} & 0 & \mathbf{i} & 0 & -\mathbf{i} & 0 & \mathbf{i} & \dots \\ \dots & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & \mathbf{1} & \dots \end{array} \quad (6.8)$$

One checks that the frieze (6.7) is tame. To do so one uses Remark 4.3 and check (4.1) and (4.3) centered at a black zero entry. The only relevant calculations are to

$$\begin{array}{l} \left| \begin{array}{ccc} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{array} \right| = 0, \\ \left| \begin{array}{ccccc} 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{array} \right| = 0, \end{array}$$

which allow to conclude that (6.7) is tame. Similarly for (6.8) one computes

$$\left| \begin{array}{cccc} i & 1 & 0 & 0 \\ 1 & -i & 1 & 0 \\ 0 & 1 & i & 1 \\ 0 & 0 & 1 & -i \end{array} \right| = -1,$$

which shows that (6.8) is not tame since (4.2) fails. Hence (i).

For $w = 2$, we start by establishing two lemmas covering the cases when the frieze has four adjacent non-zero entries or two consecutive zero entries (we believe that the first lemma should also be true for an arbitrary width and arbitrary double zig-zag).

Lemma 6.15. *Four non-zero entries in two consecutive columns, or in two consecutive diagonals, determine a unique tame symplectic 2-frieze of width 2 over the complex numbers. This symplectic 2-frieze is cluster-regular.*

Proof. Since a tame symplectic 2-frieze is invariant under a glide symmetry it has the following general form:

$$\begin{array}{cccccccccccccccc}
\dots & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\
\dots & A & b & C & d & E & f & G & h & I & j & K & l & M & n & A & \dots \\
\dots & h & I & j & K & l & M & n & A & b & C & d & E & f & G & h & \dots \\
\dots & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots
\end{array} \tag{6.9}$$

where the upper case letters are black entries and lower case letters white entries. Let us assume that the four entries C, d, j, K are non zero complex numbers. We know that there is at least one tame symplectic 2-frieze with these particular entries C, d, j, K in two consecutive columns. This is the frieze obtained by specializing the cluster $\chi = (x'_1, x_2, \mathbf{x}_3, \mathbf{x}'_4)$ to (d, j, C, K) in (6.4) for $w = 2$.

Let us show that there is at most one tame symplectic 2-frieze with these particular entries C, d, j, K in two consecutive columns. The frieze rule uniquely determines the entries A, b, E, f, h, I, l, M of the frieze. The entries G and n are possibly undefined by the frieze rule (this happens in the case where $E = I = b = l = 0$). However the tameness condition (4.1) expressed in the 3×3 minor centered at E implies that G is uniquely determined (one has $Gd - CM + 1 = E$ with $d \neq 0$). Then $n = MA - G$ is also uniquely determined. ■

Lemma 6.16. *Tame symplectic 2-friezes of width 2 over \mathbb{C} containing two consecutive zero entries in at least one of its rows are exactly of the following form*

$$\begin{array}{cccccccccccccccc}
\dots & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\
\dots & A & -1 & 0 & 0 & 1 & f & G & h & 1 & 0 & 0 & -1 & M & \dots \\
\dots & h & 1 & 0 & 0 & -1 & M & n & A & -1 & 0 & 0 & 1 & f & \dots \\
\dots & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots
\end{array} \tag{6.10}$$

with $G = f + M$, $h = f + 2M - 1$, $n = -f - M^2$, $A = 1 - M$, for some free parameters $f, M \in \mathbb{C}$.

Proof. Let us consider the general tame symplectic 2-frieze (6.9). Assume $C = d = 0$. The frieze rule immediately implies $j = K = 0$ and then $E^2 = -l = E$. So one immediately gets two possible values for E , namely $E = 0$ or $E = 1$. The tameness condition (4.1) expressed in the 3×3 -minor centered at C implies $IE - 1 = 0$. Therefore $E \neq 0$ so that $E = 1$ and $l = -1, I = 1$. Using the frieze rule one deduces $b = -1, G = f + M, n = -f - M^2$. Finally the tameness condition (4.1) expressed in the 3×3 minor centered at M gives $A = 1 - M$ from which one deduces $h = f + 2M - 1$ with the frieze rule. We have established that a tame symplectic 2-frieze of width 2 containing two consecutive zero entries are necessarily of the form (6.10). Let us show that the symplectic 2-frieze (6.10) is tame for any values of the parameters f and M . When $fM \neq 0$ the frieze (6.10) is exactly the one obtained by specializing the cluster $\chi = (x'_1, x_2, \mathbf{x}_3, \mathbf{x}'_4)$ to $(f, -1, 1, M)$ in (6.4) for $w = 2$. This means that the polynomial identities (4.1), (4.2), (4.3), required for the tameness of the frieze, hold for any non-zero values of f and M , and therefore for any values of f and M . ■

We go back to the proof of Proposition 6.14(ii). We are looking for cluster-singular symplectic 2-friezes of width 2. Let us consider a tame symplectic 2-frieze of width 2.

Case (1): there are two consecutive zeroes on the first (or last) row of the frieze. By Lemma 6.16 the array is of the form (6.10). If $(f, M) \neq (0, \frac{1}{2})$ the frieze always contains four adjacent non-zero entries, so that it is cluster-regular according to Lemma 6.15.

Case (2): the first (or last) row has a sequence of the form $(x, 0, y, 0, z, 0, t)$ where $xyzt \neq 0$ are either black or white entries. The frieze rule will imply that the array is of the form

$$\begin{array}{cccccccc} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ x & 0 & y & 0 & z & 0 & t & r \\ r & xy & s & yz & u & zt & v & \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \end{array}$$

with $rsuv \neq 0$ so that the specified entries in the second row are non zero. By glide symmetry, one would have the entry r next to t in the first row so that (t, zt, r, v) are non zero entries in two consecutive diagonals. So the frieze is cluster-regular according to Lemma 6.15.

Case (3): the first row has two consecutive non zero entries (x, y) . Let us denote by (s, t) the entries under (x, y) . If $st \neq 0$ then one has two consecutive columns of non-zero entries and we are done by Lemma 6.15. By the frieze rule one has $(s, t) \neq (0, 0)$. We can now assume that the array is locally of the form

$$\begin{array}{cccc} 1 & 1 & 1 & 1 \\ x & y & z & \\ 0 & t & u & v \\ 1 & 1 & 1 & 1 \end{array}$$

with $t \neq 0$. If $u \neq 0$ then (x, t, y, u) are non zero entries in two consecutive diagonals and we are done by Lemma 6.15. If $u = 0$ then $v \neq 0$ (otherwise it goes back to Case (1)) and therefore $z = tv \neq 0$. By induction we show that the next entries on the first row are non zero and the one of the second row alternate zero and non-zero, so that it goes back to Case (2). ■

Remark 6.17. To give a complete answer to the open problem 6.12 it remains to determine whether the following tame symplectic 2-frieze is cluster-regular or not:

$$\begin{array}{cccccccccccccccc} \dots & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ \dots & -1 & 0 & 0 & 1 & 0 & \frac{1}{2} & 0 & 1 & 0 & 0 & -1 & \frac{1}{2} & \dots \\ \dots & 1 & 0 & 0 & -1 & \frac{1}{2} & -\frac{1}{4} & \frac{1}{2} & -1 & 0 & 0 & 1 & 0 & \dots \\ \dots & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \end{array}$$

(this is the frieze (6.10) for $(f, M) = (0, \frac{1}{2})$).

Remark 6.18. The frieze (6.7) is the point of coordinates $(a_1, \dots, a_6) = (0, \dots, 0)$, $(b_1, \dots, b_6) = (-1, \dots, -1)$ in the variety described in Example 3.5. The tangent space at this point is of dimension 2 so that it is not a singular point of the variety (the variety is actually smooth).

Conjecture 6.19. *The frieze of width 7 given in (4.5) is cluster-singular.*

7 Further investigations

7.1 Symplectic 2-friezes of positive integers

A major problem in the theory of friezes is the classification of the friezes with positive integers. This problem usually leads to very nice combinatorial interpretations. The main result in the domain is the theorem of Conway–Coxeter [7] that classifies Coxeter friezes with triangulations of polygons. There are many results along the same line for the classification of other types of friezes, e.g., [3, 5, 16, 27].

There is also a nice extension of the Conway–Coxeter theorem in terms of 3d-dissections of polygons by allowing negative entries in the Coxeter friezes [26].

Open problem 7.1. *Find a combinatorial model for the classification of the symplectic 2-friezes of positive integers.*

Note that it is still an open problem in the case of classical 2-friezes and SL-friezes.

An “easy” way to produce symplectic 2-friezes of positive integers is to use the maps ϕ_χ given in Proposition 6.11 and evaluate on the point $(z_1, \dots, z_{2w}) = (1, \dots, 1)$. It was proved in [22] that when the cluster algebra \mathcal{A} is of infinite type, i.e., has infinitely many clusters, this produces infinitely many friezes of positive integers. When the cluster algebra \mathcal{A} is of finite type this procedure is in general not sufficient to produce all the friezes with positive integers (except in the case of Coxeter friezes).

Proposition 7.2. *There exist exactly 6 symplectic 2-friezes of width 1 with positive integers.*

Proof. The formal frieze F_1 , cf. (6.4) contains the 6 cluster variables of type C_2 . It is known that in this case there is only 6 ways to get positive integers simultaneously for all the cluster variables [16]. ■

The 6 friezes are all obtained from (3.2) under the action of the dihedral group D_6 .

Conjecture 7.3. *There exist exactly 112 symplectic 2-friezes of width 2 with positive integers.*

This conjecture is based on two evidences. First, Michael Cuntz kindly provide me with the 112 arrays of SL_3 -friezes which are symmetric with respect to the middle line. By Theorem 2.2 they correspond to symplectic 2-friezes of width 2. The 112 SL_3 -friezes are grouped in 9 arrays modulo the action of the dihedral group D_7 . The second evidence is that the entries of the symplectic 2-friezes of width 2 are cluster variables of type F_4 (see Example 6.2) and it has been already conjectured in [16] that there are only 112 evaluations that make all the cluster variables positive integers. Note that in this case only 14 out of 28 cluster variables appear in the frieze, so that it was not necessary to have all the cluster variables simultaneously positive integers. Note also that there are 105 clusters of type F_4 . Which would mean that 7 friezes are not obtained as $\phi_\chi(1, \dots, 1)$.

Bernhard Keller mentioned to me that the applet [18] allows to check that the cluster algebra of type $C_2 \square A_w$ is cluster-infinite whenever $w \geq 3$. By consequence, there exist infinitely many symplectic 2-friezes of positive integers of width w whenever $w \geq 3$.

7.2 The 2-friezes of type G_2

Replacing the square by a cube in the local rule of the symplectic 2-frieze, i.e., when the local rule in the array (2.1) reads

$$\mathbf{AD} - \mathbf{BC} = e, \quad eh - fg = \mathbf{D}^3,$$

leads to interesting arrays. In particular they are still periodic, of period $2(m+7)$ where m is the width, and are invariant under a glide symmetry. They are related to the combinatorics of the cluster algebras of type $G_2 \times A_m$. Such friezes could be called “2-friezes of type G_2 ” and it would be interesting to study them further from a combinatorial viewpoint and geometric viewpoint.

Open problem 7.4. *What are the configurations spaces related to the 2-friezes of type G_2 ?*

We give below examples of 2-friezes of type G_2 of width 1 and 2.

$$\begin{array}{cccccccccccccccc} \dots & 1 & \mathbf{1} & 1 & 1 & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & 1 & \mathbf{1} & 1 & 1 & \mathbf{1} & 1 & \dots \\ \dots & \mathbf{1} & 2 & \mathbf{3} & 14 & \mathbf{5} & 9 & \mathbf{2} & 1 & \mathbf{1} & 2 & \mathbf{3} & 14 & \mathbf{5} & 9 & \mathbf{2} & 1 & \dots \\ \dots & 1 & \mathbf{1} & 1 & 1 & 1 & \mathbf{1} & 1 & \mathbf{1} & 1 & 1 & \mathbf{1} & 1 & 1 & \mathbf{1} & 1 & \dots \end{array}$$

$$\begin{array}{cccccccccccccccccccc}
 \dots & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\
 \dots & 1 & 2 & 3 & 15 & 7 & 28 & 6 & 9 & 2 & 1 & 1 & 3 & 6 & 77 & 14 & 36 & 3 & 1 & \dots \\
 \dots & 1 & 1 & 3 & 6 & 77 & 14 & 36 & 3 & 1 & 1 & 2 & 3 & 15 & 7 & 28 & 6 & 9 & 2 & \dots \\
 \dots & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots
 \end{array}$$

7.3 Symplectic friezes in arbitrary dimensions

In [6], the spaces of Lagrangian configurations of n lines in \mathbb{C}^{2k} modulo Sp_{2k} are studied. These configurations should give rise to more general symplectic friezes (recall that the case of symplectic 2-frieze is when $k = 2$) and be connected to the cluster combinatorics of type $C_k \times A_m$, where $n = 2k + m + 1$. In particular the moduli space of Lagrangian configurations of $2k + 2$ lines in \mathbb{C}^{2k} should be a cluster variety of type C_k .

Open problem 7.5. *What are the friezes related to Lagrangian configurations in arbitrary dimensions?*

A Desnanot–Jacobi identity

The Desnanot–Jacobi identity, or Dodgson formula, is a classical formula involving the determinant of an $(n \times n)$ -matrix and its minors of order $n - 1$ and $n - 2$ obtained by erasing the first and/or last row/column. The formula can be pictured as follows

$$\left| \begin{array}{cccc} * & * & * & * \\ * & * & * & * \\ * & * & * & * \\ * & * & * & * \end{array} \right| \left| \begin{array}{cc} * & * \\ * & * \end{array} \right| = \left| \begin{array}{ccc} * & * & * \\ * & * & * \\ * & * & * \end{array} \right| \left| \begin{array}{ccc} * & * & * \\ * & * & * \\ * & * & * \end{array} \right| - \left| \begin{array}{ccc} * & * & * \\ * & * & * \\ * & * & * \end{array} \right| \left| \begin{array}{ccc} * & * & * \\ * & * & * \\ * & * & * \end{array} \right|,$$

where the deleted columns/rows are left blank.

It is a key identity to establish Proposition 4.1.

B Properties of SL_{k+1} -friezes

B.1 Definition of tame SL_{k+1} -friezes

An SL_{k+1} -frieze is an array of numbers consisting in a finite number of infinite rows:

$$\begin{array}{cccccccc}
 & & & \vdots & & \vdots & & \\
 & 0 & & 0 & & 0 & & 0 & \dots \\
 \dots & & 1 & & 1 & & 1 & & 1 & & 1 & & \dots \\
 & \dots & & d_{0,w-1} & & d_{1,w} & & d_{2,w+1} & & \dots & & \dots \\
 & & & \ddots & & \ddots & & \ddots & & & & & \\
 \dots & & d_{0,1} & & d_{1,2} & & d_{2,3} & & d_{3,4} & & d_{4,5} & & \\
 & d_{0,0} & & d_{1,1} & & d_{2,2} & & d_{3,3} & & d_{4,4} & & \dots \\
 \dots & & 1 & & 1 & & 1 & & 1 & & 1 & & \dots \\
 & 0 & & 0 & & 0 & & 0 & & 0 & & \dots \\
 & & & \vdots & & \vdots & & & & & & &
 \end{array} \tag{B.1}$$

where the strip is bounded by k rows of 0's at the top, and at the bottom, and where every “diamond” $(k + 1) \times (k + 1)$ -subarray forms an element of SL_{k+1} . The number of rows between the bounding rows of 1's is called the width and is denoted by w .

More precisely, the entries in the array are denoted by $(d_{i,j})$, with $i, j \in \mathbb{Z}$ such that

$$i - k - 1 \leq j \leq i + w + k.$$

The array is a tame SL_{k+1} -frieze of width w when it satisfies:

- “boundary conditions”

$$\begin{cases} d_{i,i-1} = d_{i,i+w} = 1 & \text{for all } i, \\ d_{i,j} = 0 & \text{for } i - k - 1 \leq j < i - 1 \text{ or } i + w < j \leq i + w + k. \end{cases}$$

- “ SL_{k+1} -conditions” on the $(k+1) \times (k+1)$ -adjacent minors

$$\begin{vmatrix} d_{i,j} & d_{i,j+1} & \cdots & d_{i,j+k} \\ d_{i+1,j} & d_{i+1,j+1} & \cdots & d_{i+1,j+k} \\ \cdots & \cdots & \cdots & \cdots \\ d_{i+k,j} & d_{i+k,j+1} & \cdots & d_{i+k,j+k} \end{vmatrix} = 1,$$

for all (i, j) in the index set.

- “tameness conditions” on the $(k+2) \times (k+2)$ -adjacent minors

$$\begin{vmatrix} d_{i,j} & d_{i,j+1} & \cdots & d_{i,j+k+1} \\ d_{i+1,j} & d_{i+1,j+1} & \cdots & d_{i+1,j+k+1} \\ \cdots & \cdots & \cdots & \cdots \\ d_{i+k+1,j} & d_{i+k+1,j+1} & \cdots & d_{i+k+1,j+k+1} \end{vmatrix} = 0,$$

for all (i, j) in the index set.

B.2 Tame SL_{k+1} -friezes and difference equations

An n -superperiodic difference equation of order $k+1$ is a system

$$V_i = a_i^1 V_{i-1} - a_i^2 V_{i-2} + \cdots + (-1)^{k-1} a_i^k V_{i-k} + (-1)^k V_{i-k-1}, \quad (\text{B.2})$$

where $a_i^j \in \mathbb{C}$, with $i \in \mathbb{Z}$ and $1 \leq j \leq k$, are given coefficients (note that the superscript j is an index, not a power) and V_i are unknowns, such that

- the coefficients are n -periodic, i.e., $a_{i+n}^j = a_i^j$, for all i, j .
- every solution $(V_i)_{i \in \mathbb{Z}}$ is n -(anti)periodic, i.e., $V_{i+n} = (-1)^k V_i$, for all $i \in \mathbb{Z}$.

Proposition B.1 ([24]). *An array $(d_{i,j})$ as (B.1) forms a tame SL_{k+1} frieze if and only if every diagonal*

$$d_{i,\bullet} = (0, \dots, 0, 1, d_{i,i}, d_{i,i+1}, \dots, d_{i,i+w-1}, 1)$$

gives a solution of a same $(w+k+2)$ -superperiodic equation of the form (B.2) with initial conditions

$$(V_{i-k-1}, V_{i-k}, \dots, V_{i-1}) = (0, 0, \dots, 0, 1).$$

B.3 Coefficients of the difference equations from the entries of the frieze and *vice versa*

We collect some formulas given in [24, Sections 5.2 and 5.3].

Let (B.2) be the associated equation associated to a tame SL_{k+1} -frieze (B.1). The coefficient a_{i-1}^{k-j} of the equation can be expressed as a $(j+1) \times (j+1)$ -minor of adjacent entries in the frieze:

$$a_{i-1}^{k-j} = \begin{vmatrix} d_{i+1,i+w} & 1 & & & \\ \vdots & \ddots & & & \\ d_{i+j+1,i+w} & \cdots & d_{i+j+1,i+j+w} & & \end{vmatrix}. \quad (\text{B.3})$$

Conversely, the entries of the frieze can be computed as determinant of matrices involving the coefficients of the equation. The entry $d_{i,i+j}$ can be computed using $(j+1) \times (j+1)$ -determinants

$$d_{i,i+j} = \begin{vmatrix} a_i^1 & 1 & & & & & \\ \vdots & a_{i+1}^1 & 1 & & & & \\ a_{i+k-1}^k & & \ddots & \ddots & & & \\ 1 & & & \ddots & \ddots & & \\ & \ddots & & & a_{i+j-1}^1 & 1 & \\ & & 1 & a_{i+j}^k & \cdots & a_{i+j}^1 & \end{vmatrix}, \quad (\text{B.4})$$

or alternatively, using $(w-j) \times (w-j)$ -determinants

$$d_{i,i+j} = \begin{vmatrix} a_{i-w+j-1}^k & \cdots & a_{i-w+j-1}^1 & 1 & & & \\ 1 & a_{i-w+j}^k & \cdots & a_{i-w+j}^1 & 1 & & \\ & 1 & \cdots & & \ddots & 1 & \\ & & \ddots & & \ddots & \vdots & \\ & & & 1 & a_{i-3}^k & a_{i-3}^{k-1} & \\ & & & & 1 & a_{i-2}^k & \end{vmatrix}. \quad (\text{B.5})$$

B.4 Projective duality

Given a tame SL_{k+1} -frieze $(d_{i,j})$, its projective dual is defined as the array consisting of $k \times k$ adjacent minors:

$$d_{i,j}^* = \begin{vmatrix} d_{i,j} & d_{i,j+1} & \cdots & d_{i,j+k-1} \\ d_{i+1,j} & d_{i+1,j+1} & \cdots & d_{i+1,j+k-1} \\ \vdots & & & \vdots \\ d_{i+k-1,j} & d_{i+k-1,j+1} & \cdots & d_{i+k-1,j+k-1} \end{vmatrix}.$$

Proposition B.2 ([24]).

- (i) The projective dual frieze to a tame SL_{k+1} -frieze is a tame SL_{k+1} -frieze.
- (ii) If a_i^j are the coefficients of the equation (B.2) associated to a SL_{k+1} -frieze, then

$$V_i^* = a_{i+k-1}^k V_{i-1}^* - a_{i+k-2}^{k-1} V_{i-2}^* + \cdots + (-1)^{k-1} a_i^1 V_{i-k}^* + (-1)^k V_{i-k-1}^* \quad (\text{B.6})$$

is the superperiodic equation associated to the projective dual.

(iii) *The projective dual to a tame SL_{k+1} -frieze is just the symmetric array with respect to the median horizontal axis.*

Let us make the above statement number (iii) more precise.

Proposition B.3. *A tame SL_{k+1} -frieze of width w satisfies*

$$d_{i,j} = \begin{vmatrix} d_{j-w-k,i-k-1} & \cdots & d_{j-w-k,i-2} \\ \vdots & & \vdots \\ d_{j-w-1,i-k-1} & \cdots & d_{j-w-1,i-2} \end{vmatrix} = d_{j-w-k,i-k-1}^* \quad (\text{B.7})$$

for all i, j .

Proof. First, note the diagonals

$$d_{\bullet,i} = (0, \dots, 0, 1, d_{i,i}, d_{i-1,i}, \dots, d_{i-w+1,i}, 1)$$

satisfy the dual equation (B.6). Let us consider two sequences $V = (V_\ell)_{1 \leq \ell \leq j-i+k+2}$ and $W = (W_\ell)_{1 \leq \ell \leq j-i+k+2}$ forming blocks of size $(k+1) \times (j-i+k+2)$ in the frieze that differ by a shift along the north-west diagonal:

$$\begin{array}{cccccccc} & V_1 & & V_2 & & \cdots & V_{k+1} & & V_{k+2} & & \cdots & V_{j-i+1} & & V_{j-i+k+2} \\ \left(\begin{array}{cccccccc} d_{j-w-k,i-k-1} & d_{j-w-k,i-k} & \cdots & \cdots & d_{j-w-k,i} & \cdots & 1 & 0 & 0 \\ \vdots & \vdots & & & \vdots & \cdots & & 1 & \\ \vdots & \vdots & & & \vdots & \cdots & & \ddots & 0 \\ d_{j-w,i-k-1} & d_{j-w,i-k} & \cdots & \cdots & d_{j-w,i} & \cdots & \cdots & \cdots & 1 \\ \vdots & \vdots & & & \vdots & & & & \vdots \\ 1 & d_{i-k,i-k} & \cdots & \cdots & d_{i-k,i} & \cdots & \cdots & \cdots & d_{i-k,j} \\ 0 & 1 & \ddots & & \vdots & & & & \vdots \\ \vdots & \vdots & \ddots & \ddots & \vdots & & & & \vdots \\ 0 & 0 & \cdots & 1 & d_{i,i} & \cdots & \cdots & \cdots & d_{i,j} \end{array} \right) = V, \\ & W_1 & & W_2 & & \cdots & W_{k+1} & & W_{k+2} & & \cdots & W_{j-i+1} & & W_{j-i+k+2} \end{array} \right) = W.$$

To simplify the picture we have rotated the frieze by 45 degree so that the shift between the blocks is now seen vertically.

Due to the recurrence relation in the frieze, the matrix V can be obtained from W by multiplying on the left by a product of matrices of the form

$$(-1)^k \begin{pmatrix} (-1)^k a_{\ell+k-1}^k & \cdots & -a_\ell^1 & 1 \\ 1 & & & \\ & \ddots & & \\ & & 1 & 0 \end{pmatrix}.$$

So that we have $V = TW$ with $V \in \mathrm{SL}_{k+1}$. This implies that the minors of order $(k+1)$ in V and W involving the same columns are equal. In particular, we have

$$\begin{aligned} d_{i,j} &= \det(W_1, \dots, W_k, W_{j-i+k+2}) = \det(V_1, \dots, V_k, V_{j-i+k+2}) \\ &= \begin{vmatrix} d_{j-w-k,i-k-1} & \cdots & d_{j-w-k,i-2} \\ \vdots & & \vdots \\ d_{j-w-1,i-k-1} & \cdots & d_{j-w-1,i-2} \end{vmatrix}. \end{aligned} \quad \blacksquare \quad (\text{B.8})$$

B.5 Gale duality

The Gale duality is a one-to-one correspondence between SL_{k+1} -frieze of width w and SL_{w+1} -frieze of width k . The correspondence is given by the following result.

Proposition B.4 ([24, Proposition 4.1.3]). *Consider a SL_{k+1} -frieze of width w , and its associated equation (B.2). The following array formed by the coefficients of the equation*

$$\begin{array}{cccccccc}
 \dots & 1 & & 1 & & 1 & & 1 & & 1 & & 1 & & 1 \\
 & & \dots & & a_n^1 & & a_1^1 & & a_2^1 & & \dots & & a_n^1 \\
 & & & a_n^2 & & a_1^2 & & a_2^2 & & & \dots & & a_n^2 \\
 \dots & & \ddots & & \ddots & & \ddots & & & & \ddots & & \dots \\
 & a_n^k & & a_1^k & & a_2^k & & \dots & & a_n^k & & \dots & \dots \\
 1 & & 1 & & 1 & & 1 & & 1 & & 1 & & \dots
 \end{array}$$

is an SL_{w+1} -frieze of width k .

Acknowledgements

I am deeply grateful to Valentin Ovsienko for sharing with me ideas and results of the preliminary version of [6]. I am also grateful to Michael Cuntz for computer calculations and Bernhard Keller for help with the applet [18]. I also want to thank Luc Pirio for stimulating discussions on the subject. This work is supported by the ANR project SC^3A , ANR-15-CE40-0004-01.

References

- [1] Assem I., Dupont G., Schiffler R., Smith D., Friezes, strings and cluster variables, *Glasg. Math. J.* **54** (2012), 27–60, [arXiv:1009.3341](#).
- [2] Assem I., Reutenauer C., Smith D., Friezes, *Adv. Math.* **225** (2010), 3134–3165, [arXiv:0906.2026](#).
- [3] Baur K., Marsh R.J., Frieze patterns for punctured discs, *J. Algebraic Combin.* **30** (2009), 349–379, [arXiv:1008.5329](#).
- [4] Bergeron F., Reutenauer C., SL_k -tilings of the plane, *Illinois J. Math.* **54** (2010), 263–300, [arXiv:1002.1089](#).
- [5] Bessenrodt C., Holm T., Jørgensen P., All SL_2 -tilings come from infinite triangulations, *Adv. Math.* **315** (2017), 194–245, [arXiv:1603.09103](#).
- [6] Conley C.H., Ovsienko V., Lagrangian configurations and symplectic cross-ratios, *Math. Ann.* **375** (2019), 1105–1145, [arXiv:1812.04271](#).
- [7] Conway J.H., Coxeter H.S.M., Triangulated polygons and frieze patterns, *Math. Gaz.* **57** (1973), 175–183.
- [8] Conway J.H., Guy R.K., The book of numbers, *Copernicus*, New York, 1996.
- [9] Coxeter H.S.M., Frieze patterns, *Acta Arith.* **18** (1971), 297–310.
- [10] Cuntz M., Holm T., Frieze patterns over integers and other subsets of the complex numbers, *J. Comb. Algebra* **3** (2019), 153–188, [arXiv:1711.03724](#).
- [11] Dupont G., An approach to non-simply laced cluster algebras, *J. Algebra* **320** (2008), 1626–1661, [arXiv:math.RT/0512043](#).
- [12] Fomin S., Williams L., Zelevinsky A., Introduction to cluster algebras, Chapters 1–3, [arXiv:1608.05735](#).
- [13] Fomin S., Zelevinsky A., Cluster algebras. I. Foundations, *J. Amer. Math. Soc.* **15** (2002), 497–529, [arXiv:math.RT/0104151](#).
- [14] Fomin S., Zelevinsky A., Cluster algebras. II. Finite type classification, *Invent. Math.* **154** (2003), 63–121, [arXiv:math.RA/0208229](#).
- [15] Fomin S., Zelevinsky A., Cluster algebras. IV. Coefficients, *Compos. Math.* **143** (2007), 112–164, [arXiv:math.RA/0602259](#).

-
- [16] Fontaine B., Plamondon P.G., Counting friezes in type D_n , *J. Algebraic Combin.* **44** (2016), 433–445, [arXiv:1409.3698](#).
- [17] Fuchs D., Tabachnikov S., Self-dual polygons and self-dual curves, *Funct. Anal. Other Math.* **2** (2009), 203–220, [arXiv:0707.1048](#).
- [18] Keller B., Quiver mutation in java, available at <https://webusers.imj-prg.fr/~bernhard.keller/>.
- [19] Keller B., Cluster algebras, quiver representations and triangulated categories, in *Triangulated Categories, London Math. Soc. Lecture Note Ser.*, Vol. 375, Cambridge University Press, Cambridge, 2010, 76–160, [arXiv:0807.1960](#).
- [20] Keller B., The periodicity conjecture for pairs of Dynkin diagrams, *Ann. of Math.* **177** (2013), 111–170, [arXiv:1001.1531](#).
- [21] Krichever I.M., Commuting difference operators and the combinatorial Gale transform, *Funct. Anal. Appl.* **49** (2015), 175–188, [arXiv:1403.4629](#).
- [22] Morier-Genoud S., Arithmetics of 2-friezes, *J. Algebraic Combin.* **36** (2012), 515–539, [arXiv:1109.0917](#).
- [23] Morier-Genoud S., Coxeter’s frieze patterns at the crossroads of algebra, geometry and combinatorics, *Bull. Lond. Math. Soc.* **47** (2015), 895–938, [arXiv:1503.05049](#).
- [24] Morier-Genoud S., Ovsienko V., Schwartz R.E., Tabachnikov S., Linear difference equations, frieze patterns, and the combinatorial Gale transform, *Forum Math. Sigma* **2** (2014), e22, 45 pages, [arXiv:1309.3880](#).
- [25] Morier-Genoud S., Ovsienko V., Tabachnikov S., 2-frieze patterns and the cluster structure of the space of polygons, *Ann. Inst. Fourier (Grenoble)* **62** (2012), 937–987, [arXiv:1008.3359](#).
- [26] Ovsienko V., Partitions of unity in $SL(2, \mathbb{Z})$, negative continued fractions, and dissections of polygons, *Res. Math. Sci.* **5** (2018), 21, 25 pages, [arXiv:1710.02996](#).
- [27] Tschabold M., Arithmetic infinite friezes from punctured discs, [arXiv:1503.04352](#).