

MATRIX PROBLEMS, TRIANGULATED CATEGORIES AND STABLE HOMOTOPY TYPES

YURIY A. DROZD

ABSTRACT. We show how the matrix problems can be used in studying triangulated categories. Then we apply the general technique to the classification of stable homotopy types of polyhedra, find out the “representation types” of such problems and give a description of stable homotopy types in finite and tame cases.

CONTENTS

1. Matrix problems arising in triangulated categories	2
2. Stable homotopy category	4
3. Discrete case: Whitehead–Chang Theorem	9
4. Tame case: Baues–Hennes Theorem	14
5. Torsion free polyhedra. Finite case	18
6. Torsion free polyhedra. Tame case	23
7. Wild cases	28
References	31

The technique of matrix problems, especially, of bimodule categories, has proved their efficiency in lots of problems from representation theory, algebraic geometry, group theory and other domains of modern algebra. During last years, mainly due to the works of Baues, Henn, Hennes, and the author, it has found new applications in algebraic topology, namely, in studying stable homotopy classes of polyhedra (see [19], [8], [4]–[7], [16]). In the survey [15] the author has picked out the background of this approach, which is based on the triangulated structure of the stable homotopy category. In this paper we show that the same method can be used in general situation, when we construct subcategories of a triangulated category from simpler ones (see Section 1). Then we summarize what can be done using this technique for

1991 *Mathematics Subject Classification.* 55P15, 55P42, 18E30, 16G60.

Key words and phrases. stable homotopy types, polyhedra, triangulated categories, matrix problems, representation type.

This paper summarizes the investigations mainly made during the stay of the author at the Max-Planck-Institut für Mathematik in Bonn. The author was also partially supported by the INTAS Grant 06-1000017-9093.

the classification problem of stable homotopy classes. Namely, we consider the subcategories $\mathcal{S}_n$ of the stable homotopy category consisting of polyhedra having only cells of n consecutive dimensions. We classify polyhedra from $\mathcal{S}_n$ for $n \leq 4$ and show that for $n > 4$ this problem becomes *wild* in the sense of the representation theory of algebras. Then we consider the subcategories $\mathcal{T}_n$ of $\mathcal{S}_n$ consisting of polyhedra with no torsion in integral homologies. This time we classify polyhedra from $\mathcal{T}_n$ for $n \leq 7$ and show that for $n > 7$ their classification is also a wild problem. In some sense, these results are “final,” though we are sure that this technique will be useful for some other problems of algebraic topology as well as for studying other triangulated categories.

Since the technical details of calculations are sometimes rather cumbersome and can be found in the previous papers, we usually omit them, just outlining the ideas.

1. MATRIX PROBLEMS ARISING IN TRIANGULATED CATEGORIES

Let $\mathcal{C}$ be a triangulated category with the shift $A \mapsto SA$, $\mathcal{A}$ and $\mathcal{B}$ be two fully additive (but usually not triangulated) subcategories of $\mathcal{C}$. We denote by $\mathcal{A} \dagger \mathcal{B}$ the full subcategory of $\mathcal{C}$ consisting of all objects C arising in triangles

$$(1.1) \quad A \xrightarrow{a} B \xrightarrow{b} C \xrightarrow{c} SA \quad \text{with } A \in \mathcal{A}, B \in \mathcal{B}.$$

We also denote by $\mathcal{I}$ the ideal of the category $\mathcal{C}$ consisting of all morphisms $\gamma : C \rightarrow C'$ that factorizes both through $\mathcal{B}$ and through $S\mathcal{A}$, i.e. such that $\gamma = \gamma'\alpha = \gamma''\beta$, where $\alpha : C \rightarrow SA$, $\beta : C \rightarrow B$, where $A \in \mathcal{A}$, $B \in \mathcal{B}$.

On the other hand, we consider the $\mathcal{A}$ - $\mathcal{B}$ -bimodule ${}_{\mathcal{B}}\mathcal{C}_{\mathcal{A}}$, which is the restriction of the regular $\mathcal{C}$ -bimodule $\mathcal{C}(A, B)$ for $A \in \mathcal{A}$, $B \in \mathcal{B}$. We often omit subscripts and denote this bimodule by $\mathcal{C}$ if it cannot lead to misunderstanding. Recall that the *bimodule category* $\mathbf{Bim}({}_{\mathcal{B}}\mathcal{C}_{\mathcal{A}})$ has $\bigcup_{\substack{A \in \mathcal{A} \\ B \in \mathcal{B}}} \mathcal{C}(A, B)$ as the set of objects, while the set of morphisms $\mathbf{Bim}(a, a')$, where $a : A \rightarrow B$, $a' : A' \rightarrow B'$, is defined as

$$\{(\alpha, \beta) \mid \alpha : A \rightarrow A', \beta : B \rightarrow B', \beta a = a' \alpha\}.$$

We denote by $\mathcal{J}$ the ideal of $\mathbf{Bim}({}_{\mathcal{B}}\mathcal{C}_{\mathcal{A}})$ consisting of all morphisms $(\alpha, \beta) : a \rightarrow a'$ such that α factors through a and β factors through a' .

We define a functor $F : \mathbf{Bim}({}_{\mathcal{B}}\mathcal{C}_{\mathcal{A}}) \rightarrow (\mathcal{A} \dagger \mathcal{B})/\mathcal{I}$ as follows. For every morphism $a : A \rightarrow B$, choose a triangle like (1.1) and set $C = Fa$. If $a' : A' \rightarrow B'$, $C' = Fa'$ and $(\alpha, \beta) \in \mathbf{Bim}(a, a')$, there is $\gamma : C \rightarrow C'$ such that the diagram

$$(1.2) \quad \begin{array}{ccccccccc} A & \xrightarrow{a} & B & \xrightarrow{b} & C & \xrightarrow{c} & SA & \xrightarrow{Sa} & SB \\ \alpha \downarrow & & \beta \downarrow & & \gamma \downarrow & & \downarrow S\alpha & & \downarrow S\beta \\ A' & \xrightarrow{a'} & B' & \xrightarrow{b'} & C' & \xrightarrow{c'} & SA' & \xrightarrow{Sa'} & SB' \end{array}$$

commutes. Set $F(\alpha, \beta) = \gamma \bmod \mathcal{I}$. We must check that the latter definition is consistent. Indeed, if $\gamma' : C \rightarrow C'$ is another morphism making diagram (1.2) commutative, $g = \gamma - \gamma'$, then $gb = c'g = 0$, therefore there are $f : SA \rightarrow C'$ and $h : C \rightarrow B'$ such that $g = cf = b'h$, i.e. $g \in \mathcal{I}$. Thus F is well-defined.

Suppose now that $\mathcal{C}(B, SA) = 0$ for all $A \in \mathcal{A}$, $B \in \mathcal{B}$. In this situation we define a functor $G : \mathcal{A} \dagger \mathcal{B} \rightarrow \mathbf{Bim}(\mathcal{B}\mathcal{C}_{\mathcal{A}})/\mathcal{I}$ as follows. Let $C \in \mathcal{A} \dagger \mathcal{B}$. Choose one triangle like (1.1) and set $a = GC$. If $GC' = a'$, i.e. C' occur in the triangle

$$A' \xrightarrow{a'} B' \xrightarrow{b'} C' \xrightarrow{c'} SA' \quad \text{with } A' \in \mathcal{A}, B' \in \mathcal{B},$$

and $\gamma : C \rightarrow C'$, then $c'\gamma b = 0$, hence $\gamma b = b'\beta$ for some $\beta : B \rightarrow B'$. Choose one of such triangles Since

$$B \xrightarrow{b} C \xrightarrow{c} SA \xrightarrow{-Sa} SB$$

and

$$B' \xrightarrow{b'} C' \xrightarrow{c'} SA' \xrightarrow{-Sa'} SB'$$

are also triangles, there is a morphism $\alpha : A \rightarrow A'$ that makes the diagram (1.2) commutative, thus $(\alpha, \beta) \in \mathbf{Bim}(a, a')$. Set $G\gamma = (\alpha, \beta) \bmod \mathcal{I}$. If (α', β') is another pair making (1.2) commutative, then $(\beta - \beta')b' = 0$, hence $\beta - \beta' = a'f$ for some $f : B \rightarrow A'$; in the same way $S\alpha - S\alpha' = g(Sa)$, i.e. $\alpha - \alpha' = g'a$ for some $g : SB \rightarrow SA'$ and $g' : B \rightarrow A'$ such that $Sg' = g$. Therefore $(\alpha - \alpha', \beta - \beta') \in \mathcal{I}$, so the functor G is well-defined.

Theorem 1.1. *Suppose that $\mathcal{C}(B, SA) = 0$ for all $A \in \mathcal{A}$, $B \in \mathcal{B}$. Then the functors F, G constructed above induce quasi-inverse functors $\bar{F} : \mathbf{Bim}(\mathcal{B}\mathcal{C}_{\mathcal{A}})/\mathcal{I} \rightarrow (\mathcal{A} \dagger \mathcal{B})/\mathcal{I}$ and $\bar{G} : (\mathcal{A} \dagger \mathcal{B})/\mathcal{I} \rightarrow \mathbf{Bim}(\mathcal{B}\mathcal{C}_{\mathcal{A}})/\mathcal{I}$. Thus $(\mathcal{A} \dagger \mathcal{B})/\mathcal{I} \rightarrow \mathbf{Bim}(\mathcal{B}\mathcal{C}_{\mathcal{A}})/\mathcal{I}$. Moreover, $\mathcal{I}^2 = 0$, therefore, the natural functor $\Pi : (\mathcal{A} \dagger \mathcal{B}) \rightarrow (\mathcal{A} \dagger \mathcal{B})/\mathcal{I}$ is an epivalence.*

Recall that an *epivalence* is a functor $E : \mathcal{C}_1 \rightarrow \mathcal{C}_2$, which is

- *full*, i.e. all induced maps $\mathcal{C}_1(X, Y) \rightarrow \mathcal{C}_2(EX, EY)$ are surjective;
- *dense*, i.e. every object from $\mathcal{C}_2$ is isomorphic to EX for some $X \in \mathcal{C}_1$;
- *conservative*, i.e. $f \in \mathcal{C}_1(X, Y)$ is invertible if and only if so is $Ef \in \mathcal{C}_2(EX, EY)$.

(In [2] such functors are called *detecting*.) Note that then also

- $X \simeq Y$ in $\mathcal{C}_1$ if and only if $EX \simeq EY$ in $\mathcal{C}_2$;
- if $\mathcal{C}_1$ and $\mathcal{C}_2$ are additive, then an object $X \in \mathcal{C}_1$ is indecomposable (into a nontrivial direct sum) if and only if so is EX .

Proof. One immediately sees that $F(\mathcal{I}) = 0$ and $G(\mathcal{I}) = 0$, hence $\bar{F}$ and $\bar{G}$ are well-defined. Moreover, we have already seen that, given

(α, β) , the morphism γ is defined up to a summand from $\mathcal{I}$, and given γ , the pair (α, β) is defined up to a summand from $\mathcal{I}$. It obviously implies that $\bar{F}\bar{G} \simeq \text{Id} \text{ nolimits}$ and $\bar{G}\bar{F} \simeq \text{Id} \text{ nolimits}$. If $\gamma : C \rightarrow C'$ and $\gamma' : C' \rightarrow C''$ are from $\mathcal{I}$, then $\gamma = gf$ for some $f : C \rightarrow B$ and $g : B \rightarrow C'$, where $B \in \mathcal{B}$, while $\gamma' = g'f'$ for some $f' : C' \rightarrow SA$ and $g' : A \rightarrow C''$, where $A \in \mathcal{A}$. Then $\gamma'\gamma = g'f'gf = 0$, since $f'g \in \mathcal{C}(B, SA)$. Thus $\mathcal{I}^2 = 0$ and, therefore, Π is an epivalence. $\square$

Corollary 1.2. *Under conditions of Theorem 1.1, let $\mathcal{V}$ be a subbimodule of ${}_{\mathcal{B}}\mathcal{C}_{\mathcal{A}}$ such that $f_1af_2 = 0$ whenever $a \in \mathcal{V}$, $f_i \in \mathcal{C}(B_i, A_i)$ with $A_i \in \mathcal{A}$, $B_i \in \mathcal{B}$ ($i = 1, 2$). Denote by $\mathcal{A} \dagger_{\mathcal{V}} \mathcal{B}$ the full subcategory of $\mathcal{A} \dagger \mathcal{B}$ consisting of all objects C arising in triangles (1.1) with $a \in \mathcal{V}$, $\mathcal{I}_{\mathcal{V}} = \mathcal{I} \cap (\mathcal{A} \dagger_{\mathcal{V}} \mathcal{B})$, $\mathcal{J}_{\mathcal{V}} = \mathcal{J} \cap \text{Bim}(\mathcal{V})$. Then the functor F and G constructed above induce quasi-inverse functors $\bar{F} : \text{Bim}(\mathcal{V})/\mathcal{J} \rightarrow (\mathcal{A} \dagger_{\mathcal{V}} \mathcal{B})/\mathcal{I}_{\mathcal{V}}$ and $\bar{G} : (\mathcal{A} \dagger_{\mathcal{V}} \mathcal{B})/\mathcal{I} \rightarrow \text{Bim}(\mathcal{V})/\mathcal{J}_{\mathcal{V}}$. Thus $(\mathcal{A} \dagger_{\mathcal{V}} \mathcal{B})/\mathcal{I} \simeq \text{Bim}(\mathcal{V})/\mathcal{J}_{\mathcal{V}}$. Moreover, $\mathcal{I}_{\mathcal{V}}^2 = 0$ and $\mathcal{J}_{\mathcal{V}}^2 = 0$, therefore, the natural functors $(\mathcal{A} \dagger_{\mathcal{V}} \mathcal{B}) \rightarrow (\mathcal{A} \dagger_{\mathcal{V}} \mathcal{B})/\mathcal{I}_{\mathcal{V}}$ and $\text{Bim}(\mathcal{V}) \rightarrow \text{Bim}(\mathcal{V})/\mathcal{J}_{\mathcal{V}}$ are epivalences. In particular, there is a one-to-one correspondences between isomorphism classes of objects and of indecomposable objects from $\mathcal{A} \dagger_{\mathcal{V}} \mathcal{B}$ and $\text{Bim}(\mathcal{V})$.*

2. STABLE HOMOTOPY CATEGORY

In this paper the word “*polyhedron*” is used as a synonym for “*finite cell* (or CW) *complex*”. We denote by **Hot** the category of punctured topological spaces with homotopy classes of continuous maps as morphisms and by **CW** its full subcategory consisting of polyhedra. We denote by CX the *cone over the space* X , i.e. the factor space $X \times I / X \times 1$, $I = [0, 1]$ being the unit interval. For a map $f : X \rightarrow Y$ we denote by Cf the *cone of this map*, i.e. the factor space $(Y \sqcup CX) / \sim$, where the equivalence relation $\sim$ is given by the rule $f(x) \sim (x, 0)$. Let also SX be the *suspension* of X , i.e. the factor space $CX / (X \times 0)$. This operation induces a functor $S : \text{Hot} \rightarrow \text{Hot}$. Note that for every X the space SX is an H -cogroup and the n -fold suspension $S^n X$ is a *commutative* H -cogroup for $n \geq 2$ [23, 2.21–2.26]. Therefore, $\text{Hot}(S^n X, Y)$ is a group, commutative for $n \geq 2$. The natural maps $\text{Hot}(S^n X, S^n Y) \rightarrow \text{Hot}(S^{n+1} X, S^{n+1} Y)$ are group homomorphisms. Set

$$\text{Hos}(X, Y) = \varinjlim_n \text{Hot}(S^n X, S^n Y).$$

It is a group called the group of *stable maps* from X to Y . Thus we get the *stable homotopy category* **Hos** and its full subcategory $\mathcal{S}$ consisting of polyhedra. We also denote by **CF** and $\mathcal{T}$ respectively the full subcategories of **CW** and of $\mathcal{S}$ consisting of *torsion free polyhedra* X , i.e. such that all integral homology groups $H_k(X) = H_k(X, \mathbb{Z})$ are

torsion free. The groups $\mathbf{Hos}(S^n, X)$ are called the *stable homotopy groups* of the space X and denoted by $\pi_n^S(X)$.


The category $\mathbf{Hos}$ is additive, with the *bouquet* (or *wedge*) $X \vee Y$ playing the role of direct sum. Moreover, $\mathbf{Hos}$ is *fully additive*, i.e. every idempotent in it splits [12, Theorem 4.8]. The suspension induces a functor, which we also denote by $S : \mathbf{Hos} \rightarrow \mathbf{Hos}$. Obviously, it is fully faithful. Thus we can “supplement” it so that S becomes an equivalence. To do it, we consider formal “imaginary spaces” $S^n X$ with $n < 0$ setting, for $n < 0$ or $m < 0$, $\mathbf{Hos}(S^n X, S^m Y) = \mathbf{Hos}(S^{n+k} X, S^{m+k} Y)$, where $k = -\min(n, m)$. Then we consider formal bouquets $\bigvee_{i=1}^r X_i$, where each X_i is either a “real” or an “imaginary” space, and define $\mathbf{Hos}(\bigvee_{j=1}^s Y_j, \bigvee_{i=1}^r X_i)$ as the set of $r \times s$ matrices (f_{ij}) with $f_{ij} \in \mathbf{Hos}(Y_j, X_i)$ (see [12] for details). As a result we get the category (also denoted by $\mathbf{Hos}$), where S is an auto-equivalence.

In fact, the new category is a *triangulated* category. The triangles in it are the *cofibration sequences*, i.e. those isomorphic to the *cone sequences*

$$X \xrightarrow{f} Y \xrightarrow{g} Cf \xrightarrow{h} SX,$$

where g is the natural embedding $Y \rightarrow Cf$ and h is the natural surjection $Cf \rightarrow SX \simeq Cf/Y$ [22]. Note that in the stable category $\mathbf{Hos}$ they coincide with the *fibration sequences* [12], though we do not use this fact.

We denote by $\mathbf{CW}_n^k$ the full subcategory of $\mathbf{CW}$ consisting of $(n-1)$ -connected cell complexes of dimension at most $n+k$. If $X \in \mathbf{CW}_n^k$, one can suppose that its $(n-1)$ -th skeleton X^{n-1} (the “ $(n-1)$ -dimensional part” of X) consist of a unique point and X has no cells of dimensions greater than $n+k$. Following Baues, we describe such a cell complex using its *gluing* (or *attachment*) *diagram*, which looks like (for $n = 7, k = 6$)



In this diagram each bullet on the level m corresponds to an m -dimensional cell, i.e. to a ball $\mathbf{B}^m$ glued to the $(m-1)$ -dimensional skeleton X^{m-1} by a map of its boundary $f : \mathbf{S}^{m-1} \rightarrow X^{m-1}$. The lines between this bullet and the lower ones describe the nonzero components of the map f . If there are more than one nonzero map between $\mathbf{S}^{m-1}$ and a smaller

S^l ($l < m$), these lines carry some marks precisising the corresponding maps. Especially, in our example the groups $\mathbf{Hos}(S^{l+3}, S^l)$ are cyclic of order 24, so we put the marks that show, which multiple of the generator is used for this gluing. There are no marks on other lines, since the groups $\mathbf{Hos}(S^{l+2}, S^l)$ are of order 2, so only have one nonzero element.

Every polyhedron from $\mathcal{S}$ decomposes into a direct sum of indecomposable ones. Note that such a decomposition is far from being unique (see [12, 4.2] for examples). Nevertheless, a description of indecomposable polyhedra in $\mathcal{S}$ can be a good first step towards the classification problem. Moreover, if the endomorphism ring $\mathbf{Es}(Y) = \mathbf{Hos}(Y, Y)$ is *local* and $Y \vee Y' \simeq \bigoplus_i Y_i$, there is an index i such that $Y_i \simeq Y \vee Y''$ [1, Lemma I.3.5] Hence, in all decompositions of a polyhedron X into bouquets of indecomposables the multiplicity of Y is the same. Another approach gives the notion of *congruence*. Namely, we say that two polyhedra X, Y are *congruent* if there is a polyhedron Z such that $X \vee Z \simeq Y \vee Z$. One can show, following [21] or [13], that an equivalent condition is that the images of X and Y in all *localizations* $\mathcal{S}_p$ of the stable homotopy category are isomorphic. Here $\mathcal{S}_p$ (p is a prime integer) is the category whose objects are polyhedra, but $\mathbf{Hos}_p(X, Y) = \mathbf{Hos}(X, Y) \otimes \mathbb{Z}_p$, where $\mathbb{Z}_p$ is the ring of p -adique integers. (The same notion is obtained if we replace $\mathbb{Z}_p$ by the subring $\{a/b \mid a, b \in \mathbb{Z}, p \nmid b\}$ of the rational numbers.) We call the classes of congruence *genera*, like they do in the theory of integral representations. Though genera satisfy the cancellation property (in fact, by definition), their decomposition into bouquets of indecomposable is not unique too (see the first of the cited examples from [12]).

Recall that due to the Generalized Freudenthal Theorem [12, Theorem 1.21] there is no need to go up to infinity in defining $\mathbf{Hos}(X, Y)$ if we deal with polyhedra. Namely, if Y is $(n-1)$ -connected and $\dim X \leq m$, then the map $\mathbf{Hot}(X, Y) \rightarrow \mathbf{Hot}(SX, SY)$ is bijective if $m < 2n-1$ and surjective if $m = 2n-1$. It implies that the map $\mathbf{Hot}(S^k X, S^k Y) \rightarrow \mathbf{Hos}(X, Y)$ is bijective for $k > m-2n+1$ and surjective for $k = m-2n+1$. In particular, if Y is $(n-1)$ -connected, $\pi_m^S(Y) \simeq \pi_{2(m-n+1)}^S(S^{m-n+2}Y)$. Moreover, on the subcategory of $\mathbf{Hot}$ consisting of simply connected spaces the suspension functor is conservative. Therefore, the induced functor $\mathbf{CW}_n^k \rightarrow \mathbf{CW}_{n+1}^k$ is an equivalence for $n > k+1$ and an epivalence for $n = k+1$. Denote by $\mathcal{S}_n$ the image in $\mathcal{S}$ of the category $\mathbf{CW}_n^{n-1}$. The polyhedra from $\mathcal{S}_n$ can only have cells on n consecutive levels (from n -th up to $(2n-1)$ -th) and every polyhedra having cells on n consecutive levels is isomorphic in $\mathcal{S}$ to $S^m X$ for some integer m and some $X \in \mathcal{S}_n$. We also denote by $\mathcal{T}_n$ the full subcategory of $\mathcal{S}_n$ consisting of torsion free polyhedra.

Definition 2.1 (cf. [3]). An *atom* is an indecomposable object A from $\mathcal{S}_n$, which does not belong to $S(\mathcal{S}_{n-1}) \cup S^2(\mathcal{S}_{n-1})$. (In other words, any polyhedron isomorphic to A in $\mathcal{S}$ must have cells of dimensions n

and $2n - 1$.) If A is an atom, all polyhedra of the sort $S^m A$ are called *suspended atoms*.

This definition immediately implies that every polyhedron is isomorphic in $\mathcal{S}$ to a bouquet of suspended atoms, though, as we have mentioned, such a decomposition is not unique. Note that, unlike Baues, we consider S^1 as an atom (a unique atom in $\mathcal{S}_1$), hence all spheres are considered as suspended atoms. Note also that this definition implies that all atoms are of odd dimensions: an atom from $\mathcal{S}_n$ is of dimension $2n - 1$.

To clarify the structure of $\mathcal{S}_n$ we use the technique from Section 1. Namely, choose an integer m such that $0 \leq m < n - 1$ and set

$$\begin{aligned}
 \mathcal{A} &= \mathcal{A}_{n,m} = S^{2m+1} \mathcal{S}_{n-m-1}, \\
 \mathcal{B} &= \mathcal{B}_{n,m} = S^{n-m-1} \mathcal{S}_{m+1}, \\
 \mathcal{S}_{n,m} &= \mathcal{B}_{n,m} \mathcal{S}_{\mathcal{A}_{n,m}}, \\
 (2.2) \quad \mathcal{I}_{n,m} &= \{ f : X \rightarrow Y \mid X, Y \in \mathcal{S}_n, f \text{ factors both} \\
 &\quad \text{through } \mathcal{B} \text{ and through } S\mathcal{A} \}, \\
 \mathcal{J}_{n,m} &= \{ (\alpha, \beta) \in \text{Bim}(a, a') \mid a, a' \in \mathcal{S}_{n,m}, \\
 &\quad \alpha \text{ factors through } a \text{ and } \beta \text{ factors through } a' \}.
 \end{aligned}$$

Then polyhedra from $\mathcal{A}$ only have cells in dimensions from $n + m$ up to $n - m - 2$, while those from $\mathcal{B}$ only have cell in dimensions from n up to $n + m$. If $C \in \mathcal{S}_n$, its $(n + m)$ -th skeleton B belongs to $\mathcal{B}$, while the factor space C/B belongs to $S\mathcal{A}$, i.e. $C/B \simeq SA$, $A \in \mathcal{A}$. Then $C \in \mathcal{A} \dagger \mathcal{B}$, since $A \rightarrow B \rightarrow C \rightarrow C/B \simeq A$ is a cofibration sequence. On the other hand, any object from $\mathcal{A} \dagger \mathcal{B}$ obviously belongs to $\mathcal{S}_n$. So we have proved

Theorem 2.2. $\mathcal{S}_n \simeq \mathcal{A}_{n,m} \dagger \mathcal{B}_{n,m}$. Thus $\mathcal{S}_n / \mathcal{I}_{n,m} \simeq \text{Bim}(\mathcal{S}_{n,m}) / \mathcal{J}_{n,m}$. Moreover, $\mathcal{J}_{n,m}^2 = 0$.

To consider torsion free polyhedra, we set

$$\begin{aligned}
 \mathcal{A}^0 &= \mathcal{A}_{n,m}^0 = S^{2m+1} \mathcal{S}_{n-m-1}, \\
 \mathcal{B}^0 &= \mathcal{B}_{n,m}^0 = S^{n-m-1} \mathcal{S}_{m+1}, \\
 (2.3) \quad \mathcal{S}^0 &= \mathcal{S}_{n,m}^0 = \{ a \in \mathcal{S}_{n,m} \mid H_{n+m}(a) = 0 \}, \\
 \mathcal{I}_{n,m}^0 &= \mathcal{I}_{n,m} \cap (\mathcal{A}^0 \dagger_{\mathcal{S}^0} \mathcal{B}^0), \\
 \mathcal{J}_{n,m}^0 &= \mathcal{J}_{n,m} \cap \text{Bim}(\mathcal{S}^0).
 \end{aligned}$$

To get an analogue of Theorem 2.2 we need the following lemma.

Lemma 2.3. *Let $f \in \text{Hos}(A, B)$, where A and B are torsion free polyhedra, A is $(m - 1)$ -connected, $\dim B \leq m$ and Cf is also torsion free. There are decompositions $A \simeq C \oplus A'$, $B \simeq C \oplus B'$ such that, with*

respect to this decomposition, $f = \begin{pmatrix} \text{Id } \text{nolimits} & 0 \\ 0 & g \end{pmatrix}$ with $H_m(g) = 0$ and $Cf \simeq Cg$.

Proof. Note first that if $A = kS^m$, $B = lS^m$ are bouquets of m -dimensional spheres, then $H_m(A) = m\mathbb{Z}$, $H_m(B) = l\mathbb{Z}$, and the natural map $\text{Hos}(A, B) \rightarrow \text{Hom } \text{nolimits}(H_m(A), H_m(B))$ is an isomorphism. In particular, every decomposition of $H_m(A)$ arises from a decomposition of A , and the same is true for B . In this case $H_m(f)$, or, the same, f is actually an integer matrix and there are decompositions $A \simeq C \oplus A'$, $B \simeq C \oplus B'$ (all summands are, of course, also bouquets of spheres) such that, with respect to them, $f = \begin{pmatrix} \text{Id } \text{nolimits} & 0 \\ 0 & d \end{pmatrix}$, where $d : A' \rightarrow B'$ can be presented by diagonal matrix without unit components.

In general case, the calculation of homologies of cell spaces from [23, Chapter 10] shows that the embedding $\alpha : A^m \rightarrow A$ induces a surjection $H_m(A^m) \rightarrow H_m(A)$, while the surjection $\beta : B \rightarrow \tilde{B} = B/B^{m-1}$ induces an embedding $H_m(B) \rightarrow H_m(\tilde{B})$ with torsion free cokernel. Therefore, there are decompositions $A^m \simeq A_1 \oplus A_0$, $\tilde{B} \simeq B_1 \oplus B_0$ such that the restriction of $H_m(\alpha)$ onto A_1 is an isomorphism, and that onto A_0 is 0, while $H_m(\beta)$ induces an isomorphism $H_m(B) \rightarrow \text{im } \text{nolimits} H_m(\beta) = H_m(B_1)$. Denote by $\alpha_1 : A_1 \rightarrow A$ and $\beta_1 : B \rightarrow B_1$ the corresponding components of α and β . As above, there are decompositions $A_1 \simeq C \oplus A_0$, $B_1 \simeq C \oplus B_0$ such that, with respect to them, the morphism $\beta_1 f \alpha_1 = \begin{pmatrix} \text{Id } \text{nolimits} & 0 \\ 0 & d \end{pmatrix}$, where d can be presented by diagonal matrix without unit components. Denote by $\iota : C \rightarrow A_1$ the natural embedding (presented by the matrix $\begin{pmatrix} \text{Id } \text{nolimits} \\ 0 \end{pmatrix}$) and by $\pi : B_1 \rightarrow C$ the natural projection (presented by the matrix $\begin{pmatrix} \text{Id } \text{nolimits} & 0 \end{pmatrix}$). Then $\pi \beta_1 f \alpha_1 \iota = \text{Id } \text{nolimits}$, so $B \simeq C \oplus B'$, $A \simeq C \oplus A'$, so that, with respect to these decompositions, $f = \begin{pmatrix} \text{Id } \text{nolimits} & 0 \\ 0 & g \end{pmatrix}$. Then $Cf \simeq Cg$ and $d = \beta_0 g \alpha_0$, where $\alpha_0 : A_0 \rightarrow A'$ and $\beta_0 : B' \rightarrow B_0$. Note that α_0 and β_0 also induce isomorphisms of the m -th homology groups, so $\text{Coker } \text{nolimits} H_m(g) \simeq \text{Coker } \text{nolimits} H_m(d)$. Since this cokernel embeds in $H_m(Cg)$, it is torsion free. Therefore, $d = 0$, whence $H_m(g) = 0$. $\square$

Theorem 2.4. $\mathcal{T}_n \simeq \mathcal{A}_{n,m}^0 \dagger_{\mathcal{J}^0} \mathcal{B}_{n,m}^0$. Thus $\mathcal{T}_n / \mathcal{J}_{n,m}^0 \simeq \text{Bim}(\mathcal{J}_{n,m}^0) / \mathcal{J}_{n,m}^0$. Moreover, $(\mathcal{J}_{n,m}^0)^2 = (\mathcal{J}_{n,m}^0)^2 = 0$, so this equivalence induces one-to-one correspondences between isomorphism classes of objects and of indecomposable objects in $\mathcal{T}_n$ and in $\text{Bim}(\mathcal{J}_{n,m}^0)$.

Proof. Let $C \in \mathcal{T}_n$, $B = C^{n+m}$, $SA \simeq C/B$. The triangle

$$(2.4) \quad A \xrightarrow{a} B \xrightarrow{b} C \xrightarrow{c} SA$$

gives rise to the exact sequence of homologies

$$\begin{aligned} \cdots \rightarrow H_k(A) &\xrightarrow{H_k(a)} H_k(B) \xrightarrow{H_k(b)} H_k(C) \xrightarrow{H_k(c)} H_k(SA) \simeq \\ &\simeq H_{k-1}(A) \xrightarrow{H_{k-1}(a)} H_{k-1}(B) \xrightarrow{H_{k-1}(b)} H_{k-1}(C) \rightarrow \cdots \end{aligned}$$

If $k < n+m$, then $H_k(A) = H_{k-1}(A) = 0$, so $H_k(B) \simeq H_k(C)$ is torsion free. If $k > n+m$, we get in the same way that $H_k(A) \simeq H_{k+1}(C)$ is also torsion free. Let now $k = n+m$, then we get the exact sequence

$$0 \rightarrow H_{n+m+1}(C) \rightarrow H_{n+m}(A) \xrightarrow{H_{n+m}(a)} H_{n+m}(B) \rightarrow H_{n+m}(C) \rightarrow 0.$$

Note that $H_{n+m}(B)$ is always torsion free, since B contains no cells of dimensions bigger than $n+m$, hence $B \in \mathcal{T}$. Therefore, $H_{n+m}(A)$ is torsion free too, so $A \in \mathcal{T}$. Moreover, $\text{Coker } \text{Id} \text{ } \text{Id} \text{ } H_{n+m}(a)$ is also torsion free. As both $H_{n+m}(A)$ and $H_{n+m}(B)$ are free, it means that $H_{n+m}(A) \simeq M \oplus M'$, $H_{n+m}(B) \simeq M \oplus M''$ so that $H_{n+m}(a)$ induces isomorphism $M \rightarrow M$ and is zero on M' . By Lemma 2.3, there are decompositions $A \simeq A_0 \vee A'$, $B \simeq A_0 \vee B'$ such that, with respect to them, $a = \begin{pmatrix} \text{Id} \text{ } \text{Id} \text{ } 0 \\ 0 & a' \end{pmatrix}$, where $a' \in \mathcal{S}^0$. Then $Ca' \simeq Ca \simeq C$, so $C \in \mathcal{A}^0 \upharpoonright_{\mathcal{S}^0} \mathcal{B}^0$. On the other hand, if $C \in \mathcal{A}^0 \upharpoonright_{\mathcal{S}^0} \mathcal{B}^0$, i.e. belongs to a triangle (2.4) with $A \in \mathcal{A}^0$, $B \in \mathcal{B}^0$ and $H_{n+m}(a) = 0$, the exact sequence of homologies implies that $C \in \mathcal{T}_n$.

To prove the remaining assertions, it is enough to show that $uav = 0$ for every $a \in \mathcal{S}^0(A, B)$, $v : B' \rightarrow A$, $u : B \rightarrow A'$, where $A, A' \in \mathcal{A}^0$, $B, B' \in \mathcal{B}^0$ (see Corollary 1.2). Since $H_{n+m}(a) = 0$, the induced map $A^{m+n} \rightarrow B/B^{m+n-1}$ is zero. On the other hand, $\mathcal{S}(B', A/A^{m+n}) = 0 = \mathcal{S}((B')^{m+n-1}, A^{m+n})$, so the map $v : B' \rightarrow A$ factors through a map $B'/(B')^{m+n-1} \rightarrow A^{m+n}$. Since the same holds for u , it implies that $uav = 0$. $\square$

We shall also use the following obvious lemma.

Lemma 2.5. *Let $X \in \mathcal{S}_n$, $H_i = H_i(X)$. If X is decomposable, there are decompositions $H_i = H'_i \oplus H''_i$ and indices j, k such that both $H'_j \neq 0$ and $H''_k \neq 0$.*

(Note also that $H_i(X) = 0$ for $i < n$ or $i > 2n - 1$.)

3. DISCRETE CASE: WHITEHEAD-CHANG THEOREM

We apply now Theorem 2.2 to polyhedra from $\mathcal{S}_n$ for small n . First, we recall some values of stable homotopy groups [20, Sections XI.15–16]:

- $\pi_{n+1}^S(S^n) \simeq \mathbb{Z}/2$, the generator being the (suspended) *Hopf map* $\eta = 2^{n-1}h_2$, where h_2 is the Hopf fibration $S^3 \rightarrow S^2$;

- $\pi_{n+2}^S(S^n) \simeq \mathbb{Z}/2$, the generator being the *double Hopf map* η^2 , i.e. the composition of Hopf maps $S^{n+2} \rightarrow S^{n+1} \rightarrow S^n$;
- $\pi_{n+3}^S(S^n) \simeq \mathbb{Z}/24$, the generator being $\nu = S^{n-4}h_4$, where h_4 is the Hopf fibration $S^7 \rightarrow S^4$. Moreover, the composition $\eta^3 : S^{n+3} \rightarrow S^{n+2} \rightarrow S^{n+1} \rightarrow S^n$ equals 12ν .

If $n = 1$, the only atom in $\mathcal{S}_1$ is S^1 , and every polyhedron is a bouquet of several copies of S^1 . If $n = 2$, $\mathcal{S}_2 = \mathcal{A}_{2,0} \dagger \mathcal{B}_{2,0}$, and $\mathcal{A}_{2,0} = \mathcal{B}_{2,0} = S\mathcal{S}_1$. Thus every polyhedron C from $\mathcal{S}_2$ is isomorphic to the cone of a map $a : kS^2 \rightarrow lS^2$. Since $\mathbf{Hos}(S^2, S^2) = \mathbb{Z}$, the map a can be considered as a matrix $(a_{ij}) \in \text{Mat } \textit{nolimits}(l \times k, \mathbb{Z})$. If a' is another object from $\mathcal{C}_{2,1}$, also considered as a matrix from $\text{Mat } \textit{nolimits}(l' \times k', \mathbb{Z})$, a morphism $a \rightarrow a'$ in $\mathbf{Bim}(\mathcal{C}_{2,1})$ is given by a pair of matrices $\alpha \in \text{Mat } \textit{nolimits}(k' \times k, \mathbb{Z})$, $\beta \in \text{Mat } \textit{nolimits}(l' \times l, \mathbb{Z})$ such that $a'\alpha = \beta a$. Especially, this morphism is an isomorphism if and only if both α and β are invertible. So the well-known Smith Theorem implies that every object $a \in \mathcal{C}_{2,1}$ is isomorphic to one presented by a diagonal matrix $\text{diag } \textit{nolimits}(q_1, q_2, \dots, q_r)$. Hence, every polyhedron from $\mathcal{S}_2$ is isomorphic to a bouquet of cones $\bigvee_i Cq_i$, where we identify an integer q with the corresponding map $S^2 \rightarrow S^2$. Moreover, if $q = uv$, where $\gcd(u, v) = 1$, then

$$Cq \simeq C \begin{pmatrix} 1 & 0 \\ 0 & q \end{pmatrix} \simeq C \begin{pmatrix} u & 0 \\ 0 & v \end{pmatrix} \simeq Cu \vee Cv.$$

Therefore, Cq can only be indecomposable if $q = p^s$, where p is prime. On the other hand, the exact sequence of homologies arising from the triangle

$$(3.1) \quad S^2 \xrightarrow{q} S^2 \rightarrow M^3(q) \rightarrow S^3$$

that $H_2(Cq) \simeq \mathbb{Z}/q$ and $H_3(Cq) = 0$. Hence, Lemma 2.5 implies that Cq is indecomposable. Therefore, the atoms in $\mathcal{S}_2$ are just Cq for $q = p^s$ with a prime p . These atoms are denoted by $M^3(q)$ and their suspensions $S^k M^3(q)$ by $M^{k+3}(q)$. The atoms and suspended atoms $M^d(q)$ are called *Moore spaces*¹ [12]. We also write M_s^d instead of $M^d(2^s)$ (these atoms play a special role later).

We can calculate the groups $\mathbf{Hos}(M^3(q), M^3(q'))$. Since $\pi_3^S(S^2) \simeq \mathbb{Z}/2$ [20, Theorem 15.1], the exact sequences for the functor $\mathbf{Hos}$ arising from the triangles (3.1) for q and q' imply that

$$\begin{aligned} \mathbf{Hos}(S^2, M^3(q)) &\simeq \mathbf{Hos}(M^3(q), S^3) \simeq \mathbb{Z}/q, \\ \mathbf{Hos}(S^3, M^3(q)) &\simeq \mathbf{Hos}(M^3(q), S^2) \simeq \begin{cases} \mathbb{Z}/2 & \text{if } q \text{ is even,} \\ 0 & \text{if } q \text{ is odd,} \end{cases} \\ \mathbf{Hos}(M^3(q), M^3(q')) &\simeq \mathbb{Z}/(q, q') \text{ if } q \text{ or } q' \text{ is odd,} \end{aligned}$$

¹In [20, Section XI.10] they are denoted by P_q^d and called *pseudo-projective spaces*.

and there is an exact sequence

$$(3.2) \quad 0 \rightarrow \mathbb{Z}/2 \rightarrow \text{Hos}(M_s^3, M_r^3) \rightarrow \mathbb{Z}/2^m \rightarrow 0, \text{ where } m = \min(r, s).$$

Note that the endomorphism rings $\text{Es}(M^3(q))$ are finite, hence, local. These considerations immediately imply the description of polyhedra from $\mathcal{S}_2$.

Theorem 3.1. *Every polyhedron from $\mathcal{S}_2$ uniquely (up to permutation of summands) decomposes into a bouquet of spheres S^2, S^3 and Moore atoms $M^3(q)$.*

We also need the following fact.

Proposition 3.2.

$$\pi_4^S(M^3(q)) \simeq \begin{cases} 0 & \text{if } q \text{ is odd,} \\ \mathbb{Z}/4 & \text{if } q = 2, \\ \mathbb{Z}/2 \oplus \mathbb{Z}/2 & \text{if } q = 2^s, \ s > 1. \end{cases}$$

Proof. Recall that $\pi_4^S(S^3) \simeq \pi_4^S(S^2) \simeq \mathbb{Z}/2$ [20, Theorems 15.1, 15.2]. Therefore, the exact sequence for π_4^S arising from (3.1) shows that $\pi_4^S(M^3(q)) = 0$ for q odd and, for $q = 2^s$, there is an exact sequence

$$0 \rightarrow \mathbb{Z}/2 \rightarrow \pi_4^S(M_s^3) \rightarrow \mathbb{Z}/2 \rightarrow 0.$$

Note that $\pi_4^S(M_1^3) \simeq \pi_6(M_1^5)$, so [20, Lemma 10.2] implies that it embeds into $\pi_6(S^3) \simeq \mathbb{Z}/12$ [20, Theorem 16.1]. Hence, $\pi_4^S(M_1^3) \simeq \mathbb{Z}/4$. For $r > 1$ consider the commutative diagram of triangles

$$(3.3) \quad \begin{array}{ccccccc} S^2 & \xrightarrow{2} & S^2 & \longrightarrow & M_1^3 & \longrightarrow & S^3 \\ 1 \downarrow & & 2^{r-1} \downarrow & & \downarrow & & 1 \downarrow \\ S^2 & \xrightarrow{2^r} & S^2 & \longrightarrow & M_s^3 & \longrightarrow & S^3, \end{array}$$

It induces the commutative diagram with exact rows

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathbb{Z}/2 & \longrightarrow & \pi_4^S(M_1^3) & \longrightarrow & \mathbb{Z}/2 & \longrightarrow & 0 \\ & & 0 \downarrow & & \downarrow & & \downarrow 1 & & \\ 0 & \longrightarrow & \mathbb{Z}/2 & \longrightarrow & \pi_4^S(M_s^3) & \longrightarrow & \mathbb{Z}/2 & \longrightarrow & 0, \end{array}$$

which shows that the second row is the pushdown of the first one along zero map, hence, it splits. $\square$

Proposition 3.3.

$$\text{Hos}(M_s^d, M_r^d) \simeq \begin{cases} \mathbb{Z}/4 & \text{if } r = s = 1, \\ \mathbb{Z}/2 \oplus \mathbb{Z}/2^m & \text{otherwise, where } m = \min(r, s). \end{cases}$$

Proof. Obviously, we may suppose that $m = 3$. Since $\pi_4^S(M_1^3) \simeq \mathbb{Z}/4$ is a module over the ring $\text{Hos}(M_1^3, M_1^3)$, $2\text{Hos}(M_1^3, M_1^3) \neq 0$, hence, $\text{Hos}(M_1^3, M_1^3) \simeq \mathbb{Z}/4$. On the other hand, applying the functor $\text{Hos}(_, M_1^3)$ to the diagram (3.3) with $s > 1$, we get a commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{Z}/2 & \longrightarrow & \text{Hos}(M_1^3, M_1^3) & \longrightarrow & \mathbb{Z}/2 \longrightarrow 0 \\ & & \uparrow 1 & & \uparrow & & \uparrow 0 \\ 0 & \longrightarrow & \mathbb{Z}/2 & \longrightarrow & \text{Hos}(M_s^3, M_1^3) & \longrightarrow & \mathbb{Z}/2 \longrightarrow 0. \end{array}$$

Thus its second row is the pull-back of the first one along the zero map, hence, it splits. The dual consideration shows that the sequence (3.2) for $r > 1$ can be obtained as a pushdown of the sequence for $r = 1$, hence, it splits too. $\square$

Note that the latter decomposition in this statement is that of *groups*. Taking into account the multiplication, it is convenient to present morphisms $M_s^d \rightarrow M_r^d$ as triangular matrices $\begin{pmatrix} a & b \\ 0 & c \end{pmatrix}$, with $a \in \mathbb{Z}/2^r$, $b \in \mathbb{Z}/2$, $c \in \mathbb{Z}/2^s$, $2^{s-m}a \equiv 2^{r-m}c \pmod{2^\mu}$, where $m = \min(s, r)$, $\mu = \max(s, r)$. The product of morphisms correspond then to the usual product of matrices, while the sum of morphisms correspond to the usual sum of matrices, with the only exception, when $s = r = 1$: then we must add matrices as follows:

$$\begin{pmatrix} a & b \\ 0 & a \end{pmatrix} + \begin{pmatrix} a' & b' \\ 0 & c' \end{pmatrix} = \begin{pmatrix} a + a' & b + b' + aa' \\ 0 & a + a' \end{pmatrix}.$$

Let now $n = 3, m = 1$, then $\mathcal{S}_3 = \mathcal{A}_{3,1} \upharpoonright \mathcal{B}_{3,1}$, where $\mathcal{A}_{3,1} = S^3\mathcal{S}_1$ and $\mathcal{B}_{3,1} = S\mathcal{S}_2$. Hence, polyhedra from $\mathcal{A}_{3,1}$ are just bouquets of spheres S^4 , while those from $\mathcal{B}$ are bouquets of spheres S^4, S^3 and Moore spaces $M^4(q)$. For convenience, we set $M_0^4 = S^4$ and $M_\infty^4 = S^3$ and order the set of indices by the rule $1 < 2 < \dots < \infty < 0$. As we have seen, $\text{Hos}(S^4, M^4(q)) = 0$ for q odd, $\text{Hos}(S^4, M_r^4) = H_r \simeq \mathbb{Z}/2$ for $r \neq 0$ and $\text{Hos}(S^4, S^4) = H_0 \simeq \mathbb{Z}$. Therefore, a map $a : A \rightarrow B$, where $A \in \mathcal{A}$, $B \in \mathcal{B}$ can be presented as a block matrix


$$(3.4) \quad a = \begin{pmatrix} a_0 \\ a_\infty \\ \vdots \\ a_2 \\ a_1 \end{pmatrix},$$

where a_s are matrices over H_s . One easily sees that if η_s is a generator of H_s and $\beta_{rs} : M_s^4 \rightarrow M_r^4$, then $\beta_{rs}\eta_s = 0$ if $r > s$, while for $r \leq s$ the

map β_{rs} can be so chosen that $\beta_{rs}\eta_s = \eta_r$. Set

$$H_{rs} = \begin{cases} 0 & \text{if } r > s, \\ \mathbb{Z}/2 & \text{if } 0 \neq r \leq s, \\ \mathbb{Z} & \text{if } s = r = 0. \end{cases}$$

Therefor two matrices a, a' of the form (3.4) define isomorphic objects from $\text{Bim}(\mathcal{S}_{3,1})$ if and only if there is an invertible integral matrix α and an invertible block matrix $\beta = \beta_{rs}$, where β_{rs} is a matrix over H_{rs} , such that $a' = \beta a \alpha^{-1}$. Then simple considerations show that every object from $\text{Bim}(\mathcal{S}_{3,1})$ decomposes into a direct sum of objects given by the 1×1 matrices $q \in H_0$, $\eta_r \in H_r$, $r \neq 0$ and $\begin{pmatrix} 2^s \\ \eta_r \end{pmatrix} \in H_0 \oplus H_r$, $r \neq 0, s > 0$. The first case correspond to the Moore space $M^5(q)$, while the second and the third cases define new polyhedra, respectively, $C^5(\eta)$, $C^5(2^r\eta)$, $C^5(\eta 2^s)$ and $C^5(2^r\eta 2^s)$, given by the gluing diagrams



(The words in brackets show the corresponding gluings.)

To find endomorphisms of these atoms, note that there are triangles

$$(3.5) \quad S^3 \vee S^4 \xrightarrow{\begin{pmatrix} 2^r & \eta \end{pmatrix}} S^3 \rightarrow C^5(2^r\eta) \rightarrow S^4 \vee S^5,$$

$$(3.6) \quad S^4 \xrightarrow{\begin{pmatrix} \eta \\ 2^s \end{pmatrix}} S^3 \vee S^4 \rightarrow C^5(\eta 2^s) \rightarrow S^5,$$

$$(3.7) \quad S^3 \vee S^4 \xrightarrow{\begin{pmatrix} 2^r & \eta \\ 0 & 2^s \end{pmatrix}} C^5(2^r\eta 2^s) \rightarrow S^3 \vee S^4.$$

By Theorem 1.1, $\text{Es}(C^5(2^r\eta))$, up to an ideal I such that $I^2 = 0$, is isomorphic to the endomorphism ring of the map $f = \begin{pmatrix} 2^r & \eta \end{pmatrix}$ in the category $\text{Bim}(\mathcal{S})/\mathcal{J}$. An endomorphism of f in $\text{Bim}(\mathcal{S})$ is a pair (α, β) , where $\alpha = \begin{pmatrix} a & b\eta \\ 0 & c \end{pmatrix}$ ($a, c \in \mathbb{Z}$, $b \in \mathbb{Z}/2$), $\beta \in \mathbb{Z}$, such that $\beta f = f\alpha$, i.e. $\beta = a \equiv c \pmod{2}$. Moreover, one easily sees that $\mathcal{J}$ consists of the pairs with the first component $\begin{pmatrix} 2^r x & x\eta \\ 0 & 0 \end{pmatrix}$, whence $\text{Es}(C^5(2^r\eta))/I^2$ is isomorphic to the subring of $\mathbb{Z}/2^{r+1} \oplus \mathbb{Z}$ consisting of all pairs (a, c) with $a \equiv c \pmod{2}$. This ring has no nontrivial idempotent, hence, $C^5(2^r\eta)$ is indeed indecomposable, hence, an atom. Moreover, using the triangle (3.5), one can see that $I \simeq \mathbb{Z}/2$ and

$\text{Es}(C^5(2^r\eta))$ is isomorphic to the ring of triangular matrices $\begin{pmatrix} a & b \\ 0 & c \end{pmatrix}$, where $a \in \mathbb{Z}/2^{r+1}$, $b \in \mathbb{Z}/2$, $c \in \mathbb{Z}$, $a \equiv c \pmod{2}$. The same result for $\text{Es}(C^5(\eta 2^r))$ follows from the triangle (3.6). Finally, one gets from the triangle (3.7) that $\text{Es}(C^5(2^r\eta 2^s))$ is isomorphic to the ring of triangular matrices $\begin{pmatrix} a & b \\ 0 & c \end{pmatrix}$, where $a \in \mathbb{Z}/2^r$, $b \in \mathbb{Z}/2$, $c \in \mathbb{Z}/2^s$, $a \equiv c \pmod{2}$. Therefore these polyhedra are also atoms. They are called *Chang atoms*. Moreover, the last ring is local, thus the multiplicity of $C^5(2^r\eta 2^s)$ (as well as of any its shift) in a decomposition of a polyhedron into a bouquet of indecomposables is the same for all such decompositions. Note that the same is true for suspended atoms $M^d(q)$. On the other hand, the triangles (3.5) and (3.6) imply that $H_3(C^5(2^r\eta)) \simeq H_4(C^5(\eta 2^r)) \simeq \mathbb{Z}/2^r$, while other homologies of these spaces are zero. Altogether, it gives the following description of the category $\mathcal{S}_3$.

Theorem 3.4 (Whitehead–Chang, [25, 11]). *Any polyhedron from $\mathcal{S}_3$ uniquely (up to permutation of summands) decomposes into a bouquet of spheres S^3, S^4, S^5 , suspended Moore atoms $M^4(q), M^5(q)$ and Chang atoms $C^5(\eta), C^5(2^r\eta), C^5(\eta 2^s)$ and $C^5(2^r\eta 2^s)$.*

Using terms from the representation theory, one can say that the categories $\mathcal{S}_n$, $n \leq 3$, are *discrete* (or *essentially finite*). In this context it means that there are only finitely many isomorphism classes of polyhedra in $\mathcal{S}_n$ with a prescribed exponent of the torsion part of homologies. (So it looks similar to the description of finitely generated abelian groups.)

4. TAME CASE: BAUES–HENNES THEOREM

We study now the category $\mathcal{S}_4$. By Theorem 2.2, $\mathcal{S}_4 = \mathcal{A} \dagger \mathcal{B}$, where $\mathcal{A} = S^3\mathcal{S}_2$, $\mathcal{B} = S^2\mathcal{S}_2$. By Theorem 3.1, every polyhedron from $\mathcal{A}$ (from $\mathcal{B}$) is a bouquet of spheres S^5, S^6 and Moore atoms $M^6(q)$ (respectively, S^4, S^5 and $M^5(q)$). We have already calculated morphisms between indecomposables in $\mathcal{S}_2$. Just in the same way one calculates morphisms from the objects of $S^3\mathcal{S}_2$ and those of $S^2\mathcal{S}_2$. We omit the details, which are standard; the result is presented in Table 1. Actually, the groups $\text{Hos}(M_s^6, M_r^5)$ can be naturally considered as the groups of upper triangular matrices $\begin{pmatrix} a & b \\ 0 & c \end{pmatrix}$ over $\mathbb{Z}/2$ with $b = 0$ if $s = r = 1$. Again the sum of morphisms correspond to the usual sum of matrices, with the exceptions for $s > 1, r = 1$ and $s = 1, r > 1$, when

TABLE 1.

	S^5	S^6	M_1^6	M_s^6 ($s > 1$)
S^4	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/4$	$\mathbb{Z}/2 \oplus \mathbb{Z}/2$
S^5	$\mathbb{Z}$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/2$
M_1^5	$\mathbb{Z}/2$	$\mathbb{Z}/4$	$\mathbb{Z}/2 \oplus \mathbb{Z}/2$	$\mathbb{Z}/4 \oplus \mathbb{Z}/2$
M_r^5 ($r > 1$)	$\mathbb{Z}/2$	$\mathbb{Z}/2 \oplus \mathbb{Z}/2$	$\mathbb{Z}/2 \oplus \mathbb{Z}/4$	$\mathbb{Z}/2 \oplus \mathbb{Z}/2 \oplus \mathbb{Z}/2$

the sum of matrices must be twisted as follows:

$$\begin{pmatrix} a & b \\ 0 & c \end{pmatrix} + \begin{pmatrix} a' & b' \\ 0 & c' \end{pmatrix} = \begin{pmatrix} a + a' & b + b' + aa' \\ 0 & c + c' \end{pmatrix} \text{ if } s > 1, r = 1,$$

$$\begin{pmatrix} a & b \\ 0 & c \end{pmatrix} + \begin{pmatrix} a' & b' \\ 0 & c' \end{pmatrix} = \begin{pmatrix} a + a' & b + b' + cc' \\ 0 & c + c' \end{pmatrix} \text{ if } s = 1, r > 1.$$

The multiplication of elements from $\mathbf{Hos}(M_s^6, M_r^5)$ by morphisms between objects from $\mathcal{A}$ and $\mathcal{B}$ (also presented by triangular matrices as in Section 3) correspond to the usual product of matrices. Therefore, a morphism $A \rightarrow B$ can be naturally considered as a block matrix presented in Table 2. In this table a symbol 2 (∞) shows that the

TABLE 2.


	(1)	(2)	(3)	...		...	(3)	(2)	(1)
(1)	2	2	2	...	2	2	...	2	2
(2)	2	2	2	...	2	2	...	2	2
(3)	2	2	2	...	2	2	...	2	2
	$\vdots$	$\vdots$	$\vdots$	$\ddots$	$\vdots$	$\vdots$	$\ddots$	$\vdots$	$\vdots$
	2	2	2	...	2	2	...	2	2
	0	0	0	...	∞	2	...	2	2
	$\vdots$	$\vdots$	$\vdots$	$\ddots$	$\vdots$	$\vdots$	$\ddots$	$\vdots$	$\vdots$
(3)	0	0	0	...	0	2	...	2	2
(2)	0	0	0	...	0	2	...	2	2
(1)	0	0	0	...	0	2	...	2	2

corresponding block has values from $\mathbb{Z}/2$ (respectively, from $\mathbb{Z}$). Zeros show that the corresponding block is always zero. Arrows on the left

and below symbolize the action of morphisms between the objects from $\mathcal{A}$ and $\mathcal{B}$ respectively. The labels $(_1), (_2), \dots$ (or $(^1), (^2), \dots$) show that the corresponding horizontal (respectively, vertical) stripes are of the same size and we must use the same elementary transformations in both of them. These stripes correspond to M_r^d with the same d and r . Note that there are 2 horizontal and 2 vertical stripes without such labels. They correspond to spheres S^d .


This matrix problem is a slight variation of a well-known one, namely, representations of *bunches of chains* (see [9] or [10, Appendix B]). It implies a description of indecomposable objects in the category $\text{Bim}(\mathcal{S}_{4,2})$, hence, in $\mathcal{S}_4$. We call them *strings* and *bands*, as it is usual in the representation theory of algebras. Not providing details (see [15]), we just present the corresponding attachment diagrams (Table 3). It is convenient to distinguish two types of strings: *usual* and *decorated*; I hope that the pictures show the difference. “Decorations” (one for each string) are shown with double lines. We omit integers precisising the degrees of “vertical” attachments, as well as one precisising the “long” attachment in a decorated strings of the first kind; they can be arbitrary and differ for different attachments. Certainly, each diagram is actually finite: it starts at any place on the left and stops at any place on the right.

Multiple bullets in the case of bands symbolize not a unique cell but several (say m) copies of it (the same for each ball). All attachments except the one marked by the wavy line are “natural”: the first copy of an upper cell is attached to the first copy of a lower one, the second to the second, etc. The attachment marked by the wavy line is “twisted” by an invertible Frobenius matrix Φ of size $m \times m$ over the field $\mathbb{Z}/2$ with the characteristic polynomial $f(x)$, which must be a power of an irreducible one and such that $f(0) \neq 0$. For instance, if $f(x) = x^3 + x + 1$, i.e. $m = 3$ and $\Phi = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 1 \end{pmatrix}$, this attachment is:




One can check that all strings and bands are indecomposable and pairwise non-isomorphic. Note also that all atoms from $\mathcal{S}_4$ are p -primary (2-primary, except Moore atoms $M^d(p^r)$ with odd p , which are p -primary). Therefore, we have the uniqueness of decomposition of spaces from $\mathcal{S}_4$ into bouquets of suspended atoms. So we get the following result. We call strings and bands *Baues atoms*.


TABLE 3.
usual strings




decorated strings



and



bands



Theorem 4.1 (Baues–Hennes [8]). *Any polyhedron from $\mathcal{S}_4$ decomposes uniquely into a bouquet of spheres, suspended Moore atoms, suspended Chang atoms and Baues atoms.*


In Section 7 we shall see that actually $\mathcal{S}_4$ is the last case where a “good” description of polyhedra is possible. Starting from $\mathcal{S}_5$ this problem becomes *wild*.

5. TORSION FREE POLYHEDRA. FINITE CASE

Consider now torsion free case. Note that if all $H_k(X)$ are torsion free, the attachment diagram cannot contain “Moore fragments”



In particular, among the atoms from Sections 3 and 4 only Chang atom $C^5 = C^5(\eta)$ and the *double Chang atom* $C_2^7 = C^7(\eta^2)$ with the attachment diagram



are torsion free. Therefore, if we set in Theorem 2.4 $n = 5, m = 4$, the category $\mathcal{A}^0$ consists of bouquets of spheres S^8 and the category $\mathcal{B}^0$ consists of bouquets of spheres S^d ($5 \leq d \leq 8$), suspended Chang atoms C^7, C^8 and suspended double Chang atoms C_2^8 . Obviously, $\mathcal{S}^0(S^8, S^8) = 0$. Easy calculation give the following values of the groups $\Gamma(B) = \mathcal{S}^0(S^8, B)$ for atoms B from $\mathcal{B}^0$:

B	S^5	S^6	S^7	C^7	C^8	C_2^8
Γ	$\mathbb{Z}/24$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/12$	0	$\mathbb{Z}/12$

Morphisms of these spaces induce monomorphisms $\Gamma(S^7) \rightarrow \Gamma(C^7) \rightarrow \Gamma(S^5)$ and $\Gamma(S^6) \rightarrow \Gamma(S^5)$, epimorphism $\Gamma(S^5) \rightarrow \Gamma(C_2^8)$, and isomorphisms $\Gamma(S^7) \rightarrow \Gamma(S^6)$ and $\Gamma(S^7) \rightarrow \Gamma(S^6)$. Thus, an object from $\text{Bim}(\mathcal{S}_{5,4}^0)$ can be presented by a block matrix as in Table 4. Here inside each blocks we have written the groups, wherefrom the coefficients of this block are. The arrows show the allowed transformation between blocks. An integer k in the arrows point out that, when we perform this transformation, the row must be multiplied by k . (No integer means that $k = 1$.) For instance, we can add the rows of the third matrix multiplied by 2 to the rows of the first one. Certainly, compositions of these transformations are also allowed. Thus, for instance, we can add the rows of the third matrix multiplied by 2 to the rows of the second one too. The arising matrix problem is rather simple. It is of finite type, and Table 6 shows the attachment diagrams of the corresponding atoms from $\mathcal{S}_5$. We call them *A-atoms of the 1st kind*. The integer v show, which multiple of the generator of the group $\pi_8^S(S^5) \simeq \mathbb{Z}/24$ is

TABLE 4.





TABLE 5.



used for the “long” attachment. Actually, $1 \leq v \leq 12$ in the case of $A(v)$, $1 \leq v \leq 3$ in the case of $A(\eta v)$, $1 \leq v \leq 6$ in all other cases.

So we have got a description of polyhedra from $\mathcal{T}_5$.

Theorem 5.1 (Baues–Drozd [4]). *Every polyhedron from $\mathcal{T}_5$ is a bouquet of spheres, suspended Chang and double Chang atoms, and the A -atoms of the first kind.*

Analogous is the case of $\mathcal{T}_6$, when we take $m = 4$. We omit details, just schematically presenting the arising matrix problem in Table 6. The dashed line from the 4th to the 6th level show the transformation that only acts on the left-hand column (on $\mathbb{Z}/2$ components). The resulting list of atoms (their attachment diagrams) see in the

TABLE 6.

$\mathbb{Z}/24$	0
$\mathbb{Z}/12$	0
$\mathbb{Z}/12$	0
$\mathbb{Z}/2$	$\mathbb{Z}/2$
0	$\mathbb{Z}/2$
$\mathbb{Z}/2$	$\mathbb{Z}/24$
0	$\mathbb{Z}/12$


Vertical arrows on the right are labeled 2, 6, 12, 6, 2. Vertical arrows on the left are labeled 1, 12. A dashed arrow labeled 12 points from the first module of the top row to the first module of the bottom row.

So we have got a description of polyhedra from $\mathcal{T}_6$.

In the next section we shall use the values of **Hos**-groups between Chang atoms and spheres. To deal with the Chang atom C^5 we apply the bifunctor **Hos** to the cofibration sequence

$$(5.1) \quad S^4 \xrightarrow{\eta} S^3 \rightarrow C^5 \rightarrow S^5 \xrightarrow{\eta} S^4.$$

TABLE 7.



It gives the commutative diagram with exact rows and columns (we write here (X, Y) instead of $\text{Hos}(X, Y)$)

$$\begin{array}{ccccccccc}
\mathbb{Z} & \longrightarrow & \mathbb{Z}/2 & \longrightarrow & (C^5, S^4) & \longrightarrow & 0 & \longrightarrow & \mathbb{Z} \\
\downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
\mathbb{Z}/2 & \longrightarrow & \mathbb{Z}/2 & \longrightarrow & (C^5, S^4) & \longrightarrow & \mathbb{Z} & \longrightarrow & \mathbb{Z}/2 \\
\downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
(S^4, C^5) & \longrightarrow & (S^5, C^5) & \longrightarrow & (C^5, C^5) & \longrightarrow & (S^3, C^5) & \longrightarrow & (S^4, C^5) \\
\downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & \mathbb{Z} & \longrightarrow & (C^5, S^5) & \longrightarrow & 0 & \longrightarrow & 0 \\
\downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
\mathbb{Z} & \longrightarrow & \mathbb{Z}/2 & \longrightarrow & (C^5, S^4) & \longrightarrow & 0 & \longrightarrow & \mathbb{Z},
\end{array}$$

where all maps $\mathbb{Z} \rightarrow \mathbb{Z}/2$ are surjective and all maps $\mathbb{Z}/2 \rightarrow \mathbb{Z}/2$ are bijective. It gives the following values of **Hos**-groups:

$$\begin{aligned}\text{Hos}(C^5, S^4) &= \text{Hos}(S^4, C^5) = 0 \\ \text{Hos}(S^3, C^5) &= \text{Hos}(C^5, S^5) = \mathbb{Z}, \\ \text{Hos}(C^5, S^3) &= \text{Hos}(S^5, C^5) = 2\mathbb{Z}, \\ \text{Hos}(C^5, C^5) &= \mathbb{D},\end{aligned}$$

where $\mathbb{D}$ (the “*dyad*”) is the subrings of $\mathbb{Z} \times \mathbb{Z}$ consisting of all pairs (a, b) with $a \equiv b \pmod{2}$.

Similar observations applied to the suspended versions of the sequence (5.1) and the cofibration sequence

$$S^6 \xrightarrow{\eta^2} S^4 \rightarrow C_2^7 \rightarrow S^7 \xrightarrow{\eta^2} S^5$$

give Table 8 of the values $\text{Hos}(X, Y)$ for suspended atoms from $\mathcal{T}_4$. In

TABLE 8.

	S^4	$C_2^7 : 4$	7	$C^6 : 4$	6	S^5	$C^7 : 5$	7	S^6	S^7
S^4	$\mathbb{Z}$	$2\mathbb{Z}$	$\mathbb{Z}/12$	$2\mathbb{Z}$	0	$\mathbb{Z}/2$	0	$\mathbb{Z}/12$	$\mathbb{Z}/2$	$\mathbb{Z}/24$
$C_2^7 : 4$	$\mathbb{Z}$	$\mathbb{Z}^=$	$\mathbb{Z}/12$	$2\mathbb{Z}$	0	$\mathbb{Z}/2$	0	$\mathbb{Z}/12$	0	$\mathbb{Z}/12$
7	0	0	$\mathbb{Z}^=$	0	0	0	0	$\mathbb{Z}$	0	$2\mathbb{Z}$
$C^6 : 4$	$\mathbb{Z}$	$\mathbb{Z}$	$\mathbb{Z}/12$	$\mathbb{Z}^=$	0	0	0	$\mathbb{Z}/12$	0	$2\mathbb{Z}$
6	0	0	0	0	$\mathbb{Z}^=$	0	0	0	$2\mathbb{Z}$	0
S^5	0	0	0	0	0	$\mathbb{Z}$	$2\mathbb{Z}$	0	$\mathbb{Z}/2$	$\mathbb{Z}/2$
$C^7 : 5$	0	0	0	0	0	$\mathbb{Z}$	$\mathbb{Z}^=$	0	0	0
7	0	0	$2\mathbb{Z}$	0	0	0	0	$\mathbb{Z}^=$	0	$2\mathbb{Z}$
S^6	0	0	$\mathbb{Z}/2$	0	$\mathbb{Z}$	0	0	0	$\mathbb{Z}$	$\mathbb{Z}/2$
S^7	0	0	$\mathbb{Z}$	0	0	0	0	$\mathbb{Z}$	0	$\mathbb{Z}$

this table the **Hos**-groups for suspended Chang atoms are presented in matrix form, emphasizing which components have come from the cells of given dimensions. The superscripts $=$ show that the diagonal parts of the corresponding matrices are with entries not from $\mathbb{Z} \times \mathbb{Z}$, but from

$\mathbb{D}$. For instance, $\text{Es}(C_2^7)$ is presented as the ring of triangular matrices $\begin{pmatrix} a & b \\ 0 & c \end{pmatrix}$, where $a, c \in \mathbb{Z}$, $a \equiv c \pmod{2}$, $b \in \mathbb{Z}/12$. Under such presentation the multiplication of morphisms turns into the multiplication of matrices.

6. TORSION FREE POLYHEDRA. TAME CASE

The category $\mathcal{T}_7$ is more complicated. To describe it, we use Theorem 2.4 with $n = 7, m = 3$. Then $\mathcal{A}^0$ consists of the bouquets of spheres S^d ($10 \leq d \leq 12$) and suspended Chang atoms C^{12} , while $\mathcal{B}^0$ consists of bouquets of spheres S^d ($7 \leq d \leq 10$) suspended Chang atoms C^9, C^{10} and suspended double Chang atoms C_2^{10} . The calculations similar to those of the end of preceding section give Table 9 of the values of groups $\mathcal{S}^0(X, Y)$ for the suspended atoms $X \in \mathcal{A}^0$ and $Y \in \mathcal{B}^0$, also presented in matrix form. The superscripts * show

TABLE 9.

	S^{10}	S^{11}	S^{12}	$C^{12} : 10$	12
S^7	$\mathbb{Z}/24$	0	0	$\mathbb{Z}/24$	0
$C_2^{10} : 7$	0	0	$\mathbb{Z}/2$	$\mathbb{Z}/24^*$	0
10	0	0	$\mathbb{Z}/2$	0	$\mathbb{Z}/2^*$
$C^9 : 7$	$\mathbb{Z}/12$	0	0	$\mathbb{Z}/24^*$	0
9	0	0	$\mathbb{Z}/24$	0	$\mathbb{Z}/2^*$
S^8	$\mathbb{Z}/2$	$\mathbb{Z}/24$	0	0	0
$C^{10} : 8$	0	$\mathbb{Z}/12$	0	0	0
10	0	0	0	0	0
S^9	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/24$	0	$\mathbb{Z}/12$
S^{10}	0	$\mathbb{Z}/2$	$\mathbb{Z}/2$	0	0

that in the corresponding groups we identify the elements of order 2. So actually, these values are isomorphic to $\mathbb{Z}/24$, but it is convenient to consider them as $(\mathbb{Z}/24 \oplus \mathbb{Z}/2)/(12, 1)$. Then again the action of

morphisms from $\mathcal{A}^0$ and $\mathcal{B}^0$, as presented in Table 8 (or, rather, its suspended version) turns into the multiplication of matrices. Again we obtain a bimodule problem close to that of bunches of chains, especially, in its “*decorated*” version (see [17]). To present the answer (for details see [16]), we introduce the following notations and definitions.

Definition 6.1. (1) We consider chains $\mathfrak{E}_k$ and $\mathfrak{F}_k$ ($1 \leq k \leq 4$):

$$\begin{aligned} \mathfrak{E}_1 &= \{e_1 < e_2 < e_4\}, & \mathfrak{F}_1 &= \{f_4 < f_1\}, \\ \mathfrak{E}_2 &= \{e_5 < e_9\}, & \mathfrak{F}_2 &= \{f_3 < f_5\}, \\ \mathfrak{E}_3 &= \{e_6 < e_7\}, & \mathfrak{F}_3 &= \{f_2\}, \\ \mathfrak{E}_4 &= \{e_3 < e_{10} < e'_9 < e'_6\}, & \mathfrak{F}_4 &= \{f'_1 < f'_2 < f'_3\}. \end{aligned}$$

Actually, the elements e_i (f_j) correspond to the rows (columns) of Table 9, while the relations $\overset{c}{-}$ correspond to the elements of the groups $\mathcal{C}^0(A, B)$. We need extra elements e'_i and f'_j since the entries $\mathbb{Z}/2$ in this table behave in a different way than the other ones.

We set $\mathfrak{E} = \bigcup_{i=1}^4 \mathfrak{E}_i$, $\mathfrak{F} = \bigcup_{i=1}^4 \mathfrak{F}_i$, $\mathfrak{X} = \mathfrak{E} \cup \mathfrak{F}$. $x \approx y$ means that x and y belong to the same set $\mathfrak{E}_i$ or $\mathfrak{F}_i$.

- (2) We define symmetric relations $\sim$ and $\overset{c}{-}$ on $\mathfrak{X}$ setting $x \sim y$ if $x \in \mathfrak{E}_i$, $y \in \mathfrak{F}_i$ or vice versa; $e_i \sim e'_i$ ($i \in \{6, 9\}$), $f_j \sim f'_j$ ($1 \leq j \leq 3$). We also define the symmetric relations $\overset{c}{-}$, where $c \in \{1, 2, 3, 4, 6\}$, setting $e_i \overset{c}{-} f_j$ if $e_i - f_j$ and the (ij) -th entry in Table 9 is $\mathbb{Z}/m$ with $c \mid m$. We denote by R the set of all relations $\{\sim, \overset{c}{-}\}$ and by $v(c)$ the biggest d such that 2^d divides c .
- (3) We define a *word* as a sequence $w = x_1 r_2 x_2 r_3 \dots r_l x_l$ where $x_i \in \mathfrak{X}$, $r_i \in R$ such that
- (a) $x_{k-1} r_k x_k$ in $\mathfrak{X}$ for each $1 < k \leq l$;
 - (b) if $r_k = \sim$, then $r_{k+1} = \overset{c}{-}$ and vice versa;
 - (c) if $r_2 = \overset{c}{-}$ (respectively, $r_l = \overset{c}{-}$), there is no element $y \in \mathfrak{X}$ such that $x_1 \sim y$ (respectively, $x_l \sim y$);
 - (d) if $r_k = \overset{c}{-}$ with $v(c) = 1$, then either $2 < k < l$, or $k = 2$, $x_1 = e_1$, or $k = l$, $x_l = x_1$;

- (e) if $\mathbf{r} = \overset{c}{-}$ with $v(c) = 2$, then $\mathbf{r}$ can only occur in the following words or their reverse:

$$\begin{aligned}
& e_4 \sim e_5 \mathbf{r} f_3 \sim \dots \quad (\text{of any length}), \\
& e_1 \mathbf{r} f_4 \sim f_5, \quad e_3 \sim e_2 \mathbf{r} f_4 \sim f_5, \\
& \dots \overset{c'}{-} e_4 \sim e_5 \mathbf{r} f_3 \sim \dots \quad (\text{of any length}), \\
& e_1 \mathbf{r} f_4 \sim f_5 \overset{c'}{-} \dots \quad (\text{of any length}), \\
& e_3 \sim e_2 \mathbf{r} f_4 \sim f_5 \overset{c'}{-} \dots \quad (\text{of any length}), \\
& e_1 \mathbf{r} f_4 \sim f_5 \overset{c'}{-} \dots \quad (\text{of any length}), \\
& e_1 \mathbf{r} f_1 \sim f'_1, \quad e'_6 \sim e_6 \mathbf{r} f_2 \sim f'_2, \quad e'_9 \sim e_9 \mathbf{r} f_3 \sim f'_3,
\end{aligned}$$

where $c' \equiv 0 \pmod{3}$;

- (f) if w contains a subword $e_i \overset{c}{-} f_j$, $c \in \{3, 9\}$ or its reverse, it does not contain any subword $e_{i'} \overset{c'}{-} f_{j'}$, $c' \not\equiv 0 \pmod{3}$, $e_i \approx e_{i'}$ (equivalently, $f_j \approx f_{j'}$) or its reverse.

Here the *reverse* to the word w is the word $w^* = x_l r_l x_{l-1} \dots x_2 r_2 x_1$.

We call l the *length* of the word w .

- (4) We define a *cycle* as a pair $z = (w, r_1)$, where w is a word such that $r_2 = r_l = \sim$ and $r_k \neq \overset{c}{-}$ with $v(c) = 2$, while $r_1 = \overset{c}{-}$ with $v(c) \neq 2$ and $x_l r_1 x_1$ in $\mathfrak{X}$. For such a cycle we set $x_{ql+k} = x_k$ and $r_{ql+k} = r_k$ for any q and $1 \leq k \leq l$.
- (5) The m -th *shift* of the cycle $z = (w, r_1)$ is defined as the cycle $z^{(m)} = (w^{(m)}, r_{2m+1})$, where $w^{(m)} = x_{2m+1} r_{2m+2} x_{2m+2} \dots r_{2m} x_{2m}$.
- (6) A cycle (w, r_1) is called *periodic* if w is of the form $w = v r_1 v r_1 \dots r_1 v$ for a shorter cycle (v, r_1) .
- (7) We call two words, w and $w' = x_1 r'_2 x_2 r'_3 \dots r'_l x_l$ (with the same x_k), *elementary congruent* if there are two indices k_1, k_2 such that

$$\begin{aligned}
& r_{k_1} = \overset{3c}{-}, \quad r_{k_2} = \overset{d}{-} \quad \text{for some } c \neq 3, d \neq 3, \\
& r'_{k_1} = \overset{c}{-}, \quad r'_{k_2} = \overset{3d}{-}, \\
& r'_k = r_k \quad \text{for } k \notin \{k_1, k_2\}, \\
& x_{k_1} \approx x_{k_2} \quad \text{or } x_{k_1} \approx x_{k_2-1}.
\end{aligned}$$

- (8) We call two words w, w' *congruent* and write $w \equiv w'$ if there is a sequence of words $w = w_1, w_2, \dots, w_n = w$ such that w_k and w_{k+1} are elementary congruent for $1 \leq k < n$. We call two cycles $z = (w, r_1)$ and $z' = (w', r'_1)$ *congruent* and write $z \equiv z'$ if $w' \equiv z$ and $r'_1 = r_1$.

We recall that two polyhedra X, Y are called *congruent* if $X \vee Z \simeq Y \vee Z$ for some polyhedron Z . Then we write $X \equiv Y$.

- Theorem 6.2.** (1) Every word w defines an indecomposable polyhedron $P(w)$ from $\mathcal{T}_7$, called string polyhedron.
- (2) Let $\pi(t) \neq t$ be an irreducible polynomial over the field $\mathbb{Z}/2$. Every triple $(z, \pi(t), m)$, where z is a non-periodic cycle and $m \in \mathbb{N}$, defines an indecomposable polyhedron $P(z, \pi, m)$ from $\mathcal{T}_7$, called band polyhedron.
- (3) Every indecomposable polyhedron from $\mathcal{T}_7$ is congruent either to a string or to a band one.
- (4) $P(w) \equiv P(w')$ if and only if either $w' \equiv w$ or $w' \equiv w^*$.
- (5) $P(z, \pi(t), m) \equiv P(z', \pi'(t), m)$ if and only if $m = m'$ and one of the following possibilities hold:
- (a) $\pi'(t) = \pi(t)$ and either $z' \equiv z^{(k)}$ with k even or $z' = z^{*(k)}$ with k odd;
 - (b) $\pi'(t) = t^d \pi(1/t)$, where $d = \deg \pi$, and either $z' = z^{(k)}$ with k odd or $z' = z^{*(k)}$ with k even.
- (6) Neither string polyhedron is congruent to a band one.
- (7)

The cofibration sequence

$$A \xrightarrow{f} B \rightarrow Cf \rightarrow SA, \quad A \in \mathcal{A}^0, \quad B \in \mathcal{B}^0,$$

and the attachment diagram of a string polyhedron $P(w)$ is constructed as follows.

- (1) The indecomposable summands of A correspond to the following subwords of w (or their reverse):

$$\begin{array}{lll} S^{10} & \text{to} & f_1 \sim f'_1, \\ S^{11} & \text{to} & f_2 \sim f'_2, \\ S^{12} & \text{to} & f_3 \sim f'_3, \\ C^{12} & \text{to} & f_4 \sim f_5. \end{array}$$

- (2) The indecomposable summands of B correspond to the following subwords of w (or their reverse):


$$\begin{array}{lll} S^7 & \text{to} & e_1, \\ C_2^{10} & \text{to} & e_2 \sim e_3, \\ C^9 & \text{to} & e_4 \sim e_5, \\ S^8 & \text{to} & e_6 \sim e'_6, \\ C^{10} & \text{to} & e_7, \\ S^9 & \text{to} & e_9 \sim e'_9, \\ S^{10} & \text{to} & e_{10}. \end{array}$$

- (3) The attachments correspond to the subwords $e_i \xrightarrow{c} f_j$ (or their reverse). Namely, such an attachment starts at the f -end of the corresponding subword and ends at its e -end; the number c shows which multiple of the generator of the (ij) -th group from Table 9 must be taken.

For instance, if

$$\begin{aligned} w &= e_{10} \xrightarrow{1} f'_2 \sim f_2 \xrightarrow{8} e_6 \sim e'_6 \xrightarrow{1} f'_1 \sim f_1 \xrightarrow{2} e_4 \\ &\sim e_5 \xrightarrow{6} f_5 \sim f_4 \xrightarrow{1} e_2 \sim e_3 \xrightarrow{1} f'_3 \sim f_3 \xrightarrow{2} e_5 \\ &\sim e_4 \xrightarrow{3} f_1 \sim f'_1 \xrightarrow{1} e'_9 \sim e_9 \xrightarrow{12} f_3 \sim f'_3, \end{aligned}$$

the polyhedron $P(w)$ has the attachment diagram




Let now $P(z, \pi(t), m)$ be a band polyhedron. Replacing w by w^* , we may suppose that $x_1 \in \mathfrak{E}$, $x_n \in \mathfrak{F}$. Let also Φ be the Frobenius matrix with the characteristic polynomial $\pi(t)^m$. Then the cofibration sequence and the attachment diagram are constructed as follows.

- (1) Do the construction as above for the word w .
- (2) Replace every summand A_j of A and every summand B_i of B by m copies of it, $A_{j1}, \dots, A_{jm}$ and $B_{i1}, \dots, B_{im}$.
- (3) If there was an attachment $A_j \xrightarrow{c} B_i$, replace it by the attachments $A_{jk} \xrightarrow{c} B_{ik}$ ($1 \leq k \leq m$).
- (4) If A_j is the last summand of A , B_i is the first summand of B and $r_1 = \frac{c}{-}$, add new attachments $A_{jk} \xrightarrow{c} B_{il}$ in all cases, when the (lk) -th coefficient of the matrix Φ is nonzero.


For instance, consider the band polyhedron $P(z, t^2 + t + 1, 3)$ $z = (w, \frac{1}{-})$, where

$$w = e_2 \sim e_3 \xrightarrow{1} f'_3 \sim f_3 \xrightarrow{2} e_9 \sim e'_9 \xrightarrow{1} f'_1 \sim f_1 \xrightarrow{3} e_4 \sim e_5 \xrightarrow{6} f_5 \sim f_4.$$


Then the attachment diagram is



Here the double lines show the attachments like



while the wavy line shows the attachment



ruled by the Frobenius matrix with the characteristic polynomial $\pi(t)^2 = t^4 + t^2 + 1$, namely,

$$\begin{pmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

7. WILD CASES

Since we are dealing with additive categories that are not categories over a field, we have to precise the notion of wildness. The following one seems to work in all known cases.

Definition 7.1. We call an additive category $\mathcal{C}$ *wild* if, there is a field $\mathbb{k}$ such that for every finitely generated $\mathbb{k}$ -algebra Λ there is a full subcategory $\mathcal{C}_\Lambda \subseteq \mathcal{C}$ and an epivalence $\mathcal{C}_\Lambda \rightarrow \Lambda\text{-mod}$ (the category of Λ -modules that are finite dimensional over $\mathbb{k}$).

One can see that for algebras over a field this definition is equivalent to the usual one (see, for instance [14]). One can also easily show that if a category $\mathcal{D}$ is wild and there is an epivalence $\mathcal{C}' \rightarrow \mathcal{D}$ for a full subcategory $\mathcal{C}' \subseteq \mathcal{C}$, then $\mathcal{C}$ is wild as well.

Now we present the results on wildness of categories $\mathcal{S}_n$ and $\mathcal{T}_n$.

Theorem 7.2 (Baues [6]). *If $n > 4$, the category $\mathcal{S}_n$ is wild.*

Proof. Obviously, one only has to prove the claim for $n = 5$. The category $\mathcal{S}_5$ contains the full subcategory $\mathcal{C} = \mathcal{A} \uparrow \mathcal{B}$, where $\mathcal{A}$ consists of bouquets of suspended Moore atom $A = M^6(2)$ and $\mathcal{B}$ consists of bouquets of suspended Moore atoms $B = M^8(2)$. Let $\mathcal{V} = \mathcal{A} \mathcal{S}_\mathcal{B}$. Since

$\text{Hos}(B, A) = 0$, Corollary 1.2 is applicable. Moreover, the ideal $\mathcal{I}$ in this case is zero, so $\mathcal{C}/\mathcal{I} \simeq \text{Bim}(\mathcal{V})$ with $\mathcal{I}^2 = 0$, hence, the natural functor $\mathcal{C} \rightarrow \text{Bim}(\mathcal{V})$ is an epivalence.

Consider the cofibration sequence

$$(7.1) \quad S^7 \xrightarrow{2} S^7 \rightarrow A \rightarrow S^8 \xrightarrow{2} S^8.$$

Apply to it the functors $\text{Hos}(-, S^6)$ and $\text{Hos}(-, S^5)$. Taking into account the Hopf map $\eta : S^6 \rightarrow S^5$ we get the commutative diagram with exact rows

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathbb{Z}/2 & \longrightarrow & \text{Hos}(A, S^6) & \longrightarrow & \mathbb{Z}/2 & \longrightarrow & 0 \\ & & \eta_* \downarrow & & \downarrow & & \downarrow \wr & & \\ 0 & \longrightarrow & \mathbb{Z}/2 & \longrightarrow & \text{Hos}(A, S^5) & \longrightarrow & \mathbb{Z}/2 & \longrightarrow & 0. \end{array}$$

Since $\eta^3 = 4\nu$, where ν is the element of order 8 in $\text{Hos}(S^8, S^5)$, the map η_* in this diagram is zero, therefore, the lower exact sequence splits and $\text{Hos}(A, S^5) \simeq \mathbb{Z}/2 \oplus \mathbb{Z}/2$. Quite similarly, one shows that $\text{Hos}(S^8, B) \simeq \mathbb{Z}/2 \oplus \mathbb{Z}/2$. Now apply the functors $\text{Hos}(-, S^5)$ and $\text{Hos}(-, B)$ to the exact sequence (7.1) and take into account the map $S^5 \rightarrow B$ from the definition of $B = M^6(2)$. Since $\text{Hos}(S^7, B) \simeq \mathbb{Z}/2$, we get the commutative diagram with exact rows

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathbb{Z}/2 & \longrightarrow & \text{Hos}(A, S^5) & \longrightarrow & \mathbb{Z}/2 & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow \wr & & \\ 0 & \longrightarrow & \mathbb{Z}/2 \oplus \mathbb{Z}/2 & \longrightarrow & \text{Hos}(A, B) & \longrightarrow & \mathbb{Z}/2 & \longrightarrow & 0. \end{array}$$

We know that the upper row of this diagram splits. Hence, the lower row splits too, so $\text{Hos}(A, B) \simeq (\mathbb{Z}/2)^3$. Recall that $\text{Es}(A) \simeq \text{Es}(B) \simeq \mathbb{Z}/4$ (Proposition 3.3). Hence, there is an epivalence $\text{Bim}(\mathcal{V}) \rightarrow \Lambda\text{-mod}$, where Λ is the path algebra of the quiver $\bullet \rightrightarrows \bullet$ over the field $\mathbb{Z}/2$. The latter is well-known to be wild, therefore, so is also $\mathcal{S}_5$. $\square$

Theorem 7.3 ([16]). *The category $\mathcal{T}_n$ is wild for $n > 7$.*

Proof. Again we only have to prove it for $n = 8$. The category $\mathcal{T}_8$ contains the full subcategory $\mathcal{C} = \mathcal{A} \uparrow_{\mathcal{V}} \mathcal{B}$, where $\mathcal{A}$ consists of bouquets of Chang atoms C^1_4 and $\mathcal{B}$ consists of bouquets of spheres $\mathbb{S}^8$ and S^1 , and $\mathcal{V} = \mathcal{S}_{8,3}^0 \mathcal{A}$. Moreover, $\mathcal{S}_{8,3}^0 \cap \text{Bim } \mathcal{V} = 0$, so there is an epivalence $\mathcal{C} \rightarrow \text{Bim}(\mathcal{V})$. Consider the cofibration sequence

$$S^{13} \xrightarrow{\eta^2} S^{11} \rightarrow C_2^{14} \rightarrow S^{14} \rightarrow S^{12}$$

and apply to it the functor $\text{Hos}(-, S^{11})$. We get the exact sequence

$$\mathbb{Z}/2 \xrightarrow{(\eta^2)^*} \mathbb{Z}/24 \rightarrow \text{Hos}(C_2^{14}, S^{11}) \rightarrow \mathbb{Z} \rightarrow \mathbb{Z}/2,$$

wherefrom $\mathcal{S}^0(C_2^{14}, S^1) \simeq \mathbb{Z}/12$. Moreover, there is a commutative diagram of cofibration sequences

$$\begin{array}{ccccccccc}
 S^{13} & \xrightarrow{\eta} & S^{12} & \longrightarrow & C^{14} & \longrightarrow & S^{14} & \xrightarrow{\eta} & S^{13} \\
 \text{Id } nolimits \downarrow & & \eta \downarrow & & \downarrow & & \downarrow \text{Id } nolimits & & \downarrow \eta \\
 S^{13} & \xrightarrow{\eta^2} & S^{11} & \longrightarrow & C_2^{14} & \longrightarrow & S^{14} & \xrightarrow[\eta^2]{} & S^{12}.
 \end{array}$$

Applying the functor $\text{Hos}(-, S^8)$, we get the commutative diagram with exact rows

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \mathbb{Z}/2 & \longrightarrow & \text{Hos}(C_2^{14}, S^8) & \longrightarrow & \mathbb{Z}/24 \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & \mathbb{Z}/2 & \longrightarrow & \text{Hos}(C^{14}, S^8) & \longrightarrow & 0 \longrightarrow 0.
 \end{array}$$

(Recall that $\pi_{d+4}^S(S^d) = \pi_{d+5}^S(S^d) = 0$ and $\pi_{d+6}^S(S^d) = \mathbb{Z}/2$ [24]). Therefore, $\mathcal{S}^0(C_2^{14}, S^8) \simeq \mathbb{Z}/24 \oplus \mathbb{Z}/2$. So we present maps $a \in \mathcal{V}(A, B)$, where $A \in \mathcal{A}$, $B \in \mathcal{B}$, as block-triangular matrices

$$a = \begin{pmatrix} a_1 & a_2 \\ 0 & a_3 \end{pmatrix},$$

where a_1 is with the coefficients from $\mathbb{Z}/24$, a_2 is with coefficients from $\mathbb{Z}/2$ and a_3 with coefficients from $\mathbb{Z}/12$. On the other hand, maps $\alpha : A \rightarrow A'$, where $A, A' \in \mathcal{A}$, and $\beta : B \rightarrow B'$, where $B, B' \in \mathcal{B}$ can be presented by block-triangular matrices

$$\alpha = \begin{pmatrix} \alpha_1 & \alpha_2 \\ 0 & \alpha_3 \end{pmatrix} \quad \text{and} \quad \beta = \begin{pmatrix} \beta_1 & \beta_2 \\ 0 & \beta_3 \end{pmatrix},$$

where α_2 has coefficients from $\mathbb{Z}/12$, β_2 has coefficients from $\mathbb{Z}/24$, other blocks have components from $\mathbb{Z}$ and $\alpha_1 \equiv \alpha_3 \pmod{2}$.

We consider the full subcategory $\mathcal{C} \subset \text{Bim}(\mathcal{V})$ consisting of all maps a such that the corresponding blocks a_1, a_2, a_3 are of the form

$$a_1 = \begin{pmatrix} 6I & 0 & 0 \\ 0 & 12 & 0 \end{pmatrix}, \quad a_2 = \begin{pmatrix} 0 & I & 0 \\ 0 & 0 & u \end{pmatrix} \quad a_3 = \begin{pmatrix} 6v_1 & 6v_2 & 0 \end{pmatrix},$$

where the entries I stand for identity matrices (not necessary of the same dimensions) and u, v_1, v_2 are arbitrary matrices with coefficients from $\mathbb{Z}/2$ of proper sizes. We write $a = a(u, v_1, v_2)$. One can verify that if (α, β) is a morphism $a(u, v_1, v_2) \rightarrow a(u', v'_1, v'_2)$, there are integral matrices $\gamma_1, \gamma_2, \gamma_3$ such that $v_i \gamma_1 = \gamma_2 v_i$ ($i = 1, 2$) and $u \gamma_3 = \gamma_1 u$. Conversely, any given triple $\gamma_1, \gamma_2, \gamma_3$ with these properties can be accomplished to a morphism $a(u, v_1, v_2) \rightarrow a(u', v'_1, v'_2)$. It gives rise to an epivalence $\mathcal{C} \rightarrow \Lambda\text{-mod}$, where Λ is the path algebra of the quiver $\bullet \longrightarrow \bullet \rightleftarrows \bullet$. It is known to be wild. Therefore, $\mathcal{T}_8$ is wild as well. $\square$

REFERENCES

- [1] H. Bass. Algebraic K -theory. W. A. Benjamin, Inc. New York–Amsterdam, 1968.
- [2] H.-J. Baues. Homotopy types. Handbook of Algebraic Topology. Elsevier, Amsterdam, 1995, 1–72.
- [3] H.-J. Baues. Atoms in Topology. Jahresber. Dtsch. Math.-Ver. **104** (2002), 147–164.
- [4] H.-J. Baues and Y. A. Drozd. The homotopy classification of $(n-1)$ -connected $(n+4)$ -dimensional polyhedra with torsion free homology. Expo. Math. **17** (1999), 161–179.
- [5] H.-J. Baues and Y. A. Drozd. Representation theory of homotopy types with at most two non-trivial homotopy groups. Math. Proc. Cambridge Phil. Soc. **128** (2000), 283–300.
- [6] H.-J. Baues and Y. A. Drozd. Indecomposable homotopy types with at most two non-trivial homology groups. Groups of Homotopy Self-Equivalences and Related Topics. Contemporary Mathematics, **274** (2001), 39–56.
- [7] H.-J. Baues and Y. A. Drozd. Classification of stable homotopy types with torsion-free homology. Topology **40** (2001), 789–821.
- [8] H. J. Baues and M. Hennes. The homotopy classification of $(n-1)$ -connected $(n+3)$ -dimensional polyhedra, $n \geq 4$. Topology, **30** (1991), 373–408.
- [9] V. M. Bondarenko. Representations of bundles of semichained sets and their applications. St. Petersburg Math. J. **3** (1992), 973–996.
- [10] I. Burban and Yu. Drozd. Derived categories of nodal algebras. J. Algebra **272** (2004), 46–94, arXiv:math/0307060.
- [11] S. C. Chang. Homotopy invariants and continuous mappings. Proc. R. Soc. London **202** (1950), 253–263.
- [12] J. M. Cohen. Stable Homotopy. Lecture Notes in Math. **165** (1970).
- [13] Y. A. Drozd. Adèles and integral representations. Izv. Akad. Nauk SSSR, Ser. Mat. **33** (1969), 1080–1088. (Math. USSR, Izv. **3** (1969), 1019–1026.)
- [14] Y. A. Drozd. Reduction algorithm and representations of boxes and algebras. Comptes Rendue Math. Acad. Sci. Canada, **23** (2001), 97–125.
- [15] Y. A. Drozd. Matrix problems and stable homotopy types of polyhedra. Central European J. Math. **2** (2004), 420–447, arXiv:math/0404252.
- [16] Y. A. Drozd. On classification of torsion free polyhedra. Preprint MPI 2005-99. Max-Planck-Institut für Mathematik, 2005.
- [17] O. Y. Drozd-Koroleva. Representations of decorated bunches of chains. Preprint. Kiev, 2009.
- [18] R. Hartshorn. Residues and Duality. Lecture Notes in Math. **20** (1966).
- [19] H.-W. Henn. Classification of p -local low-dimensional spectra. J. Pure Appl. Algebra **45** (1987), 45–71.
- [20] S.-T. Hu. Homotopy Theory. Academic Press. New York–London, 1959.
- [21] H. Jacobinski. Genera and decompositions of lattices over orders. Acta Math. **121** (1968), 1–29.
- [22] D. Puppe. On the structure of stable homotopy theory. Colloquium on Algebraic Topology. Aarhus Universitet Matematisk Institut, 1962, 65–71.
- [23] R. M. Switzer. Algebraic Topology — Homotopy and Homology. Springer-Verlag. Berlin–Heidelberg–New York, 1975.
- [24] H. Toda. Composition Methods in the Homotopy Groups of Spheres. Ann. Math. Studies, Vol. 49, Princeton, 1962.
- [25] J. H. C. Whitehead. The homotopy type of a special kind of polyhedron. Ann. Soc. Polon. Math. **21** (1949), 176–186.

INSTITUTE OF MATHEMATICS, NATIONAL ACADEMY OF SCIENCES OF UKRAINE,
TERESCHENKIVSKA 3, 01601 KIEV, UKRAINE

E-mail address: drozd@imath.kiev.ua, y.a.drozd@gmail.com

URL: www.imath.kiev.ua/~drozd