

Koszul duality for extension algebras of standard modules

Yuriy Drozd and Volodymyr Mazorchuk

Abstract

We define and investigate a class of Koszul quasi-hereditary algebras for which there is a natural equivalence between the bounded derived category of graded modules and the bounded derived category of graded modules over (a proper version of) the extension algebra of standard modules. Examples of such algebras include, in particular, the multiplicity free blocks of the BGG category \mathcal{O} , and some quasi-hereditary algebras with Cartan decomposition in the sense of König.

1 Introduction

For a finite-dimensional Koszul algebra, A , of finite global dimension there is a natural equivalence between the bounded derived category $\mathcal{D}^b(A\text{-gmod})$ of graded A -modules and the bounded derived category of graded modules over the Yoneda extension algebra $E(A)$ of A , see [BGS]. This equivalence is produced by the so-called Koszul duality functor. If A is quasi-hereditary and satisfies some natural assumptions on the resolutions of standard and costandard modules (see [ADL]), then the algebra $E(A)$ is also quasi-hereditary and the Koszul duality functor behaves well with respect to this structure. Some time ago S. Ovsienko in a private communication expressed a hope that for (some) graded Koszul quasi-hereditary algebras it might be possible that $\mathcal{D}^b(A\text{-gmod})$ is equivalent to the bounded derived category of graded modules for the extension algebra $\text{Ext}_A^*(\Delta, \Delta)$ of the direct sum Δ of all standard modules for A . The reason for this hope is the fact that every quasi-hereditary algebra has two natural families of homologically orthogonal modules, namely standard and costandard modules. Both these families generate $\mathcal{D}^b(A\text{-gmod})$ as a triangulated category. The idea of Ovsienko was to organize the equivalence between the derived categories such that the standard A -modules become projective objects and the corresponding costandard A -modules become simple objects. In particular, it should follow automatically that $\text{Ext}_A^*(\Delta, \Delta)$ is Koszul, and its Koszul dual should be isomorphic to the extension algebra $\text{Ext}_A^*(\nabla, \nabla)$ of the costandard module ∇ for A .

In the present paper we define and investigate a big family of graded quasi-hereditary algebras for which Ovsienko's idea works. However, the passage from A to $\text{Ext}_A^*(\Delta, \Delta)$ is not painless. There is of course a trivial case, when A is directed. In this case we have either $A \cong \text{Ext}_A^*(\Delta, \Delta)$ or $E(A) \cong \text{Ext}_A^*(\Delta, \Delta)$. In all other cases one quickly comes to

the problem that the “natural” gradation induced on $\text{Ext}_A^*(\Delta, \Delta)$ from $\mathcal{D}^b(A\text{-gmod})$ is a \mathbb{Z}^2 -gradation and not a \mathbb{Z} -gradation. In fact, we were not able to find any “natural” copy of the category of graded $\text{Ext}_A^*(\Delta, \Delta)$ -modules inside $\mathcal{D}^b(A\text{-gmod})$. However, under special conditions (I)-(IV) see Subsection 2.5, which we impose on the algebras we consider in this paper, we single out inside $\mathcal{D}^b(A\text{-gmod})$ a subcategory of graded modules over certain \mathbb{Z} -graded category (not algebra!), \mathcal{B} , whose bounded derived category is naturally equivalent to $\mathcal{D}^b(A\text{-gmod})$. Additionally, the category \mathcal{B} carries a natural free action of \mathbb{Z} . The quotient modulo this action happens to be exactly $\text{Ext}_A^*(\Delta, \Delta)$ with the induced \mathbb{Z}^2 -gradation. As a consequence, we have to extend our setup and consider modules over categories rather than those over algebras. This forces us to reformulate and extend many classical notions and results (like Koszul algebras, quasi-hereditary algebras, Rickard-Morita Theorem etc.) in our more general setup.

The paper is organized as follows: in Section 2 we collect all necessary preliminaries about the categories and algebras we consider. In Section 3 we get some preliminary information about the quasi-hereditary categories satisfying (I)-(IV). In Section 4 we formulate and prove our main result. We finish the paper with a discussion on several applications of our result in Section 5.

2 Some generalities

2.1 Modules over categories and Rickard–Morita Theorem

Let \mathbb{k} be an algebraically closed field and $D = \text{Hom}_{\mathbb{k}}(-, \mathbb{k})$ denote the usual duality. Since we need not only modules over algebras, but also those over categories, we include the main definitions concerning them. All categories under consideration will be *linear \mathbb{k} -categories*. It means that all sets of morphisms $\mathcal{A}(x, y)$ in such a category, \mathcal{A} , are \mathbb{k} -vector spaces and the multiplication is \mathbb{k} -bilinear. Moreover, we suppose that these categories are *small*, i.e. their classes of objects are sets. All functors are supposed to be \mathbb{k} -linear. An *\mathcal{A} -module* is, by definition, a functor $M : \mathcal{A} \rightarrow \mathcal{V}ec$ (the category of \mathbb{k} -vector spaces). For an element, $m \in M(x)$, and a morphism, $\alpha : x \rightarrow y$, we write αm instead of $M(\alpha)m$, etc. We denote by $\mathcal{A}\text{-Mod}$ the category of all \mathcal{A} -modules. A *representable* module is one isomorphic to $\mathcal{A}^x = \mathcal{A}(x, -)$ for some object x . Such functors are projective objects in the category $\mathcal{A}\text{-Mod}$ and every projective object in this category is a direct summand of a direct sum (maybe infinite) of representable functors. Just in the same way, the functors $\mathcal{A}_x = \mathcal{A}(-, x)$ are projective objects in the category of \mathcal{A}^{op} -modules, where \mathcal{A}^{op} denotes the *opposite category*. A *set of generators* of an \mathcal{A} -module, M , is a subset, $S \subseteq \bigcup_{x \in \text{ob } \mathcal{A}} M(x)$, such that any element $m \in M$ can be expressed as $\sum_{u \in S} \alpha_u u$, where all $\alpha_u \in \text{mor } \mathcal{A}$ and only finitely many of these morphisms are nonzero. Especially, $\{1_x\}$ is a set of generators of \mathcal{A}^x , as well as of \mathcal{A}_x .

Recall that, if a category, \mathcal{C} , has infinite direct sums, an object, C , is called *compact* if the functor \mathcal{C}^C preserves arbitrary direct sums. For instance, finitely generated modules are compact objects of $\mathcal{A}\text{-Mod}$. Suppose now that \mathcal{A} is *basic*, i.e. different objects of \mathcal{A} are

non-isomorphic and there are no nontrivial idempotents in all algebras $\mathcal{A}(x, x)$, $x \in \text{ob } \mathcal{A}$. We denote by $\mathcal{A}\text{-mod}$ the category of *finite dimensional* \mathcal{A} -modules, that is those modules M for which all spaces $M(x)$ are finite dimensional and $M(x) = 0$ for all but a finite number of objects x . Equivalently, $\bigoplus_{x \in \text{ob } \mathcal{A}} M(x)$ is finite dimensional. If all modules \mathcal{A}^x and \mathcal{A}_x are finite dimensional, we call \mathcal{A} a *bounded category*.

We denote by $\mathcal{D}\mathcal{A}$ the *derived category* of the category $\mathcal{A}\text{-Mod}$; by $\mathcal{D}^+ \mathcal{A}$, $\mathcal{D}^- \mathcal{A}$ and $\mathcal{D}^b \mathcal{A}$, respectively, its full subcategories consisting of *right bounded*, *left bounded* and (two-sided) bounded complexes. The shift in $\mathcal{D}\mathcal{A}$ will be denoted by $C^\bullet \mapsto C^\bullet[1]$; actually $C^n[1] = C^{n+1}$. By $\mathcal{D}^{\text{per}} \mathcal{A}$ we denote the full subcategory of $\mathcal{D}\mathcal{A}$ consisting of *perfect complexes*, i.e. those isomorphic (in $\mathcal{D}\mathcal{A}$) to bounded complexes of finitely generated projective modules. The perfect complexes are just the compact objects of $\mathcal{D}\mathcal{A}$. The category $\mathcal{D}^{\text{per}} \mathcal{A}$ can be identified with the *bounded homotopy category* $\mathcal{H}^b(\mathcal{A}\text{-proj})$, i.e. the factorcategory of the category of finite complexes of finitely generated projective \mathcal{A} -modules modulo homotopy. The projective modules (or rather their canonical images) generate $\mathcal{D}^{\text{per}} \mathcal{A}$ as a triangulated category. We recall the following theorem by Rickard [Ric], which we present in a slightly more general context of \mathbb{k} -linear categories, see for example [Ke, Corollary 9.2].

Theorem 2.1 (Rickard–Morita Theorem). *Let \mathcal{A} and \mathcal{B} be two small \mathbb{k} -linear categories. Then the following conditions are equivalent:*

1. *There is a triangle equivalence $\mathcal{D}\mathcal{A} \xrightarrow{\sim} \mathcal{D}\mathcal{B}$.*
2. *There is a triangle equivalence $\mathcal{D}^* \mathcal{A} \xrightarrow{\sim} \mathcal{D}^* \mathcal{B}$, where $*$ can be replaced by any of the symbols $+$, $-$, b or per .*
3. *There is a full subcategory $\mathcal{X} \subset \mathcal{D}^{\text{per}} \mathcal{A}$ such that*
 - (a) $\mathcal{X} \simeq \mathcal{B}^{\text{op}}$;
 - (b) $\text{Hom}_{\mathcal{D}\mathcal{A}}(X, X'[k]) = 0$ for any $X, X' \in \mathcal{X}$ and any $k \neq 0$;
 - (c) \mathcal{X} generates $\mathcal{D}^{\text{per}} \mathcal{A}$ as a triangulated category.

Moreover, in this case, any equivalence $T : \mathcal{X} \xrightarrow{\sim} \mathcal{B}^{\text{op}}$ can be extended to a triangle equivalence $F : \mathcal{D}\mathcal{A} \xrightarrow{\sim} \mathcal{D}\mathcal{B}$ such that $FX = \mathcal{B}^{TX}$ for every $X \in \mathcal{X}$. In particular, if $\mathcal{B} = \mathcal{X}^{\text{op}}$, $FX = \mathcal{X}_X = \text{Hom}_{\mathcal{D}\mathcal{A}}(-, X)$.

In fact, given an equivalence, $\Phi : \mathcal{D}\mathcal{B} \xrightarrow{\sim} \mathcal{D}\mathcal{A}$, one can set $\mathcal{X} = \{\Phi \mathcal{B}^x \mid x \in \text{ob } \mathcal{B}\}$. Note that, since \mathcal{B}^{TX} is a finitely generated projective \mathcal{B} -module, then one also has $\text{RHom}_{\mathcal{A}}(X, -) \simeq \text{RHom}_{\mathcal{B}}(\mathcal{B}^{TX}, F_-)$. Thus, for every complex C^\bullet of \mathcal{A} -modules we have $\text{RHom}_{\mathcal{A}}(X, C^\bullet) \simeq FC^\bullet(TX)$ in $\mathcal{D}\mathcal{B}$.

The set of objects of a full subcategory $\mathcal{X} \subseteq \mathcal{D}\mathcal{A}$ satisfying conditions (3b) and (3c) will be called a *tilting subset* in $\mathcal{D}\mathcal{A}$.

2.2 Graded categories, graded modules and group actions

Let \mathbf{G} be a semigroup. A \mathbf{G} -grading of a category, \mathcal{A} , consists of decompositions $\mathcal{A}(x, y) = \bigoplus_{\sigma \in \mathbf{G}} \mathcal{A}(x, y)_\sigma$ given for any objects $x, y \in \mathcal{A}$, such that, for every $x, y, z \in \text{ob } \mathcal{A}$ and for every $\sigma, \tau \in \mathbf{G}$, $\mathcal{A}(y, z)_\tau \mathcal{A}(x, y)_\sigma \subseteq \mathcal{A}(x, z)_{\sigma\tau}$. A category, \mathcal{A} , with a fixed \mathbf{G} -grading is called a \mathbf{G} -graded category. The morphisms $\alpha \in \mathcal{A}(x, y)_\sigma$ are called *homogeneous of degree* σ , and we shall write $\text{deg } \alpha = \sigma$. If \mathcal{A} is a \mathbf{G} -graded category, a \mathbf{G} -graded module (or simply a *graded module*) over \mathcal{A} is an \mathcal{A} -module, M , with fixed decompositions $M(x) = \bigoplus_{\sigma \in \mathbf{G}} M(x)_\sigma$, given for all objects $x \in \mathcal{A}$, such that $\mathcal{A}(x, y)_\tau M(x)_\sigma \subseteq M(y)_{\sigma\tau}$ for any x, y, σ, τ . We denote by $\mathcal{A}\text{-GMod}$ the category of graded \mathcal{A} -modules and by $\mathcal{A}\text{-gmod}$ the category of finite dimensional graded \mathcal{A} -modules. Again we call elements $u \in M(x)_\sigma$ *homogeneous elements* of degree σ and write $\text{deg } u = \sigma$. For any graded \mathcal{A} -module M and an element, $\tau \in \mathbf{G}$, we define the *shifted graded module* $M\langle\tau\rangle$, which coincide with M as \mathcal{A} -module, but the grading is given by the rule: $M\langle\tau\rangle_\sigma = M_{\tau\sigma}$. Obviously, the shift $M \mapsto M\langle\tau\rangle$ is an autoequivalence of the category $\mathcal{A}\text{-GMod}$.

We shall usually consider the case, when \mathbf{G} is a group (mainly \mathbb{Z} or \mathbb{Z}^2). Such group gradings are closely related to the *group actions*. We say that a group, \mathbf{G} , *acts on a category*, \mathcal{A} , if a map $T : \mathbf{G} \rightarrow \text{Func}(\mathcal{A}, \mathcal{A})$ is given such that $T1 = \text{Id}$, where 1 is the unit of \mathbf{G} , and $T(\tau\sigma) = T(\tau)T(\sigma)$ for all $\sigma, \tau \in \mathbf{G}$. We do not consider here more general actions with systems of factors, when in the last formula the equality is replaced by an isomorphism of functors. We shall write σx instead of $T(\sigma)x$ both for objects and for morphisms from \mathcal{A} . Given such an action, we can define the *quotient category* \mathcal{A}/\mathbf{G} as follows:

- The objects of \mathcal{A}/\mathbf{G} are the orbits of \mathbf{G} on $\text{ob } \mathcal{A}$.
- $(\mathcal{A}/\mathbf{G})(\mathbf{G}x, \mathbf{G}y)$ is defined as the factorspace of $\bigoplus_{\substack{x' \in \mathbf{G}x \\ y' \in \mathbf{G}y}} \mathcal{A}(x', y')$ modulo the subspace generated by all differences $\alpha - \sigma\alpha$ ($\sigma \in \mathbf{G}$).
- The product of morphisms is defined in the obvious way using representatives (one can easily check that their choice does not affect the result).

The action is called *free* if $\sigma x \neq x$ for every object $x \in \mathcal{A}$ and any $\sigma \neq 1$ from \mathbf{G} . In this case it is easy to see that

$$(\mathcal{A}/\mathbf{G})(\mathbf{G}x, \mathbf{G}y) \simeq \bigoplus_{y' \in \mathbf{G}y} \mathcal{A}(x, y') \simeq \bigoplus_{x' \in \mathbf{G}x} \mathcal{A}(x', y).$$

This allows us to define a \mathbf{G} -grading of \mathcal{A}/\mathbf{G} . Namely, we fix a representative $\widehat{\mathbf{x}}$ in every orbit \mathbf{x} and consider morphisms $\widehat{\mathbf{x}} \rightarrow \sigma\widehat{\mathbf{y}}$ as homogeneous morphisms $\mathbf{x} \rightarrow \mathbf{y}$ of degree σ . One can verify that, whenever the action is free, the quotient category \mathcal{A}/\mathbf{G} is equivalent to the skew group category $\mathcal{A} * \mathbf{G}$ as defined, for instance, in [RR].

Moreover, if the action is free, there is a good correspondence between \mathcal{A} -modules and graded \mathcal{A}/\mathbf{G} -modules. Given an \mathcal{A} -module, M , we define the graded \mathcal{A}/\mathbf{G} -module GM putting $GM(\mathbf{x})_\sigma = M(\sigma\widehat{\mathbf{x}})$ and, for $u \in GM(\mathbf{x})_\sigma$ and $\alpha \in (\mathcal{A}/\mathbf{G})_\tau(\mathbf{x}, \mathbf{y})$, defining their

product as $(\sigma\alpha)u$. It gives a functor, $G : \mathcal{A}\text{-Mod} \rightarrow \mathcal{A}/\mathbf{G}\text{-GMod}$. Conversely, given a graded \mathcal{A}/\mathbf{G} -module N , we define the \mathcal{A} -module $G'N$ putting $G'N(x) = N(\mathbf{G}x)_\sigma$, where $x = \sigma\widehat{\mathbf{G}}x$. One immediately checks that G and G' are mutually inverse equivalences between $\mathcal{A}\text{-Mod}$ and $\mathcal{A}/\mathbf{G}\text{-GMod}$ (cf. [RR]). Moreover, the restrictions of these functors to the categories $\mathcal{A}\text{-mod}$ and $\mathcal{A}/\mathbf{G}\text{-gmod}$ induce an equivalence of these categories as well.

If the category \mathcal{A} has already been \mathbf{H} -graded with a grading semigroup \mathbf{H} and the action of \mathbf{G} preserves this grading, the factorcategory \mathcal{A}/\mathbf{G} becomes $\mathbf{H} \times \mathbf{G}$ graded, and the functors G, G' above induce an equivalence of the categories of \mathbf{H} -graded \mathcal{A} -modules and of $\mathbf{H} \times \mathbf{G}$ -graded \mathcal{A}/\mathbf{G} -modules.

Actually, any group grading can be obtained as the result of a free group action. Namely, given a \mathbf{G} -graded category \mathcal{A} , define a new category $\widetilde{\mathcal{A}}$ with a \mathbf{G} -action as follows:

- The *objects* of $\widetilde{\mathcal{A}}$ are pairs (x, σ) with $x \in \text{ob } \mathcal{A}$, $\sigma \in \mathbf{G}$.
- A morphism, $(x, \sigma) \rightarrow (y, \tau)$, is a pair, (α, σ) , where α is a homogeneous morphism $x \rightarrow y$ of degree $\sigma^{-1}\tau$.
- The product $(\beta, \tau)(\alpha, \sigma)$ is defined as $(\beta\alpha, \sigma)$.
- $\tau(x, \sigma) = (x, \sigma\tau)$, where x is an object or a morphism from \mathcal{A} , $\sigma, \tau \in \mathbf{G}$.

Obviously, this action is free and \mathcal{A} can be identified with $\widetilde{\mathcal{A}}/\mathbf{G}$ as a graded category. Just in the same way, given any graded \mathcal{A} -module M , we turn it into an $\widetilde{\mathcal{A}}$ -module, denoted by \widetilde{M} , setting

- $\widetilde{M}(x, \sigma) = M(x)_\sigma$.
- $(\alpha, \sigma)m = \alpha m$ if $m \in \widetilde{M}(x, \sigma)$, $(\alpha, \sigma) \in \widetilde{\mathcal{A}}((x, \sigma), (y, \tau))$.

This correspondence gives the same equivalence $\widetilde{\mathcal{A}}\text{-Mod} \xrightarrow{\sim} \mathcal{A}\text{-GMod}$ as above.

This allows us to extend all results about module categories to the categories of graded modules. Especially, we can apply the Rickard–Morita Theorem to the category $\mathcal{A}\text{-GMod}$ (note that the category \mathcal{B} from this theorem remains ungraded). We denote by $\mathcal{D}_{gr}\mathcal{A}$ the derived category of $\mathcal{A}\text{-GMod}$. The grading shift $M \mapsto M\langle\sigma\rangle$ naturally extends to the category $\mathcal{D}_{gr}\mathcal{A}$ and commutes with the triangle shift $M \mapsto M[1]$.

There is an important class of gradings, defined as follows.

Definition 2.2. *Let \mathcal{A} be a \mathbf{G} -graded category. We say that it is naturally graded if the category $\widetilde{\mathcal{A}}$ defined above contains a full subcategory $\widetilde{\mathcal{A}}^0 \simeq \mathcal{A}$ such that $\widetilde{\mathcal{A}} = \bigsqcup_{\sigma \in \mathbf{G}} \sigma(\widetilde{\mathcal{A}}^0)$, i.e.*

- $\text{ob } \widetilde{\mathcal{A}} = \bigsqcup_{\sigma \in \mathbf{G}} \sigma(\text{ob } \widetilde{\mathcal{A}}^0)$ (a disjoint union).
- $\widetilde{\mathcal{A}}(x, \sigma y) = 0$ if $x, y \in \widetilde{\mathcal{A}}^0$ and $\sigma \neq 1$.

Actually, it means that one can prescribe a degree, $\deg x \in \mathbf{G}$, to every object $x \in \mathcal{A}$ so that $\mathcal{A}(x, y) = \mathcal{A}(x, y)_{\sigma^{-1}\tau}$ whenever $\deg x = \sigma$, $\deg y = \tau$. In this case also $\mathcal{A}\text{-GMod} \simeq \bigsqcup_{\sigma \in \mathbf{G}} \sigma(\widetilde{\mathcal{A}}^0)\text{-Mod}$ and every component of this coproduct is equivalent to $\mathcal{A}\text{-Mod}$.

Finally, if \mathcal{A} is a \mathbb{Z}^2 -graded category, there are many ways to make \mathcal{A} into a \mathbb{Z} -graded category, taking a kind of “total” grading. Let $\varphi : \mathbb{Z}^2 \rightarrow \mathbb{Z}$ be any epimorphism. We can for example set $\mathcal{A}(x, y)_n = \bigoplus_{\varphi((i,j))=n} \mathcal{A}(x, y)_{(i,j)}$, where $x, y \in \text{ob } \mathcal{A}$, $n \in \mathbb{Z}$ and $(i, j) \in \mathbb{Z}^2$. All such induces \mathbb{Z} -gradings will be called *total*.

2.3 Yoneda categories

For any triangulated category \mathcal{C} and any set $\mathfrak{X} \subseteq \text{ob } \mathcal{C}$ we define the *Yoneda category* $\mathcal{E} = \mathcal{E}(\mathfrak{X})$, which is a \mathbb{Z} -graded category, as follows:

- $\text{ob } \mathcal{E}(\mathfrak{X}) = \mathfrak{X}$.
- $\mathcal{E}(X, Y)_n = \text{Hom}_{\mathcal{C}}(X, Y[n])$.
- The product $\beta\alpha$, where $\alpha : X \rightarrow Y[n]$, $\beta : Y \rightarrow Z[m]$, is defined as $\beta[n]\alpha : X \rightarrow Z[n+m]$.

Note that if $\mathcal{C} = \mathcal{D}\mathcal{A}$ and $\mathfrak{X} \subseteq \mathcal{A}\text{-Mod}$, then $\text{Hom}_{\mathcal{C}}(X, Y[n]) = \text{Ext}_{\mathcal{A}}^n(X, Y)$ and the product $\beta\alpha$ defined above coincides with the Yoneda product $\text{Ext}_{\mathcal{A}}^m(Y, Z) \times \text{Ext}_{\mathcal{A}}^n(X, Y) \rightarrow \text{Ext}_{\mathcal{A}}^{n+m}(X, Z)$.

If \mathcal{A} is a \mathbf{G} -graded category and $\mathcal{C} = \mathcal{D}_{gr}\mathcal{A}$, we also define the *graded Yoneda category* $\mathcal{E}_{gr}(\mathfrak{X})$, which is a $(\mathbb{Z} \times \mathbf{G})$ -graded category, setting $\mathcal{E}_{gr}(X, Y)_{(n, \sigma)} = \text{Hom}_{\mathcal{D}_{gr}\mathcal{A}}(X, Y\langle\sigma\rangle[n])$, which coincide with $\text{Ext}_{\mathcal{A}\text{-GMod}}^n(X, Y\langle\sigma\rangle)$ if X and Y are graded \mathcal{A} -modules. The product of the elements $\alpha : X \rightarrow Y\langle\sigma\rangle[n]$ and $\beta : Y \rightarrow Z\langle\tau\rangle[k]$ is then defined as $\beta\langle\sigma\rangle[n]\alpha : X \rightarrow Z\langle\tau\sigma\rangle[n+k]$. For example, let $\mathfrak{P} = \{\mathcal{A}^x \mid x \in \text{ob } \mathcal{A}\}$, then

$$\mathcal{E}_{gr}(\mathcal{A}^x, \mathcal{A}^y)_{(n, \sigma)} = \begin{cases} 0 & \text{if } n \neq 0, \\ \mathcal{A}(y, x)_{\sigma} & \text{if } n = 0. \end{cases}$$

Thus $\mathcal{E}_{gr}(\mathfrak{P}) \simeq \mathcal{A}^{\text{op}}$ as graded categories.

2.4 Koszul categories

In this subsection we consider \mathbb{Z} -graded categories \mathcal{A} . Moreover, we suppose that \mathcal{A} is basic, bounded and *positively graded*, i.e. $\mathcal{A}(x, y)_n = 0$ if either $n < 0$ or $n = 0$ and $x \neq y$, while $\mathcal{A}(x, x)_0 = \mathbb{k}$. In particular, the objects of \mathcal{A} are pairwise non-isomorphic and their endomorphism algebras contain no nontrivial idempotents. Then the modules $S(x)\langle 0 \rangle = \text{top } \mathcal{A}^x = \mathcal{A}(x, -)_0$ and their shifts $S(x)\langle m \rangle$ are the only simple graded \mathcal{A} -modules. If we consider them as \mathcal{A} -modules *without grading*, we write $S(x)$ for them. Let $\mathfrak{S} = \{S(x)\}$ and $\mathfrak{S}_{gr} = \{S(x)\langle m \rangle\}$. We call the Yoneda category $\mathcal{E}(\mathfrak{S})$ and the graded Yoneda category $\mathcal{E}_{gr}(\mathfrak{S}_{gr})$ respectively the *Yoneda category* and the *graded Yoneda*

category of the positively graded category \mathcal{A} and denote them by $\mathcal{E}(\mathcal{A})$ and by $\mathcal{E}_{gr}(\mathcal{A})$ respectively .

Let \mathcal{A}_+ be the ideal of \mathcal{A} consisting of morphisms of *positive degree*, i.e. $\mathcal{A}_+(x, y) = \sum_{n>0} \mathcal{A}(x, y)_n$, and $V = V_{\mathcal{A}} = \mathcal{A}_+/\mathcal{A}_+^2$. Then V is an \mathcal{A} -bimodule. Set $V^x = V(x, -)$, which is a semisimple gradable \mathcal{A} -module, hence it splits into a direct sum of copies of $S(y)$ for $y \in \text{ob } \mathcal{A}$. We denote by $\nu(x, y)$ the multiplicity of $S(y)$ in $V(x)$ and define the *species* (or the *Gabriel quiver*) of \mathcal{A} as the graph $\Gamma(\mathcal{A})$ such that its set of vertices is $\text{ob } \mathcal{A}$ and there are $\nu(x, y)$ arrows from a vertex x to a vertex y . Equivalently,

$$\nu(x, y) = \dim_{\mathbb{k}} \text{Ext}_{\mathcal{A}}^1(S(x), S(y)) = \sum_{m=1}^{\infty} \dim_{\mathbb{k}} \text{Ext}_{\mathcal{A}\text{-GMod}}^1(S(x)\langle 0 \rangle, S(y)\langle -m \rangle).$$

Note that \mathcal{A}_1 embeds into $V_{\mathcal{A}}$; hence, $\nu(x, y) \geq \dim_{\mathbb{k}} \mathcal{A}(x, y)_1$. If $V(x, y) = \mathcal{A}(x, y)_1$ for all x, y , we say that \mathcal{A} is *generated in degree 1*.

Evidently, the Yoneda category $\mathcal{E}(\mathcal{A})$ is always positively graded. Therefore, the coefficients $\nu(S(x), S(y))$ defining its species are not smaller than $\dim_{\mathbb{k}} \mathcal{E}(S(x), S(y))_1 = \dim_{\mathbb{k}} \text{Ext}_{\mathcal{A}}^1(S(x), S(y))$. Thus the species of \mathcal{A} naturally embed into those of $\mathcal{E}(\mathcal{A})$.

Proposition 2.3. *Suppose that \mathcal{A} is generated in degree 1 and that $\dim_{\mathbb{k}} V_{\mathcal{A}}(x, y) < \infty$ for all $x, y \in \text{ob } \mathcal{A}$. Then the following properties are equivalent:*

- (i) *The Yoneda category $\mathcal{E}(\mathcal{A})$ is generated in degree 1.*
- (ii) *For each object $x \in \text{ob } \mathcal{A}$ there is a projective resolution $\mathcal{P}^\bullet(x)$ of $S(x)\langle 0 \rangle$ such that, for every integer n , $\mathcal{P}^{-n}(x)$ is a direct sum of modules $\mathcal{A}^y\langle -n \rangle$, or, the same, is generated in degree $-n$ (such resolution will be called linear).*
- (iii) *For each object $x \in \text{ob } \mathcal{A}$ there is an injective resolution $\mathcal{I}^\bullet(x)$ of $S(x)\langle 0 \rangle$ such that, for every integer n , $\mathcal{I}^n(x)$ is a direct sum of modules $D\mathcal{A}_y\langle n \rangle$.*
- (iv) *For all x, y, l, m , and n the inequality $\text{Ext}_{\mathcal{A}\text{-GMod}}^n(S(x)\langle l \rangle, S(y)\langle m \rangle) \neq 0$ implies $n = l - m$.*
- (v) $\Gamma(\mathcal{E}(\mathcal{A})) = \Gamma(\mathcal{A})$.
- (vi) $\mathcal{E}(\mathcal{E}(\mathcal{A})) \simeq \mathcal{A}$.

Proof. The equivalence of the properties (i)–(v) is straightforward and well known (cf. [BGS, ADL]), at least if \mathcal{A} contains finitely many objects (i.e. arises from a graded \mathbb{k} -algebra). In the general case the arguments are the same. The equivalence of (v) and (vi) follows immediately from the fact that $\Gamma(\mathcal{A})$ embeds into $\Gamma(\mathcal{E}(\mathcal{A}))$ and the last one embeds into $\Gamma(\mathcal{E}(\mathcal{E}(\mathcal{A})))$. It must also be well known, but we have not found any reference for it. \square

Remark 2.4. We call \mathcal{A} *weakly bounded* if $\dim V(x, y) < \infty$, and both sets $\{z : V(x, z) \neq 0\}$ and $\{z : V(z, y) \neq 0\}$ are finite for all x, y . If the category \mathcal{A} is not bounded, the modules \mathcal{A}^x are usually infinite dimensional, though, if \mathcal{A} is weakly bounded, all spaces $\mathcal{A}^x(y)_n$ are finite dimensional as well. Let M be a graded \mathcal{A} -module such that $\dim M(x)_n < \infty$ for all x . We define the *dual* \mathcal{A}^{op} -module DM by setting $DM(x)_n = D(M(x)_{-n})$ with the natural action of \mathcal{A}^{op} . Obviously, there is a natural isomorphism, $DDM \simeq M$. Especially, the dual modules $\mathcal{I}^x = D\mathcal{A}_x$ are just indecomposable injective modules over \mathcal{A} if \mathcal{A} is weakly bounded. It is easy to see that Proposition 2.3 extends, without any changes, to weakly bounded categories.

The condition Proposition 2.3(vi) is even more powerful than the other conditions in Proposition 2.3. Namely, we have the following (compare with [BGS, Lemma 3.9.2]):

Proposition 2.5. *Let \mathcal{A} be a basic, bounded and positively graded category such that $\mathcal{E}(\mathcal{E}(\mathcal{A})) \simeq \mathcal{A}$ as graded categories. Then \mathcal{A} is generated in degree 1.*

Proof. For an \mathcal{A}_0 -bimodule, B , we set $\dim_{\mathcal{A}} B = (\dim_{\mathbb{k}} 1_x B 1_y)_{x, y \in \text{ob } \mathcal{A}}$. Obviously, we have $\dim_{\mathcal{A}} \mathcal{A}_1 \leq \dim_{\mathcal{A}} \mathcal{A}_+ / \mathcal{A}_+^2$. Further $\dim_{\mathcal{A}} \mathcal{A}_+ / \mathcal{A}_+^2 = \dim_{\mathcal{A}} \mathcal{E}(\mathcal{A})_1$ (note that \mathcal{A}_0 is a subcategory of $\mathcal{E}(\mathcal{A})$ in the natural way and hence the latter notation makes sense). Analogous arguments applied to $\mathcal{E}(\mathcal{A})$ give

$$\dim_{\mathcal{A}} \mathcal{E}(\mathcal{A})_1 \leq \dim_{\mathcal{A}} \mathcal{E}(\mathcal{A})_+ / \mathcal{E}(\mathcal{A})_+^2 = \dim_{\mathcal{A}} \mathcal{E}(\mathcal{E}(\mathcal{A}))_1.$$

Since $\mathcal{E}(\mathcal{E}(\mathcal{A})) \simeq \mathcal{A}$ as graded categories, we obtain $\dim_{\mathcal{A}} \mathcal{E}(\mathcal{E}(\mathcal{A}))_1 = \dim_{\mathcal{A}} \mathcal{A}_1$ and hence all the inequalities above must be in fact equalities. This means that $\dim_{\mathcal{A}} \mathcal{A}_1 = \dim_{\mathcal{A}} \mathcal{A}_+ / \mathcal{A}_+^2$ and thus \mathcal{A} is generated in degree 1. \square

A category, \mathcal{A} , satisfying one of the equivalent conditions of Proposition 2.3 (and hence all of them), will be called *Koszul category*, and the category $\mathcal{E}(\mathcal{A})$ will be called the *Koszul dual* of \mathcal{A} (the word "dual" is justified by the property (vi)). The equivalence of (ii) and (iii) implies that \mathcal{A} is Koszul if and only if so is \mathcal{A}^{op} .

Let \mathcal{A} be a Koszul category of finite global dimension and $S(x, l) = S(x)\langle l \rangle[-l]$, where $x \in \text{ob } \mathcal{A}$, $l \in \mathbb{Z}$. The property (iv) shows that the set $\{S(x, l)\}$ is a tilting subset in $\mathcal{D}_{\text{gr}} \mathcal{A}$. Hence Rickard–Morita Theorem can be applied to the full subcategory \mathcal{S} consisting of these objects. The group \mathbb{Z} acts on \mathcal{S} : $T_n S(x, l) = S(x, l + n)$, and the set $\{S(x, 0)\}$ can be chosen as a set of representatives of the orbits of \mathbb{Z} on $\text{ob } \mathcal{S}$. Moreover,

$$\text{Ext}_{\mathcal{A}}^n(S(x), S(y)) \simeq \bigoplus_{l \in \mathbb{Z}} \text{Ext}_{\mathcal{A}\text{-GMod}}^n(S(x)\langle 0 \rangle, S(y)\langle l \rangle) = \text{Ext}_{\mathcal{A}\text{-GMod}}^n(S(x, 0), S(y, -n)).$$

This implies the following result (mostly also well known).

Theorem 2.6 (Koszul duality). *If \mathcal{A} is a basic and bounded Koszul category of finite global dimension. Then*

1. $\mathcal{D}_{\text{gr}} \mathcal{A} \simeq \mathcal{D} \mathcal{S}^{\text{op}}$.
2. $\mathcal{S} / \mathbb{Z} \simeq \mathcal{E}(\mathcal{A})$ as \mathbb{Z} -graded categories.
3. $\mathcal{D}_{\text{gr}} \mathcal{A} \simeq \mathcal{D}_{\text{gr}} \mathcal{E}(\mathcal{A})^{\text{op}}$.

2.5 Quasi-hereditary categories

Let now \mathcal{A} be a bounded category and let a function, $\text{ht} : \text{ob } \mathcal{A} \rightarrow \mathbb{N} \cup \{0\}$, be given. For every object x define the *standard module* $\Delta(x)$ as the quotient of \mathcal{A}^x modulo the trace of all \mathcal{A}^y with $\text{ht}(y) > \text{ht}(x)$, and the *costandard module* $\nabla(x)$ as $D\Delta^{\text{op}}(x)$, where $\Delta^{\text{op}}(x)$ denotes the standard module for \mathcal{A}^{op} . Set $\mathbf{\Delta} = \{\Delta(x) \mid x \in \text{ob } \mathcal{A}\}$ and $\mathbf{\nabla} = \{\nabla(x) \mid x \in \text{ob } \mathcal{A}\}$. For a set, \mathfrak{X} , of \mathcal{A} -modules, we denote by $\mathcal{F}(\mathfrak{X})$, the full subcategory of \mathcal{A} -mod consisting of the modules which have a filtration with subfactors from \mathfrak{X} (an \mathfrak{X} -filtration). We call the category \mathcal{A} *quasi-hereditary* (with respect to the function ht) if $\text{End}_{\mathcal{A}}(\Delta(x)) = \mathbb{k}$, all composition subquotients of $\text{Rad}(\Delta(x))$ have the form $S(y)$, $\text{ht}(y) < \text{ht}(x)$, and $\mathcal{A}^x \in \mathcal{F}(\mathbf{\Delta})$; or, equivalently, if $\text{End}_{\mathcal{A}}(\nabla(x)) = \mathbb{k}$, all composition subquotients of $\nabla(x)/\text{Soc}(\nabla(x))$ have the form $S(y)$, $\text{ht}(y) < \text{ht}(x)$, and $\mathcal{I}^x \in \mathcal{F}(\mathbf{\nabla})$, where $\mathcal{I}^x = D\mathcal{A}_x$. Obviously, in this case both $\mathbf{\Delta}$ and $\mathbf{\nabla}$ form a set of generators for $\mathcal{D}^{\text{per}} \mathcal{A}$. The notion of a quasi-hereditary category is a natural generalization to this setup of the notion of a quasi-hereditary algebra, [DR1]. One should not confuse it with the notion of a highest weight category from [CPS]. A highest weight category is the category of modules over a quasi-hereditary algebra (or category).

Assume now that \mathcal{A} is a quasi-hereditary category. The arguments of [Rin] can be easily extended to show that for each $x \in \text{ob } \mathcal{A}$ there exists a unique (up to isomorphism) indecomposable module $T(x) \in \mathcal{F}(\mathbf{\Delta}) \cap \mathcal{F}(\mathbf{\nabla})$, called *tilting module*, whose arbitrary standard filtration starts with $\Delta(x)$.

Assume further that \mathcal{A} is positively graded. Following [MO, Section 5] one shows that in this case all simple, projective, standard, injective, costandard, and tilting modules admit graded lifts. For indecomposable modules such lift is unique up to isomorphism and a shift of grading. The grading on \mathcal{A} gives natural graded lifts for projective, standard and simple modules such that we have natural projections $\mathcal{A}^x \twoheadrightarrow \Delta(x)\langle 0 \rangle \twoheadrightarrow S(x)\langle 0 \rangle$ in \mathcal{A} -gmod. Let $x \in \text{ob } \mathcal{A}$. We fix the grading on \mathcal{I}^x and on $\nabla(x)$ such that the natural inclusions $S(x)\langle 0 \rangle \hookrightarrow \nabla(x)\langle 0 \rangle \hookrightarrow \mathcal{I}^x\langle 0 \rangle$ are in \mathcal{A} -gmod. Finally we fix a grading on $T(x)$ such that the natural inclusion $\Delta(x) \hookrightarrow T(x)$ is in \mathcal{A} -gmod and remark that it follows that the natural projection $T(x) \twoheadrightarrow \nabla(x)$ is in \mathcal{A} -gmod.

We have to remark that the lifts above are not coordinated with the isomorphism classes of modules. For example it might happen that some indecomposable \mathcal{A} -module is projective, injective and tilting at the same time. If it is not simple, this module will have different graded lifts when considered as projective module (having the top in degree 0), as injective module (having the socle in degree 0), and as tilting module (having the top in a negative degree and the socle in a positive degree).

The Ringel dual \mathcal{R} is defined as a full subcategory of \mathcal{A} -Mod whose objects are the $T(x)$, $x \in \text{ob } \mathcal{A}$. Since all $T(x)$, $x \in \text{ob } \mathcal{A}$, admit graded lifts, the category \mathcal{R} has a natural structure of a graded category (morphism of degree k from $T(x)$ to $T(y)$ are homogeneous morphisms of degree 0 from $T(x)$ to $T(y)\langle k \rangle$). If \mathcal{A} has finitely many objects, we have the *characteristic tilting module* $T = \bigoplus_{x \in \text{ob } \mathcal{A}} T(x)$ and the category \mathcal{R} corresponds to the (graded) algebra $\text{End}_{\mathcal{A}}(T)$. In the present paper we will always consider \mathcal{R} as a graded category with respect to the above grading.

Now we are ready to formulate the principal assumption for the algebras we consider. They are motivated by the study of the category of linear complexes of tilting modules, associated with a graded quasi-hereditary algebra, see [MO]. From now on we assume that

- (I) for all $x \in \text{ob } \mathcal{A}$ the minimal graded tilting coresolution $\mathcal{T}^\bullet(\Delta(x))$ of $\Delta(x)\langle 0 \rangle$ satisfies $\mathcal{T}^k(\Delta(x)) \in \text{add}(\bigoplus_{y: \text{ht}(y)=\text{ht}(x)-k} T(y)\langle k \rangle)$ for all $k \geq 0$;
- (II) for all $x \in \text{ob } \mathcal{A}$ the minimal graded tilting resolution $\mathcal{T}^\bullet(\nabla(x))$ of $\nabla(x)\langle 0 \rangle$ satisfies $\mathcal{T}^k(\nabla(x)) \in \text{add}(\bigoplus_{y: \text{ht}(y)=\text{ht}(x)+k} T(y)\langle k \rangle)$ for all $k \leq 0$.
- (III) for all $x \in \text{ob } \mathcal{A}$ the minimal graded projective resolution $\mathcal{P}^\bullet(\Delta(x))$ of $\Delta(x)\langle 0 \rangle$ satisfies $\mathcal{P}^k(\Delta(x)) \in \text{add}(\bigoplus_{y: \text{ht}(y)=\text{ht}(x)-k} \mathcal{A}^y\langle k \rangle)$ for all $k \leq 0$;
- (IV) for all $x \in \text{ob } \mathcal{A}$ the minimal graded injective coresolution $\mathcal{I}^\bullet(\nabla(x))$ of $\nabla(x)\langle 0 \rangle$ satisfies $\mathcal{I}^k(\nabla(x)) \in \text{add}(\bigoplus_{y: \text{ht}(y)=\text{ht}(x)+k} \mathcal{I}^y\langle k \rangle)$ for all $k \geq 0$.

Because of [ADL, Theorem 1], the conditions (III) and (IV) are enough to guarantee that the category \mathcal{A} is Koszul, in particular, that it is generated in degree 1.

3 Basic properties of graded quasi-hereditary categories satisfying (I)-(IV)

During this section we always assume that \mathcal{A} is a bounded graded quasi-hereditary category and that (I)-(IV) are satisfied.

Proposition 3.1. *Let $x \in \text{ob } \mathcal{A}$.*

- (i) *All subquotients of any standard filtration of $T(x)\langle 0 \rangle$ have the form $\Delta(y)\langle k \rangle$, where $k \geq 0$ and $\text{ht}(y) = \text{ht}(x) - k$; moreover, $k = 0$ is possible only if $x = y$.*
- (ii) *All subquotients of any costandard filtration of $T(y)\langle 0 \rangle$ have the form $\nabla(y)\langle k \rangle$, where $k \leq 0$ and $\text{ht}(y) = \text{ht}(x) + k$; moreover, $k = 0$ is possible only if $x = y$.*

Proof. We prove (i) using $\mathcal{T}^\bullet(\Delta(x))$ and (I), and the arguments for (ii) are similar (using $\mathcal{T}^\bullet(\nabla(x))$ and (II)). Proceed by induction in $\text{ht}(x)$. If $\text{ht}(x) = 0$, then $T(x)\langle 0 \rangle$ is a standard module and the statement is obvious. Now assume that the statement is proved for all y with $\text{ht}(y) = l - 1$, and let $\text{ht}(x) = l$. Denote by C the cokernel of the graded inclusion $\Delta(x)\langle 0 \rangle \hookrightarrow T(x)\langle 0 \rangle$. By (I), C embeds into a direct sum of several $T(y)\langle 1 \rangle$ with $\text{ht}(y) = l - 1$, such that the cokernel of this embedding has a standard filtration. From the inductive assumption it follows that every subquotient of every standard filtration of such $T(y)\langle 1 \rangle$ has the form $\Delta(z)\langle k + 1 \rangle$, where $k \geq 0$ and $\text{ht}(z) = \text{ht}(y) - k$. Since $\text{ht}(y) = \text{ht}(x) - 1$, the statement follows. \square

Corollary 3.2. *The grading on \mathcal{R} , induced from the category \mathcal{A} -gmod, is positive and \mathcal{R} satisfies (I)-(IV). In particular, the category \mathcal{R}_0 with the same objects as \mathcal{R} and whose morphisms are homogeneous morphisms from \mathcal{R} of degree 0, is semi-simple.*

Proof. Since each $T(x)$ has both a standard and a costandard filtration, from [DR2, Section 1] it follows that every morphism from $T(x)$ to $T(y)$ is a linear combination of morphisms, each of which corresponds to a map from a subquotient of a standard filtration of $T(x)$ to a subquotient of a costandard filtration of $T(y)$. By Proposition 3.1 all subquotients in all standard filtrations of $T(x)\langle 0 \rangle$ live in non-positive degrees and all subquotients in all costandard filtrations of $T(y)\langle 0 \rangle$ live in non-negative degrees. This implies that the grading on \mathcal{R} , induced from $\mathcal{A}\text{-gmod}$, is non-negative. Moreover, from Proposition 3.1 it also follows that the only non-zero graded maps from $T(x)\langle 0 \rangle$ to $T(x)\langle 0 \rangle$ are scalar multiplications, while there are no non-zero graded maps from $T(x)\langle 0 \rangle$ to $T(y)\langle 0 \rangle$ if $x \neq y$. This implies that the zero component of the grading is semi-simple and hence that the grading is in fact positive. That \mathcal{R} satisfies (I)-(I) follows from the fact that (I) and (II) are Ringel dual to (III) and (IV). \square

Corollary 3.3. *Let $x, y \in \text{ob } \mathcal{A}$.*

(i) *The canonical inclusion $\Delta(x) \hookrightarrow T(x)$ induces the following isomorphism:*

$$\text{Hom}_{\mathcal{A}\text{-mod}}(\Delta(y), \Delta(x)) \cong \text{Hom}_{\mathcal{A}\text{-mod}}(\Delta(y), T(x)).$$

(ii) *The canonical projection $T(x) \twoheadrightarrow \nabla(x)$ induces the following isomorphism:*

$$\text{Hom}_{\mathcal{A}\text{-mod}}(\nabla(x), \nabla(y)) \cong \text{Hom}_{\mathcal{A}\text{-mod}}(T(x), \nabla(y)).$$

Proof. Again we will prove (i) and (ii) is proved by similar arguments. The inclusion $\Delta(x) \hookrightarrow T(x)$ induces the inclusion

$$\text{Hom}_{\mathcal{A}\text{-mod}}(\Delta(y), \Delta(x)) \hookrightarrow \text{Hom}_{\mathcal{A}\text{-mod}}(\Delta(y), T(x)),$$

and we have only to verify that the latter inclusion is surjective.

Set $k = \text{ht}(x) - \text{ht}(y)$. Any map from $\Delta(y)$ to $T(x)$ is induced by the unique (up to scalar) map from $\Delta(y)$ to some subquotient of the form $\nabla(y)$ of some costandard filtration of $T(x)$. Hence by Proposition 3.1 the inequality $\text{Hom}_{\mathcal{A}\text{-gmod}}(\Delta(y)\langle i \rangle, T(x)\langle 0 \rangle) \neq 0$ implies $i = -k$ and $k \geq 0$.

Let $f \in \text{Hom}_{\mathcal{A}\text{-mod}}(\Delta(y)\langle k \rangle, T(x)\langle 0 \rangle)$. Consider the tilting coresolution $\mathcal{T}^\bullet(\Delta(x))$, where $\mathcal{T}^0(\Delta(x)) = T(x)\langle 0 \rangle$. Composing f with the differential in this resolution we get a homomorphism, $\bar{f} : \Delta(y)\langle k \rangle \rightarrow \mathcal{T}^1(\Delta(x))$. By (I), we have that $\mathcal{T}^1(\Delta(x))$ is a direct sum of modules of the form $T(z)\langle 1 \rangle$, where $\text{ht}(z) = \text{ht}(x) - 1$. If $\bar{f} \neq 0$, then $\text{Hom}_{\mathcal{A}\text{-gmod}}(\Delta(y)\langle -k \rangle, T(z)\langle 1 \rangle) \neq 0$ for some z . From Proposition 3.1 we hence derive $\text{ht}(y) = \text{ht}(z) - (k + 1)$. Taking $\text{ht}(y) = \text{ht}(x) - k$ into account we get $\text{ht}(x) - k = \text{ht}(x) - 1 - k - 1$, that is $0 = 2$, a contradiction. This implies that $\bar{f} = 0$ that is the image of f is contained in $\Delta(x)\langle 0 \rangle$. The statement follows. \square

Proposition 3.4. *Let $x \in \text{ob } \mathcal{A}$.*

(i) *All subquotients of any standard filtration of $\mathcal{A}^x\langle 0 \rangle$ have the form $\Delta(y)\langle k \rangle$, where $k \leq 0$ and $\text{ht}(y) = \text{ht}(x) - k$; moreover, $k = 0$ is possible only if $x = y$.*

(ii) All subquotients of any costandard filtration of $\mathcal{I}^x\langle 0 \rangle$ have the form $\nabla(y)\langle k \rangle$, where $k \geq 0$ and $\text{ht}(y) = \text{ht}(x) + k$; moreover, $k = 0$ is possible only if $x = y$.

Proof. Analogous to that of Proposition 3.1 using (III) and (IV). \square

Corollary 3.5. For all $x, y \in \text{ob } \mathcal{A}$ the inequality $\text{Ext}_{\mathcal{A}}^1(S(x)\langle 0 \rangle, S(y)\langle k \rangle) \neq 0$ implies $k = -1$ and $|\text{ht}(x) - \text{ht}(y)| = 1$.

Proof. Since \mathcal{A} is quasi-hereditary, $\text{Ext}_{\mathcal{A}}^1(S(x)\langle 0 \rangle, S(y)\langle k \rangle) \neq 0$, in particular, implies $\text{ht}(x) \neq \text{ht}(y)$. Let us first assume that $\text{ht}(x) < \text{ht}(y)$. Then $\text{Ext}_{\mathcal{A}}^1(S(x)\langle 0 \rangle, S(y)\langle k \rangle) \neq 0$ implies that $S(y)\langle k \rangle$ occurs in the top of the kernel K of the canonical projection $\mathcal{A}^x \twoheadrightarrow \Delta(x)\langle 0 \rangle$ since all composition subquotients of $\Delta(x)\langle 0 \rangle$ have the form $S(z)\langle m \rangle$ with $\text{ht}(z) < \text{ht}(x)$. From (III) it follows that the top of K consists of modules of the form $S(z)\langle -1 \rangle$ with $\text{ht}(z) = \text{ht}(x) + 1$. This proves the necessary statement.

In the case $\text{ht}(x) > \text{ht}(y)$ one uses the dual arguments with injective resolutions. \square

Proposition 3.6. Both \mathcal{A} and \mathcal{R} are standard Koszul in the sense of [ADL], in particular, they both are Koszul.

Proof. Follows from (III), (IV), [ADL, Theorem 1], and Corollary 3.2. \square

Proposition 3.7. (i) For every $x \in \text{ob } \mathcal{A}$ the module $\Delta(x)\langle 0 \rangle$ is directed in the following way: for all $l > 0$ we have $[\Delta(x)\langle 0 \rangle_l : S(y)\langle -l \rangle] \neq 0$ implies $\text{ht}(y) = \text{ht}(x) - l$.

(ii) $\dim_{\mathbb{k}} \text{Hom}_{\mathcal{A}\text{-gmod}}(\Delta(y)\langle -l \rangle, \Delta(x)) = [\Delta(x)_l : S(y)\langle -l \rangle]$ for all $y \in \Lambda$.

Proof. To prove the first statement let us first show that $[\Delta(x)\langle 0 \rangle_l : S(y)\langle -l \rangle] \neq 0$ implies $\text{ht}(y) \leq \text{ht}(x) - l$. Indeed, let l be maximal such this statement fails for $\Delta(x)_l$, that is $[\Delta(x)\langle 0 \rangle_l : S(y)\langle -l \rangle] \neq 0$ for some y such that $\text{ht}(y) > \text{ht}(x) - l$. Using Corollary 3.5 we obtain that $\text{Ext}_{\mathcal{A}\text{-gmod}}^1(S(y)\langle -l \rangle, \Delta(x)\langle 0 \rangle_{l+1}) = 0$, that is $S(y)\langle -l \rangle$ is in the socle of $\Delta(x)\langle 0 \rangle$. This implies the existence of a non-zero homomorphism from $\Delta(y)\langle -l \rangle$ to $\Delta(x)$ and hence to $T(x)$ via the canonical inclusion $\Delta(x) \hookrightarrow T(x)$. Thus $T(x)$ must contain $\nabla(y)\langle -l \rangle$ as a subquotient of some costandard filtration. Since $\text{ht}(y) > \text{ht}(x) - l$, this contradicts Proposition 3.1.

Now let us show that $[\Delta(x)\langle 0 \rangle_l : S(y)\langle -l \rangle] \neq 0$ implies $\text{ht}(y) \geq \text{ht}(x) - l$. From the definition of $\Delta(x)$ it follows that $\Delta(x)\langle 0 \rangle$ is obtained by a sequence of universal extensions, which starts from $S(x)\langle 0 \rangle$, and where we are allowed to extend with modules $S(z)\langle m \rangle$ for $\text{ht}(z) \leq \text{ht}(x)$. Applying recursively Corollary 3.5 we see that all simple subquotients, which can be obtained after at most l steps must have the form $S(z)\langle m \rangle$, where $-l \leq m \leq 0$ and $\text{ht}(x) - l \leq \text{ht}(z) \leq \text{ht}(x)$. This gives the necessary inequality.

To prove the second statement we observe that $\dim_{\mathbb{k}} \text{Hom}_{\mathcal{A}\text{-gmod}}(\mathcal{A}^y\langle -l \rangle, \Delta(x)\langle 0 \rangle) = [\Delta(x)\langle 0 \rangle_l : S(y)\langle -l \rangle]$ for all $y \in \Lambda$. Because of (i) the image of any homomorphism $f \in \text{Hom}_{\mathcal{A}\text{-gmod}}(\mathcal{A}^y\langle -l \rangle, \Delta(x)\langle 0 \rangle)$ does not contain simple subquotients $S(z)\langle t \rangle$ with $\text{ht}(z) \geq \text{ht}(y)$. Hence f factors through $\Delta(y)\langle -l \rangle$ and the statement follows. \square

4 Main theorem

Throughout this section we suppose that \mathcal{A} is a bounded graded category, which is quasi-hereditary with respect to some function $\text{ht} : \text{ob } \mathcal{A} \rightarrow \mathbb{N} \cup \{0\}$ and satisfies conditions (I)-(IV). We will use the following notation:

$$\begin{aligned}\kappa(x, l) &= \lfloor (\text{ht}(x) - l)/2 \rfloor; \\ \tilde{\kappa}(x, l) &= \lfloor (-\text{ht}(x) - l)/2 \rfloor; \\ \delta(x, l) &= \begin{cases} 0, & \text{if } \text{ht}(x) \equiv l \pmod{2}, \\ 1 & \text{otherwise;} \end{cases} \\ \Delta(x, l) &= \Delta(x) \langle l \rangle [\kappa(x, l)]; \\ \nabla(x, l) &= \nabla(x) \langle l \rangle [\tilde{\kappa}(x, l)]; \\ T(x, l) &= T(x) \langle l \rangle [\kappa(x, l)]; \\ \mathcal{B} &= \mathcal{B}(\mathcal{A}) = \{ \Delta(x, l) \mid x \in \text{ob } \mathcal{A}, l \in \mathbb{Z} \}; \\ \mathcal{B}' &= \mathcal{B}'(\mathcal{A}) = \{ \nabla(x, l) \mid x \in \text{ob } \mathcal{A}, l \in \mathbb{Z} \}.\end{aligned}$$

We use the same symbols \mathcal{B} and \mathcal{B}' for the full subcategories of $\mathcal{D}_{gr} \mathcal{A}$ with the sets of objects \mathcal{B} and \mathcal{B}' . We also denote by \mathcal{K} the ideal of \mathcal{B} consisting of all morphisms $\Delta(x, l) \rightarrow \Delta(y, m)$ with $\kappa(x, l) \neq \kappa(y, m)$ and $\mathcal{B}^{\text{ver}} = \mathcal{B}/\mathcal{K}$.

Theorem 4.1 (Main Theorem). *In the described situation the following hold:*

- (i) *There is an equivalence, $F : \mathcal{D}_{gr} \mathcal{A} \xrightarrow{\sim} \mathcal{D}(\mathcal{B}^{\text{op}}\text{-Mod})$, of categories such that*
 - (a) $F\Delta(x, l) \simeq \mathcal{B}_{\Delta(x, l)}$;
 - (b) $F\nabla(x, l) \simeq \text{top } \mathcal{B}_{\Delta(x, l)}[-\text{ht}(x)]$;
 - (c) $FT(x, l) \simeq \mathcal{B}_{\Delta(x, l)}^{\text{ver}}$;
- (ii) *Setting $\text{deg} \Delta(x, l) = \kappa(x, l)$ defines a natural \mathbb{Z} -grading on \mathcal{B} , in other words we have $\mathcal{B}(\Delta(x, l), \Delta(y, m)) = \mathcal{B}(\Delta(x, l), \Delta(y, m))_{\kappa(y, m) - \kappa(x, l)}$.*
- (iii) *The group \mathbb{Z} acts on \mathcal{B} in the following way: $T_n \Delta(x, l) = \Delta(x, l + n)$, in particular, \mathcal{B}/\mathbb{Z} becomes a \mathbb{Z}^2 -graded category. Moreover, $\mathcal{B}/\mathbb{Z} \simeq \mathcal{E}_{gr}(\Delta)$ as \mathbb{Z}^2 -graded categories.*
- (iv) *The statements, analogous to (i)-(iii) hold for \mathcal{B}' (and $\mathcal{E}_{gr}(\nabla)$).*
- (v) *There exist total \mathbb{Z} -gradings, associated with the \mathbb{Z}^2 -gradings from (iii) and (iv) respectively, with respect to which the categories \mathcal{B}/\mathbb{Z} and \mathcal{B}'/\mathbb{Z} are Koszul.*
- (vi) *The Koszul dual of the Koszul category \mathcal{B}/\mathbb{Z} is isomorphic to the category \mathcal{B}'/\mathbb{Z} and vice versa.*

The proof of this theorem includes several propositions, which will be stated separately. Most of them consist of some statements about the category \mathcal{B} (especially, the modules $\Delta(x, l)$) and analogous statements about the category \mathcal{B}' (especially, the modules $\nabla(x, l)$). We shall always prove the statements about \mathcal{B} ; those about \mathcal{B}' follow by duality (or can be proved quite in the same way).

Proposition 4.2. 1. *If $\text{Hom}_{\mathcal{D}_{gr}\mathcal{A}}(\Delta(x')\langle l' \rangle[k'], \Delta(x)\langle l \rangle[k]) \neq 0$, then $\text{ht}(x') - 2k' - l' = \text{ht}(x) - 2k - l$.*

2. *If $\text{Hom}_{\mathcal{D}_{gr}\mathcal{A}}(\nabla(x')\langle l' \rangle[k'], \nabla(x)\langle l \rangle[k]) \neq 0$, then $\text{ht}(x') + 2k' + l' = \text{ht}(x) + 2k + l$.*

Proof. Certainly, we may suppose that $k' = l' = 0$. If $\text{Hom}_{\mathcal{D}_{gr}\mathcal{A}}(\Delta(x'), \Delta(x)\langle l \rangle[k]) \neq 0$, also $\text{Hom}_{\mathcal{A}\text{-gmod}}(\Delta(x'), \mathcal{T}^k(\Delta(x))\langle l \rangle) \neq 0$, i.e. $\text{Hom}_{\mathcal{A}\text{-gmod}}(\Delta(x'), \Delta(y)\langle l + k \rangle) \neq 0$ for some y with $\text{ht}(y) = \text{ht}(x) - k$. Proposition 3.7(i) implies that $\text{ht}(x') = \text{ht}(y) - (k + l) = \text{ht}(x) - 2k - l$. \square

Since \mathcal{A} is quasi-hereditary, the sets of objects \mathcal{B} and \mathcal{B}' generate $\mathcal{D}_{gr}\mathcal{A}^{\text{per}}$ as a triangulated category. We denote by $\mathcal{D}_{\text{even}}$ and \mathcal{D}_{odd} the triangulated subcategories of $\mathcal{D}_{gr}\mathcal{A}$ generated by $\{\Delta(x, l) \mid \delta(x, l) = 0\}$ and $\{\Delta(x, l) \mid \delta(x, l) = 1\}$ respectively.

Corollary 4.3. $\text{Hom}_{\mathcal{D}_{gr}\mathcal{A}}(\mathcal{X}, \mathcal{X}') = 0$ if $\mathcal{X} \in \mathcal{D}_{\text{even}}$, $\mathcal{X}' \in \mathcal{D}_{\text{odd}}$ or vice versa. Thus $\mathcal{D}_{gr}\mathcal{A} = \mathcal{D}_{\text{even}} \bigsqcup \mathcal{D}_{\text{odd}}$.

Corollary 4.4. *The categories $\mathcal{D}_{\text{even}}$ and \mathcal{D}_{odd} are generated (as triangular categories) by $\{\nabla(x, l) \mid \delta(x, l) = 0\}$ and $\{\nabla(x, l) \mid \delta(x, l) = 1\}$ respectively.*

Proof. This follows from the fact that $\text{Hom}_{\mathcal{D}_{gr}\mathcal{A}}(\Delta(x, l), \nabla(y, m)[k]) = 0$ if $k \neq \text{ht}(x)$ or $(x, l) \neq (y, m)$. \square

Corollary 4.5. *The sets \mathcal{B} and \mathcal{B}' are tilting subsets of $\mathcal{D}_{gr}\mathcal{A}$.*

Proof. Indeed, it follows from Proposition 4.2 that $\text{Hom}_{\mathcal{D}_{gr}\mathcal{A}}(\Delta(x, l), \Delta(y, m)[n]) = 0$ if $n \neq 0$. \square

Therefore, Rickard–Morita Theorem can be applied to these sets, which implies statements (i) and (ia) of the Main Theorem as well as their analogues for \mathcal{B}' . Moreover, if the functor F satisfies (ia), then

$$\begin{aligned} \text{Hom}_{\mathcal{D}\mathcal{B}^{\text{op}}}(F\Delta(x, l), F\nabla(y, m)[n]) &\simeq \\ &\simeq \text{Hom}_{\mathcal{D}_{gr}\mathcal{A}}(\Delta(x, l), \nabla(y, m)[n]) = \begin{cases} \mathbb{k}, & \text{if } (x, l) = (y, m) \text{ and } n = \text{ht}(x), \\ 0 & \text{otherwise.} \end{cases} \end{aligned} \quad (1)$$

These values coincide with $\text{Hom}_{\mathcal{D}\mathcal{B}^{\text{op}}}(\mathcal{B}_{\Delta(x, l)}, \text{top } \mathcal{B}_{\Delta(y, m)}[n - \text{ht}(x)])$, which gives the statement (ib). It also implies that

$$\begin{aligned} \text{the Yoneda category of } \mathcal{B} \text{ is isomorphic to } \mathcal{B}' \\ \text{and vice versa as ungraded categories.} \end{aligned} \quad (2)$$

Recall that $\text{Hom}_{\mathcal{D}_{gr}\mathcal{A}}(\Delta(x)\langle l\rangle[k], T(y)\langle m\rangle[n]) = 0$ if $k \neq n$. Together with Corollary 3.3 it gives

$$\begin{aligned} \text{Hom}_{\mathcal{D}^{\mathcal{B}^{op}}}(F\Delta(x, l), FT(y, m)[n]) &\simeq \text{Hom}_{\mathcal{D}_{gr}\mathcal{A}}(\Delta(x, l), T(y, m)[n]) = \\ &= \begin{cases} \text{Hom}_{\mathcal{D}_{gr}\mathcal{A}}(\Delta(x, l), \Delta(y, m)), & \text{if } n = 0 \text{ and } \kappa(x, l) = \kappa(y, m), \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

These values coincide with $\text{Hom}_{\mathcal{D}^{\mathcal{B}^{op}}}(\mathcal{B}_{\Delta(x, l)}, \mathcal{B}_{\Delta(y, m)}^{\text{ver}}[n])$, which implies the statement (ic).

Now we define a \mathbb{Z} -grading in \mathcal{B} setting $\text{deg } f = \kappa(y, m) - \kappa(x, l)$ for every morphism $f : \Delta(x, l) \rightarrow \Delta(y, m)$, and consider the corresponding covering $\widetilde{\mathcal{B}}$ (cf. Subsection 2.2). The objects $\Delta(x, l)\langle\kappa(x, l)\rangle \in \widetilde{\mathcal{B}}$ form in $\widetilde{\mathcal{B}}$ a full subcategory $\mathcal{B}^0 \simeq \mathcal{B}$. Moreover, by definition, $\widetilde{\mathcal{B}}(\Delta(x, l)\langle\kappa(x, l)\rangle, \Delta(y, m)\langle n + \kappa(y, m)\rangle)$ consists of the pairs $(f, 0)$, where $f : \Delta(x, l) \rightarrow \Delta(y, m)$ is of degree $n + \kappa(y, m) - \kappa(x, l)$. But every morphism between these modules is of degree $\kappa(y, m) - \kappa(x, l)$. Hence, if $n \neq 0$, there are no nonzero morphisms in $\widetilde{\mathcal{B}}(\Delta(x, l)\langle\kappa(x, l)\rangle, \Delta(y, m)\langle n + \kappa(y, m)\rangle)$. It means that \mathcal{B} is naturally graded thus statement (ii) holds.

Proposition 4.2 also implies that

$$\text{Hom}_{\mathcal{D}_{gr}\mathcal{A}}(\Delta(x, l), \Delta(y, m)) \simeq \text{Hom}_{\mathcal{D}_{gr}\mathcal{A}}(\Delta(x, l + 1), \Delta(y, m + 1)).$$

So the functors $T_n : \Delta(x, l) \mapsto \Delta(x, l + n)$ define a free action of \mathbb{Z} on \mathcal{B} compatible with the grading introduced above. Then the quotient \mathcal{B}/\mathbb{Z} is well defined as a \mathbb{Z}^2 -graded category. The orbits of \mathbb{Z} on the objects of \mathcal{B} are the sets $\{\Delta(x, l) \mid l \in \mathbb{Z}\}$ (with fixed x). So we can identify them with the modules $\Delta(x)$ and choose $\Delta(x, 0)$ as the representative of such an orbit. Then

$$\begin{aligned} (\mathcal{B}/\mathbb{Z})(\Delta(x), \Delta(y))_{(n, m)} &= \text{Hom}_{\mathcal{D}_{gr}\mathcal{A}}(\Delta(x, 0), \Delta(y, m))_n = \\ &= \text{Hom}_{\mathcal{D}_{gr}\mathcal{A}}(\Delta(x, 0), \Delta(y, m))_n = \\ &= \begin{cases} \text{Ext}_{\mathcal{A}\text{-gmod}}^n(\Delta(x)\langle 0\rangle, \Delta(y)\langle m\rangle), & \text{if } n = \kappa(y, m) - \kappa(x, 0), \\ 0 & \text{otherwise,} \end{cases} \end{aligned}$$

which coincides with $\mathcal{E}_{gr}(\Delta(x), \Delta(y))_{(n, m)}$. Thus we have proved statement (iii). Certainly, the statement (iv) follows from (i)-(iii) by duality.

Observe that both \mathcal{B}/\mathbb{Z} and \mathcal{B}'/\mathbb{Z} are naturally \mathbb{Z}^2 -graded and not \mathbb{Z} -graded. Since the formula (1) is compatible with the \mathbb{Z}^2 -grading above, the isomorphisms (2) between the Yoneda category of \mathcal{B} and \mathcal{B}' and vice versa as ungraded categories give rise to isomorphisms between the graded Yoneda category of \mathcal{B}/\mathbb{Z} and \mathcal{B}'/\mathbb{Z} and vice versa as \mathbb{Z}^2 -graded categories. Now we would like to make this \mathbb{Z}^2 -grading into a positive \mathbb{Z} -grading. We will do this for \mathcal{B}/\mathbb{Z} and for \mathcal{B}'/\mathbb{Z} one uses analogous construction: the elements of degree 1 will be non-zero morphisms $\text{Hom}_{\mathcal{D}_{gr}\mathcal{A}}(\Delta(x, l), \Delta(y, m))$, where $\text{ht}(x) = \text{ht}(y) - 1$ and $l = m \pm 1$ (it is easy to see that this grading is given by assigning to $\Delta(x, l)$ the degree $\text{ht } x$). This uniquely determines a total \mathbb{Z} -grading, induced from the original \mathbb{Z}^2 -grading.

Using the positivity of the grading on \mathcal{A} it is straightforward to verify that the grading, defined in this way, is positive. Moreover, it is also easy to see the above isomorphisms between the Yoneda category of \mathcal{B}/\mathbb{Z} and \mathcal{B}'/\mathbb{Z} and vice versa are compatible with this construction (this also follows from the ext-hom duality for standard and constandard modules over quasi-hereditary algebras, see [MO, Theorem 1] and [MO, Theorem 6]). Therefore \mathcal{B}'/\mathbb{Z} is isomorphic to the Yoneda category of \mathcal{B}/\mathbb{Z} and vice versa, now as \mathbb{Z} -graded categories. Applying now Proposition 2.5 and Proposition 2.3, we get both (v) and (vi). This completes the proof of the Main Theorem.

5 Applications of the main result

5.1 Multiplicity free blocks of the BGG category \mathcal{O}

Let \mathfrak{g} be a semi-simple finite-dimensional complex Lie algebra with a fixed triangular decomposition $\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$ and $\lambda \in \mathfrak{h}^*$ be an integral dominant weight. Denote by W_λ the stabilizer of λ with respect to the dot-action of the Weyl group W of \mathfrak{g} on \mathfrak{h}^* . Let A_λ be the basic associative algebra, whose module category is equivalent to the block \mathcal{O}_λ of the BGG-category \mathcal{O} , which corresponds to λ , see [BGG, So1]. Let Δ denote the direct sum of all Verma modules in \mathcal{O}_λ . Let further S denote the set of simple roots associated with W_λ and \mathcal{O}_S denote the corresponding S -parabolic subcategory of \mathcal{O}_0 (see [RC, BGS]). Let $\tilde{\Delta}$ denote the direct sum of all generalized Verma modules in \mathcal{O}_S . Finally, let us denote by B_λ the basic associative algebra, associated with \mathcal{O}_S . In [So1, BGS] it was shown that the algebras A_λ and B_λ are Koszul and even Koszul dual to each other. A quasi-hereditary algebra (or the corresponding highest weight category) is said to be *multiplicity free* if all indecomposable standard modules are multiplicity free.

Theorem 5.1. *Assume that \mathcal{O}_λ is multiplicity free. Then the following holds:*

- (i) \mathcal{O}_S is multiplicity free.
- (ii) The algebra $\text{Ext}_{\mathcal{O}_S}^*(\tilde{\Delta}, \tilde{\Delta})$ is Koszul and even Koszul self-dual.
- (iii) The algebra $\text{Ext}_{\mathcal{O}_\lambda}^*(\Delta, \Delta)$ is Koszul and even Koszul self-dual.

Proof. The primitive idempotents of A_λ are indexed by the highest weights of Verma modules in \mathcal{O}_λ , which are $w \cdot \lambda$, where w is a representative of a cosets W/W_λ . For the antidominant $\mu = w_0 \cdot \lambda$ (here w_0 is the longest element of W) we set $\text{ht}(\mu) = 0$ and for all other $\nu = w \cdot \lambda$ we define $\text{ht}(\nu)$ and the smallest k such that there exist simple reflections s_1, \dots, s_k in W such that $\nu = s_k \dots s_1 \cdot \mu$.

The primitive idempotents of B_λ are indexed by the highest weights of generalized Verma modules in \mathcal{O}_S , which are $w \cdot 0$, where w is the shortest representative of a cosets $W_\lambda \backslash W$. Let w_0^λ be the longest element of W_λ . For the weight $\mu = w_0^\lambda w_0 \cdot \lambda$ we set $\text{ht}(\mu) = 0$ and for all other $\nu = w \cdot \lambda$ as above we define $\text{ht}(\nu)$ and the smallest k such that there exist simple reflections s_1, \dots, s_k in W such that $\nu = s_k \dots s_1 \cdot \mu$.

By [MO, Sections 6,7] and [MO, Appendix], the B_λ -module $\tilde{\Delta}$ admits a linear tilting coresolution, which, under the Koszul duality, becomes the A_λ -module Δ by [ADL]. Moreover, the A_λ -module Δ admits a linear tilting coresolution, which, under the Koszul duality, becomes the B_λ -module $\tilde{\Delta}$ for B_λ .

Assume now that A_λ is multiplicity free. Then the condition (I) for B_λ follows from the known structure of usual Verma modules (see for example [Di, Section 7]). Using the usual duality \star on B_λ (and on A_λ) we also obtain (II). The conditions (III) and (IV) follow from (I) and (II) since B_λ is Ringel self-dual by [So2]. Now Theorem 4.1 implies that $\text{Ext}_{\mathcal{O}_S}^*(\tilde{\Delta}, \tilde{\Delta})$ is Koszul with Koszul dual $\text{Ext}_{\mathcal{O}_S}^*(\tilde{\nabla}, \tilde{\nabla})$, where $\tilde{\nabla}$ is the direct sum of all costandard modules in \mathcal{O}_S . Applying \star induces an isomorphism of these two algebras, which proves (ii).

Further, from the above proof of (ii) and Proposition 3.7 it follows that $\tilde{\Delta}$ is directed in the sense of Proposition 3.7. Now [Di, Section 7] implies that B_λ is multiplicity free, which gives (i).

Finally, let us prove (iii). Again it is enough to prove (I) for A_λ (as A_λ has a duality and is Ringel self-dual by [So2]). If (I) is not satisfied, going to the Koszul dual B_λ we obtain a “wrong” occurrence of a simple in some standard B_λ -module $\tilde{\Delta}(\nu)$. This implies that the original Verma module $\Delta(\nu)$, which surjects onto $\tilde{\Delta}(\nu)$ must have higher multiplicities. Using the Kazhdan-Lusztig Theorem and induction in $\text{ht}(\nu)$, we can further assume that the “wrong” occurrence of a simple in $\tilde{\Delta}(\nu)\langle 0 \rangle$ is in degree 1. This, in turn, would mean that for some standard A_λ -module the condition (I) fails already on the first step. However, in the multiplicity-free case all standard A_λ -modules are directed in the sense of Proposition 3.7 by [Di, Section 7]. Further from the Kazhdan-Lusztig Theorem it follows that on the first step of the construction of the tilting module $T(\nu)$ we extend $\Delta(\nu)$ with $\Delta(\xi)$ for all ξ such that $S(\xi)\langle -1 \rangle$ is a subquotient of $\Delta(\nu)\langle 0 \rangle$. The directness of the standard modules and the already mentioned fact that all standard A_λ -modules have linear tilting coresolutions now imply that the first step of the tilting coresolution of every standard A_λ -module is always correct. A contradiction. This completes the proof of (iii) and of the whole theorem. \square

Remark 5.2. The Koszul grading on both $\text{Ext}_{\mathcal{O}_S}^*(\tilde{\Delta}, \tilde{\Delta})$ and $\text{Ext}_{\mathcal{O}_\lambda}^*(\Delta, \Delta)$ is given by Theorem 4.1 and can be described as follows: Both algebras are generated by elements of degree 0 and 1, and the elements of degree 0 are just scalar automorphisms of generalized Verma and Verma modules respectively. Let l denote the length function on W . Then for $w, w' \in W$ the elements of degree 1 are homomorphisms $\text{Hom}_{\mathcal{O}}(\Delta(w \cdot \lambda), \Delta(w' \cdot \lambda)\langle 1 \rangle)$ and extensions $\text{Ext}_{\mathcal{O}}^1(\Delta(w \cdot \lambda), \Delta(w' \cdot \lambda)\langle -1 \rangle)$ under the additional condition $l(w) = l(w') + 1$. Analogously for generalized Verma modules.

For more information on multiplicity free blocks of \mathcal{O} and \mathcal{O}_S (in particular for classification in the case of maximal stabilizer) we refer the reader to [BC].

Corollary 5.3. *If \mathcal{O}_0 is multiplicity-free (which is the case if and only if $\text{rank}(\mathfrak{g}) \leq 2$) then $\text{Ext}_{\mathcal{O}_0}^*(\Delta, \Delta)$ is Koszul and even Koszul self-dual.*

Proof. By [So1] we have $A_\lambda \cong B_\lambda$ in this case and the statement follows from Theorem 5.1. \square

We would like to emphasize that the algebras $\text{Ext}_{\mathcal{O}_S}^*(\tilde{\Delta}, \tilde{\Delta})$ and $\text{Ext}_{\mathcal{O}_\lambda}^*(\Delta, \Delta)$ in Theorem 5.1 are not Koszul dual to each other in general, though the algebras A_λ and B_λ are.

5.2 Some Koszul quasi-hereditary algebras with Cartan decomposition

Let A be a basic quasi-hereditary algebra over \mathbb{k} with duality and a fixed Cartan decomposition $A = B \otimes_S B^{\text{op}}$, where B is a strong exact Borel subalgebra of A , see [Ko]. Let Λ be the indexing set of simple A - (and hence also of simple B -) modules.

Proposition 5.4. *Assume in the above situation that*

- (1) B is Koszul;
- (2) there is a function, $\text{ht} : \Lambda \rightarrow \{0\} \cup \mathbb{N}$, such that the l -th term of the minimal injective resolution of the simple B -module $L(x)$, $x \in \Lambda$, contains only indecomposable injective modules $I(y)$ such that $\text{ht}(y) = \text{ht}(x) - l$;
- (3) $A \otimes_B -$ sends indecomposable injective B -modules to indecomposable tilting A -modules.

Then A satisfies (I)-(IV). In particular, for the direct sum Δ of all standard A -modules we have that $\text{Ext}_A^*(\Delta, \Delta)$ is Koszul and even Koszul self-dual.

Proof. Since B is an exact Borel subalgebra of A , the functor $A \otimes_B -$ sends simple B -modules to standard A -modules and is exact. This implies that the linear injective coresolution of any simple B -module is sent by $A \otimes_B -$ to a linear tilting coresolution of the corresponding standard A -module. This shows that A satisfies (I) and (II) follows by duality. Since B is an exact Borel subalgebra of A , the functor $A \otimes_B -$ sends indecomposable projective B -modules to indecomposable projective A -modules (see [Ko, Page 408]). Thus the linear projective resolution of any simple B -module is sent by $A \otimes_B -$ to a linear projective resolution of the corresponding standard A -module. This shows that A satisfies (III) and (IV) follows by duality.

Now Theorem 4.1 implies that $\text{Ext}_A^*(\Delta, \Delta)$ is Koszul with Koszul dual $\text{Ext}_A^*(\nabla, \nabla)$, where ∇ is a direct sum of all costandard A -modules. Koszul self-duality of $\text{Ext}_A^*(\nabla, \nabla)$ follows by applying the duality for A . \square

We note that the condition (2) is satisfied for example for incidence algebras, associated with a regular cell decomposition of the sphere \mathbb{S}^n , where $\text{ht}(x)$ denotes the dimension of the cell x , see [KM]. All such algebras are also Koszul, see [KM], so the condition (1) is also satisfied. However, the condition (3) for such algebras fails in the general case.

Acknowledgments

The research was done during the visit of the first author to Uppsala University, which was partially supported by the Faculty of Natural Science, Uppsala University, the Royal Swedish Academy of Sciences, and The Swedish Foundation for International Cooperation in Research and Higher Education (STINT). These supports and the hospitality of Uppsala University are gratefully acknowledged. The second author was also partially supported by the Swedish Research Council. We are in debt to the referee for pointing out several gaps in the original version of the paper and for many useful remarks and suggestions which led to the improvements in the paper.

References

- [ADL] *I. Ágoston, V. Dlab, E. Lukács*, Quasi-hereditary extension algebras. *Algebr. Represent. Theory* 6 (2003), no. 1, 97–117.
- [BGS] *A. Beilinson, V. Ginzburg, W. Soergel*, Koszul duality patterns in representation theory. *J. Amer. Math. Soc.* 9 (1996), no. 2, 473–527.
- [BGG] *I. Bernstein, I. Gelfand, S. Gelfand*, A certain category of \mathfrak{g} -modules. (Russian) *Funkcional. Anal. i Priložen.* 10 (1976), no. 2, 1–8.
- [BC] *B. Boe, D. Collingwood*, Multiplicity free categories of highest weight representations. I, II. *Comm. Algebra* 18 (1990), no. 4, 947–1032, 1033–1070.
- [CPS] *E. Cline, B. Parshall, L. Scott*, Finite-dimensional algebras and highest weight categories. *J. Reine Angew. Math.* 391 (1988), 85–99.
- [Di] *J. Dixmier*, Enveloping algebras. Revised reprint of the 1977 translation. *Graduate Studies in Mathematics*, 11. American Mathematical Society, Providence, RI, 1996.
- [DR1] *V. Dlab, C. M. Ringel*, Quasi-hereditary algebras. *Illinois J. Math.* 33 (1989), no. 2, 280–291.
- [DR2] *V. Dlab, C. M. Ringel*, The module theoretical approach to quasi-hereditary algebras. *Representations of algebras and related topics (Kyoto, 1990)*, 200–224, *London Math. Soc. Lecture Note Ser.*, 168, Cambridge Univ. Press, Cambridge, 1992.
- [Ke] *B. Keller*, Deriving DG categories, *Ann. Sci. École Norm. Sup.* 27 (1994), no.1, 63–102.
- [Ko] *S. König*, Exact Borel subalgebras of quasi-hereditary algebras. I. With an appendix by Leonard Scott. *Math. Z.* 220 (1995), no. 3, 399–426.
- [KM] *H. Kovilyanskaya, V. Mazorchuk*, On incidence algebras associated with regular cell decomposition of \mathbb{S}^n . *Publ. Math. Debrecen* 54 (1999), no. 3-4, 391–402.

- [MO] *V. Mazorchuk and S. Ovsienko*, A pairing in homology and the category of linear tilting complexes for a quasi-hereditary algebra, With an appendix written by C. Stroppel, *J. Math. Kyoto Univ.* 45 (2005), 711–743.
- [RR] *I. Reiten and Ch. Riedtmann*, Skew group algebras in the representation theory of Artin algebras. *J. Algebra* 92 (1985), 224–282.
- [Ric] *J. Rickard*, Morita theory for derived categories. *J. London Math. Soc.* 39 (1989), 436–456.
- [Rin] *C. M. Ringel*, The category of modules with good filtrations over a quasi-hereditary algebra has almost split sequences. *Math. Z.* 208 (1991), no. 2, 209–223.
- [RC] *A. Rocha-Caridi*, Splitting criteria for \mathfrak{g} -modules induced from a parabolic and the Bernstein-Gelfand-Gelfand resolution of a finite-dimensional, irreducible \mathfrak{g} -module. *Trans. Amer. Math. Soc.* 262 (1980), no. 2, 335–366.
- [So1] *W. Soergel*, Kategorie \mathcal{O} , perverse Garben und Moduln über den Koinvarianten zur Weylgruppe. (German) *J. Amer. Math. Soc.* 3 (1990), no. 2, 421–445.
- [So2] *W. Soergel*, Character formulas for tilting modules over Kac-Moody algebras. *Represent. Theory* 2 (1998), 432–448.

Yuriy Drozd, Department of Mechanics and Mathematics, Kyiv Taras Shevchenko University, 64, Volodymyrska st., 01033, Kyiv, Ukraine, e-mail: yuriy@drozd.org,
url: <http://bearlair.drozd.org/~yuriy>.

Volodymyr Mazorchuk, Department of Mathematics, Uppsala University, Box 480, 751 06, Uppsala, SWEDEN, e-mail: mazor@math.uu.se, url: <http://www.math.uu.se/~mazor>.