

On the P - and T -non-invariant two-component equation for the neutrino

W.I. FUSHCHYCH

The relativistic two-component equation describing the free motion of particles with zero mass and spin $\frac{1}{2}$, which is P - and T -non-invariant but C -invariant, is found. The representation of the Poincaré group for zero mass and discrete spin is constructed. The position operator for such a particle is defined.

1. Introduction

As is known, the Dirac equation for a particle with zero mass:

$$i \frac{\partial \Psi(t, \mathbf{x})}{\partial t} = \gamma_0 \gamma_k p_k \Psi(t, \mathbf{x}), \quad k = 1, 2, 3, \quad (1.1)$$

is invariant with respect to the space-time reflections. If one chooses for the Dirac matrices the Weyl representation eq. (1.1) decomposes into a system of two equations

$$i \frac{\partial \Psi_{\pm}(t, \mathbf{x})}{\partial t} = \pm \sigma_k p_k \Psi_{\pm}(t, \mathbf{x}), \quad (1.2)$$

where σ_k are the Pauli matrices and Ψ_{\pm} is a two-component spinor. The Weyl equation (1.2) for Ψ_+ (or Ψ_-) is not invariant under space reflection P and charge conjugation C but is invariant under the CP - and T -operations.

Due to the fact that the space parity in the weak interactions is not conserved it is usually assumed that neutrino is described, not by the four-component eq. (1.1), but by a two-component one (1.2). Therefore, in papers [1] an hypothesis was put forward that the weak interactions are invariant with respect to the CP operation and consequently to the T operation, if the CPT theorem is valid.

In this paper the two-component equation for a particle with zero mass and spin $\frac{1}{2}$, which is non-invariant under the time reflection of T and the CP operation, is found.

2. Equation for a neutrino with “variable mass”

On the solutions of eq. (1.1) the generators of the Poincaré group $P(1, 3)$ have the form

$$P_0^{\Psi} = \mathcal{H}^{\Psi} = \gamma_0 \gamma_k p_k, \quad P_k^{\Psi} = p_k, \quad (2.1)$$

$$J_{kl}^{\Psi} = x_k p_l - x_l p_k + S_{kl}, \quad J_{0k}^{\Psi} = x_0 p_k - \frac{1}{2} [x_k, \mathcal{H}^{\Psi}]_+,$$

$$S_{\mu\nu} = \frac{1}{4} i (\gamma_{\mu} \gamma_{\nu} - \gamma_{\nu} \gamma_{\mu}), \quad S_{\mu 4} = \frac{1}{4} i (\gamma_{\mu} \gamma_4 - \gamma_4 \gamma_{\mu}), \quad (2.1')$$

$$S_{\mu 5} = \frac{1}{2} i \gamma_{\mu}, \quad S_{45} = \frac{1}{2} i \gamma_4, \quad \mu = 0, 1, 2, 3,$$

where γ_{μ} and γ_4 are the Dirac matrices.

If one performs a unitary transformation [2] over eq. (1.1)

$$U_1 = \exp \left\{ \frac{1}{2} i \pi S_{53} e_3 \right\}, \quad e_3 = \frac{p_3}{|p_3|}, \quad p_3 \neq 0, \quad (2.2)$$

or

$$U_1 = \frac{1}{\sqrt{2}} (1 + \gamma_3 e_3), \quad (2.2')$$

eq. (1.1) has the form

$$i \frac{\partial \chi(t, \mathbf{x})}{\partial t} = (\gamma_0 \gamma_a p_a + \gamma_0 |p_3|) \chi(t, \mathbf{x}), \quad a = 1, 2, \quad (2.3)$$

$$\chi = U_1 \Psi, \quad \chi \equiv \begin{pmatrix} \chi_+ \\ \chi_- \end{pmatrix}, \quad (2.4)$$

where χ_{\pm} is a two-component spinor.

The Poincaré group generators $P(1, 3)$ on $\{\chi\}$ being the solution of eq. (2.3) have the form

$$\begin{aligned} P_0^X &= \mathcal{H}^X = \gamma_0 \gamma_a p_a + \gamma_0 |p_3|, & P_k^X &= p_k, \\ J_{ab}^X &= x_a p_b - x_b p_a + S_{ab}, & J_{a3}^X &= x_a p_3 - x_3 p_a - e_3 S_{a3} \gamma_3, \\ J_{0k}^X &= x_0 p_k - \frac{1}{2} [x_k, \mathcal{H}^X]_+. \end{aligned} \quad (2.5)$$

Choosing for the Dirac matrices somewhat unusual representation

$$\begin{aligned} \gamma_0 &= \begin{pmatrix} \sigma_3 & 0 \\ 0 & -\sigma_3 \end{pmatrix}, & \gamma_a &= \begin{pmatrix} i\sigma_a & 0 \\ 0 & -i\sigma_a \end{pmatrix}, \\ \gamma_3 &= \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}, & \gamma_4 &= \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}, \end{aligned} \quad (2.6)$$

eq. (2.3) decomposes into a system of two equations

$$\begin{aligned} i \frac{\partial \chi_{\pm}(t, \mathbf{x})}{\partial t} &= \{i\sigma_3 \sigma_a p_a \pm \sigma_3 |p_3|\} \chi_{\pm}(t, \mathbf{x}), \\ \chi_{\pm} &= Q_{\pm} \chi, \quad Q_{\pm} = \frac{1}{2} \pm i S_{43} = \frac{1}{2} (1 \pm \gamma_3 \gamma_4). \end{aligned} \quad (2.7)$$

Eq. (2.7) for the functions $\chi_+(t, \mathbf{x})$ (or $\chi_-(t, \mathbf{x})$) has quite the other properties relative to the discrete transformations than the Weyl equation (1.2).

We note the following:

(i) It is possible to arrive at eq. (2.3) (or (2.7)) in another way. If we “extract the square root” from the operator equation

$$(p_0^2 - p_a^2) \chi = p_3^2 \chi,$$

we obtain eqs. (2.3) (or eqs. (2.7)).

(ii) The fact that the Dirac equations for zero and non-zero mass are invariant under the P -, T - and C -transformations is the consequence of the fact that they, besides being invariant with respect to the group $P(1, 3)$, are invariant under the

group $SU(2) \otimes SU(2) \sim O(4)$ (this question will be considered in detail in a following paper).

Eq. (2.3) coincides in form with a usual Dirac equation for zero mass if $|p_3|$ is considered as the mass of a particle. Therefore it is possible to say that eq. (2.3) describes a “flat neutrino” with variable mass $|p_3|$. Really the operator $|p_3|$ is the Casimir operator of the group $P(1,2)$ but not of the group $P(1,3)$.

Before passing to an investigation of the P -, T - and C -properties of eqs. (2.7) we shall construct the operator of the position in the space.

For eq. (2.3) the operator of the Foldy–Wouthuysen type has the form

$$U_2 = \exp \left\{ \frac{S_{5a} p_a}{\sqrt{p_a^2}} \operatorname{arctg} \frac{\sqrt{p_a^2}}{|p_3|} \right\}. \quad (2.8)$$

If the matrix S_{5a} have the form of (2.1') then

$$U_2 = \frac{E + |p_3| + \gamma_a p_a}{\{2E(E + |p_3|)\}^{1/2}}, \quad E = \sqrt{p_1^2 + p_2^2 + p_3^2}. \quad (2.9)$$

Eq. (2.3) after the transformation (2.9) transfers into

$$i \frac{\partial \Phi(t, \mathbf{x})}{\partial t} = \mathcal{H}^\Phi(t, \mathbf{x}) = \gamma_0 E \Phi(t, \mathbf{x}), \quad \Phi(t, \mathbf{x}) = U_2 \chi(t, \mathbf{x}). \quad (2.10)$$

The generators of the group $P(1,3)$ on $\{\Phi\}$ have the form

$$\begin{aligned} P_0^\Phi &= \mathcal{H}^\Phi = \gamma_0 E, & P_k^\Phi &= p_k, \\ J_{ab}^\Phi &= x_a p_b - x_b p_a + S_{ab}, & J_{a3}^\Phi &= x_a p_3 - x_3 p_a - e_3 \frac{S_{ab} p_b}{E + |p_3|}, \\ J_{0a}^\Phi &= x_0 p_a - \frac{1}{2} [x_a, \mathcal{H}^\Phi]_+ - \gamma_0 \frac{S_{ab} p_b}{E + |p_3|}, & J_{03}^\Phi &= x_0 p_3 - \frac{1}{2} [x_3, \mathcal{H}^\Phi]_+. \end{aligned} \quad (2.11)$$

It must be noted that the operators (2.11), as it can be immediately verified, satisfy the algebra $P(1,3)$ commutation relations not depending on the matrices S_{ab} explicit form, i.e. the operators (2.11), if γ_0 is substituted for 1 (or -1) and realize irreducibly the algebra $P(1,3)$ representation which is characterized by zero mass and discrete spin. The representation (2.11) differs from the corresponding Shirokov [3], Lomont–Moses [4] ones but is certainly equivalent to them.

The position operator on a set $\{\chi\}$ looks as

$$\begin{aligned} X_a^\chi &= U_2^{-1} x_a U_2 = x_a - \frac{S_{5a}}{E} + \frac{S_{5c} p_c p_a}{E^2 (E + |p_3|)} + \frac{S_{ac} p_c}{E (E + |p_3|)}, \\ X_3^\chi &= U_2^{-1} x_3 U_2 = x_3 + e_3 \frac{S_{5c} p_c}{E^2}, & S_{5c} &= -\frac{1}{2} i \gamma_c. \end{aligned} \quad (2.12)$$

The position operator on a set of solution $\{\Psi\}$ of eq. (1.1) looks as follows

$$\begin{aligned} X_a^\Psi &= U_1^{-1} X_a^\chi U_1 = x_a + e_3 \frac{\gamma_3 S_{5a}}{E} - e_3 \frac{\gamma_3 S_{5c} p_c p_a}{E^2 (E + |p_3|)} + \frac{S_{ac} p_c}{E (E + |p_3|)}, \\ X_3^\Psi &= U_1^{-1} X_3^\chi U_1 = x_3 - \frac{\gamma_3 S_{5c} p_c}{E^2}. \end{aligned} \quad (2.13)$$

(iii) If one performs a transformation on eq. (1.1)

$$\tilde{U}_1 = \frac{1}{\sqrt{2}}(1 + \gamma_3) \quad (2.14)$$

and then a transformation

$$\tilde{U}_2 = \frac{E + p_3 + \gamma_a p_a}{\{2E(E + |p_3|)\}^{1/2}}, \quad (2.15)$$

it will transform into the equation

$$i \frac{\partial \tilde{\Phi}(t, \mathbf{x})}{\partial t} = \gamma_0 \tilde{\Phi}(t, \mathbf{x}), \quad \tilde{\Phi} = \tilde{U}_2 \tilde{U}_1 \Psi. \quad (2.16)$$

The generators of the group $P(1, 3)$ on $\{\tilde{\Phi}\}$ coincide with (2.11) where the substitution was made $e_3 \rightarrow 1$, $|p_3| \rightarrow p_3$.

3. P-, T- and C-properties of two-component equation

Here we shall study the properties of one of the two-component eqs. (2.7)¹

$$i \frac{\partial \chi(t, \mathbf{x})}{\partial t} = (i\sigma_3 \sigma_a p_a + \sigma_3 |p_3|) \chi(t, \mathbf{x}), \quad (3.1)$$

under the discrete transformations.

We shall denote through $P^{(k)}$ ($k = 1, 2, 3$) the space inversion operator of one axis which is determined as

$$P^{(1)} \chi(t, x_1, x_2, x_3) = r^{(1)} \chi(t, -x_1, x_2, x_3). \quad (3.2)$$

Analogously $P^{(2)}$ and $P^{(3)}$ are determined.

As is well known, two non-equivalent definitions of the time-reflection operator exist. According to Wigner the time-inversion operator is

$$T^{(1)} \chi(t, \mathbf{x}) = \tau^{(1)} \chi^*(-t, \mathbf{x}). \quad (3.3)$$

According to Pauli it is:

$$T^{(2)} \chi(t, \mathbf{x}) = \tau^{(2)} \chi(-t, \mathbf{x}). \quad (3.4)$$

The operator of the charge conjugation can be defined as the product of the operators $T^{(1)}$, $T^{(2)}$ or as

$$C \chi(t, \mathbf{x}) = \tau^{(3)} \chi^*(t, \mathbf{x}), \quad (3.5)$$

where $r^{(k)}$, $\tau^{(k)}$ are the 2×2 matrices.

The operators P , T , C with the group $P(1, 3)$ generators satisfy the usual commutation relations.

The generators of the group $P(1, 3)$ on the solutions $\{\chi\}$ of eq. (3.1) have the form of eq. (2.5) where

$$\begin{aligned} \mathcal{H}^\chi &\rightarrow i\sigma_3 \sigma_a p_a + \sigma_3 |p_3| = -\sigma_2 p_1 + \sigma_2 p_2 + \sigma_3 |p_3|, \\ S_{ab} &\rightarrow \frac{1}{4} i(\sigma_b \sigma_a - \sigma_a \sigma_b), \quad S_{a3} \gamma_3 \rightarrow -\frac{1}{2} \sigma_a, \end{aligned} \quad (3.6)$$

and the matrix γ_0 is substituted for the matrix σ_3 .

¹In what follows, under χ we shall understand the two-component spinor χ_+ .

Using the definitions (3.2)–(3.5) it is not difficult to verify that eq. (3.1) is $P^{(3)}$ -, C -invariant but $P^{(1)}$ -, $P^{(2)}$ -, $T^{(1)}$ -, $T^{(2)}$ -non-invariant.

Thus, eq. (3.1) is $P^{(3)}C$ -, $P^{(1)}P^{(2)}P^{(3)}C$ - and $P^{(a)}CT^{(a)}$ -invariant but $P^{(3)}CT^{(a)}$ - and $P^{(a)}C$ -non-invariant.

We note the following:

(i) The result obtained is a consequence of the fact that the projection operators Q_{\pm} , with the operators of the discrete transformations, satisfy the following relations

$$\begin{aligned} P^{(a)}Q_{\pm} &= Q_{\mp}P^{(a)}, & T^{(a)}Q_{\pm} &= Q_{\mp}T^{(a)}, \\ P^{(3)}Q_{\pm} &= Q_{\pm}P^{(3)}, & CQ_{\pm} &= Q_{\pm}C. \end{aligned} \quad (3.7)$$

(ii) The two-component equations for the functions χ_+ and χ_- are equivalent to the four-component one (2.3) with the subsidiary relativistic-invariant conditions

$$Q_-\chi = \left(\frac{1}{2} - iS_{43}\right)\chi = \frac{1}{2}(1 - \gamma_3\gamma_4)\chi = 0, \quad (3.8)$$

$$Q_+\chi = \left(\frac{1}{2} + iS_{43}\right)\chi = \frac{1}{2}(1 + \gamma_3\gamma_4)\chi = 0, \quad (3.9)$$

respectively. For eq. (1.1) these conditions look like

$$\left(\frac{1}{2} + ie_3S_{45}\right)\Psi = \frac{1}{2}(1 - e_3\gamma_4)\Psi = 0, \quad (3.8')$$

$$\left(\frac{1}{2} - ie_3S_{45}\right)\Psi = \frac{1}{2}(1 + e_3\gamma_4)\Psi = 0. \quad (3.9')$$

Eq. (1.1) with the subsidiary conditions (3.8') and (3.9') can be joined and can be written in the form of two $P^{(a)}$ - and $T^{(b)}$ -non-invariant but $P^{(3)}$ - and C -invariant equations

$$\{\gamma_{\mu}p^{\mu} + \varkappa(1 + e_3\gamma_4)\}\Psi_1(t, \mathbf{x}) = 0, \quad \{\gamma_{\mu}p^{\mu} + \varkappa(1 - e_3\gamma_4)\}\Psi_2(t, \mathbf{x}) = 0,$$

where \varkappa is some constant value. The four-component equations for the neutrino, which are the union of eq. (1.1) and the usual subsidiary condition, were recently considered in ref. [6]. These equations, as well as the Weyl equations (1.2), are P - and C -non-invariant but $T^{(1)}$ -invariant.

The unitary operator of type U_2 for the two-component eq. (3.1) has the form

$$V_1 = \exp\left\{i\frac{S_a p_a}{\sqrt{p_a^2}} \arctg \frac{\sqrt{p_a^2}}{|p_3|}\right\}, \quad S_k = \frac{1}{2}\varepsilon_{klm}S_{lm}, \quad (3.10)$$

or

$$V_1 = \frac{E + |p_3| + i\sigma_a p_a}{\{2E(E + |p_3|)\}^{1/2}}. \quad (3.11)$$

The position operator on the set of solutions $\{\chi\}$ of eqs. (3.1) looks as follows

$$\begin{aligned} X_a^{\chi+} &= V_1^{-1}x_a V_1 = x_a - \frac{\sigma_a}{2E} + \frac{\sigma_c p_c p_a}{2E^2(E + |p_3|)} - i\frac{(\sigma_a \sigma_c - \sigma_c \sigma_a)p_c}{4E(E + |p_3|)}, \\ X_3^{\chi+} &= V_1^{-1}x_3 V_1 = x_3 + e_3 \frac{\sigma_b p_b}{2E^2}. \end{aligned} \quad (3.12)$$

To complete our treatment, we find the position operator for the neutrino which is described by the Weyl equation (1.2), for example for the function Ψ_+ . This equation under a transformation

$$V = \frac{E + |p_3| + i\sigma_k \xi_k}{2\sqrt{\xi_k p_k}}, \quad (3.13)$$

where the vector ξ has the following components

$$\xi_k \equiv \{p_1 - p_2 e_3, p_2 + e_3 p_1, e_3(E + |p_3|)\},$$

takes a canonical form

$$i \frac{\partial \Phi_+(t, \mathbf{x})}{\partial t} = \sigma'_3 E \Phi_+(t, \mathbf{x}), \quad \sigma'_3 = \sigma_3 e_3, \quad \Phi_+(t, \mathbf{x}) = V \Psi_+(t, \mathbf{x}). \quad (3.14)$$

The position operator for a neutrino which is described by the Weyl equation (1.2) (for Ψ_+) looks like

$$X_a^W = V^{-1} x_a V = x_a + i e_3 \frac{\sigma_3 \sigma_a}{2E} - i \frac{e_3 \sigma_3 \sigma_c p_c p_a}{2E^2(E + |p_3|)} - i \frac{(\sigma_a \sigma_c - \sigma_c \sigma_a) p_c}{4E(E + |p_3|)},$$

$$X_3^W = V^{-1} x_3 V = x_3 - i \frac{\sigma_3 \sigma_b p_b}{2E^2}.$$

The other definitions of the operators X_k and V for the neutrino are given in ref. [5].

(iii) From Dirac eq. (1.1) one can, generally speaking, obtain three types of non-equivalent two-component equations. On the set of solutions of eq. (1.1) a direct sum of four irreducible representations $D^\varepsilon(s)$ of the group $P(1,3)$

$$D^{\varepsilon=1} \left(s = \frac{1}{2} \right) \oplus D^{\varepsilon=-1} \left(s = -\frac{1}{2} \right) \oplus D^{\varepsilon=1} \left(s = -\frac{1}{2} \right) \oplus D^{\varepsilon=-1} \left(s = \frac{1}{2} \right) \quad (3.15)$$

is realized, where ε is an energy sign, s is a helicity. Hence it follows that there exist three types of two-component equations on the set of which the following representation of the group $P(1,3)$

$$D^{\varepsilon=1} \left(s = \frac{1}{2} \right) \oplus D^{\varepsilon=-1} \left(s = -\frac{1}{2} \right),$$

or

$$D^{\varepsilon=1} \left(s = -\frac{1}{2} \right) \oplus D^{\varepsilon=-1} \left(s = \frac{1}{2} \right), \quad D^{\varepsilon=1} \left(s = \frac{1}{2} \right) \oplus D^{\varepsilon=-1} \left(s = \frac{1}{2} \right), \quad (3.16)$$

or

$$D^{\varepsilon=1} \left(s = -\frac{1}{2} \right) \oplus D^{\varepsilon=-1} \left(s = -\frac{1}{2} \right), \quad D^{\varepsilon=1} \left(s = \frac{1}{2} \right) \oplus D^{\varepsilon=-1} \left(s = -\frac{1}{2} \right), \quad (3.17)$$

or

$$D^{\varepsilon=1} \left(s = \frac{1}{2} \right) \oplus D^{\varepsilon=-1} \left(s = -\frac{1}{2} \right) \quad (3.18)$$

are realized. If on the solutions of two-component equation there realizes the representation (3.16) then this equation will be $T^{(1)}$ -invariant but C -, P -, $T^{(2)}$ -non-invariant,

if the representation (3.17) does then it will be $T^{(1)}$ -, $T^{(2)}$ -, C -invariant but P -non-invariant, and if the representation (3.18) it will be $T^{(1)}$ -, P -invariant but C -, $T^{(2)}$ -non-invariant. This problem will be considered in more detail in another paper.

4. Equation for a flat neutrino

The motion group in the Minkovski three-space is the $P(1,2)$ group of rotations and translations conserving the form

$$x^2 = x_0^2 - x_1^2 - x_2^2.$$

In this case the simplest spinor equation is

$$i \frac{\partial \chi_{\pm}(t, x_1, x_2)}{\partial t} = (i\sigma_3 \sigma_a p_a \pm \sigma_3 m) \chi_{\pm}(t, x_1, x_2), \quad (4.1)$$

χ_{\pm} is the two-component spinor and m is the eigenvalue of the operator $\sqrt{P_{\mu}^2}$.

Eq. (4.1) for χ_+ (or χ_-) like eq. (3.1) is invariant under the $P^{(1)}P^{(2)}$ - and C -operations but non-invariant under the $P^{(a)}$ and $T^{(b)}$ -operations.

Thus, eq. (4.1) for the wave function χ_+ (or χ_-) is $P^{(1)}P^{(2)}C$ -, $T^{(a)}P^{(b)}$ - and $P^{(a)}CT^{(b)}$ -invariant but $P^{(a)}C$ - and $CT^{(a)}$ -non-invariant.

It should be noted that the equation being the "direct sum" of the equation for $\chi_+(t, x_1, x_2)$ and $\chi_-(t, x_1, x_2)$ is invariant under the P -, T - and C -transformations [7].

Finally, we quote one more example of the P - and C -non-invariant equation which is invariant with respect to the inhomogeneous De Sitter group. Such is the Dirac equation:

$$i \frac{\partial \Psi(t, \mathbf{x}, x_4)}{\partial t} = (\gamma_0 \gamma_k p_k + \gamma_0 \varkappa) \Psi(t, \mathbf{x}, x_4), \quad k = 1, 2, 3, 4. \quad (4.2)$$

This equation as is shown in refs. [2, 7] is $T^{(1)}$ -, $T^{(2)}C$ -invariant but $P^{(k)}$ -, $T^{(2)}$ - and C -non-invariant.

All the results obtained in this paper can be generalized for the arbitrary spin s case, if one uses for this the purpose the equation (ref. [2]):

$$i \frac{\partial \Psi(t, \mathbf{x})}{\partial t} = \lambda S_{0l} p_l \Psi(t, \mathbf{x}), \quad l = 1, 2, 3, \quad (4.3)$$

where λ is some fixed parameter (for the Dirac equation $\lambda = -2i$), and $S_{\mu\nu}$, $S_{\mu 4}$, S_{45} are the matrices (not 4×4 ones) realizing the algebra $O(1,5)$ representation.

(i) If we transform the usual Dirac equation describing the motion of the non-zero mass particle m with a spin $\frac{1}{2}$ as

$$V_2 = \frac{\gamma_3 p_3 + q_3 + m}{\{2q_3(q_3 + m)\}^{1/2}}, \quad q_3 \equiv \sqrt{p_3^2 + m^2}, \quad (4.4)$$

it has the form

$$i \frac{\partial \Psi'(t, \mathbf{x})}{\partial t} = H' \Psi'(t, \mathbf{x}), \quad (4.5)$$

$$H' = \gamma_0 \gamma_a p_a + \gamma_0 q_3, \quad \Psi' = V_2 \Psi, \quad a = 1, 2. \quad (4.6)$$

Choosing the representation (2.6) for the Dirac matrices eq. (4.5) is decomposed into the set of two independent equations

$$i \frac{\partial \Psi'_+(t, \mathbf{x})}{\partial t} = (-\sigma_2 p_1 + \sigma_1 p_2 + \sigma_3 q_3) \Psi'_+(t, \mathbf{x}), \quad (4.7)$$

$$i \frac{\partial \Psi'_-(t, \mathbf{x})}{\partial t} = (-\sigma_2 p_1 + \sigma_1 p_2 - \sigma_3 q_3) \Psi'_-(t, \mathbf{x}), \quad (4.8)$$

where Ψ'_+ and Ψ'_- are two-component wave functions.

Eq. (4.7) or (4.8) describes a free motion of spinless particle and antiparticle with the mass m . Thus besides of the Klein–Gordon equation there exist the other equations of the type (4.7) and (4.8) which are also relativistically invariant and describe the spinless particle motion with non-zero mass. The two-component eq. (4.7) is equivalent to the four-component Dirac equation

$$i \frac{\partial \Psi(t, \mathbf{x})}{\partial t} = (\gamma_0 \gamma_k p_k + \gamma_0 m) \Psi(t, \mathbf{x}), \quad k = 1, 2, 3 \quad (4.9)$$

with such subsidiary condition

$$\left(1 - \frac{\gamma_3 \gamma_4 m + \gamma_4 p_3}{q_3} \right) \Psi(t, \mathbf{x}) = 0. \quad (4.10)$$

The author expresses his gratitude to Professor O.S. Parasjuk for the benefits of many discussions.

1. Lee T.D., Yang C.N., *Phys. Rev.*, 1957, **105**, 1671;
Landau L., *Nucl. Phys.*, 1957, **3**, 127;
Salam A., *Nuovo Cimento*, 1957, **5**, 1207.
2. Fushchych W.I., Kiev, preprint ITF-70-40, 1970.
3. Shirokov Yu.M., *JETP (Sov. Phys.)*, 1958, **6**, 664.
4. Lomont J.S., Moses H.E., *J. Math. Phys.*, 1962, **3**, 405.
5. Fronsdal C., *Phys. Rev.*, 1959, **113**, 1367;
Voisin J., *Nuovo Cimento*, 1964, **34**, 1257.
6. Tokuoka Z., *Prog. Theor. Phys.*, 1967, **37**, 603;
Sen Gupta N.D., *Nucl. Phys. B*, 1968, **4**, 147;
Santhanam T.S., Chandrasekaran P.S., *Prog. Theor. Phys.*, 1969, **41**, 264.
7. Fushchych W.I., Kiev, preprint ITF-69-17, 1969.