UDC 517.9

A FRACTIONAL ORDER COVID-19 EPIDEMIC MODEL WITH MITTAG-LEFFLER KERNEL ЕПІДЕМІОЛОГІЧНА МОДЕЛЬ ДРОБОВОГО ПОРЯДКУ COVID-19 ІЗ ЯДРОМ МІТТАГ-ЛЕФФЛЕРА

H. Khan, M. Ibrahim

Dep. Math., Shaheed Benazir Bhutto Univ., Sheringal, District Dir(Upper), Khyber Pakhtunkhwa, Pakistan e-mail: hasibkhan13@yahoo.com, ibrahimsbbu1989@gmail.com

A. Khan

Dep. Math. and Gen. Sci., Prince Sultan Univ., 11586, Riyadh, Saudi Arabia e-mail: akhan@psu.edu.sa

O. Tunç

Dep. Comput. Progr., Baskale Vocational School, Van Yuzuncu Yil Univ., Van, Turkey e-mail: osmantunc89@gmail.com

Th. Abdeljawad

Dep. Math. and Gen. Sci., Prince Sultan Univ., 11586, Riyadh, Saudi Arabia, Dep. Med. Research, China Med. Univ., Taichung, Taiwan, Dep. Comput. Sci. and Inform. Eng., Asia Univ., Taichung, Taiwan e-mail: tabdeljawad@psu.edu.sa

We have considered a nonlinear fractional order Covid-19 model in the Atagana–Baleanu fractional derivative sense for the analytical and computational studies. The model consists of six classes including susceptible, protected susceptible, asymptomatic infected, symptomatic infected, quarantined and the recovered individuals. The model is studied for the existence of solution with the help of the successive iterative technique with the limit point as the solution of the model and the Hyers–Ulam stability is studied with the basic results notions. A numerical scheme is produced and tested with the support of the available literature. The graphical results show prediction of the curtail of the spread in the next 5000 days. And there is a gradual increase in the population of the protected susceptible.

Розглянуто нелінійну модель дробового порядку Covid-19 у розумінні дробової похідної Атагана – Балеану для аналітичних і комп'ютерних досліджень. Модель складається з шести класів людей, які сприйнятливі, захищено сприйнятливі, безсимптомно інфіковані, симптомно інфіковані, ізольовані на карантині та одужавші. За допомогою методу послідовних наближень досліджено існування розв'язку моделі з граничною точкою, а також стійкість Хайерса – Улама. З використанням наявної літератури запропоновано й випробувано чисельну схему. Графічні результати прогнозують скорочення поширення через 5000 днів. Також поступово збільшується кількість людей, які мають захищену сприйнятливість.

© H. Khan, M. Ibrahim, A. Khan, O. Tunç, Th. Abdeljawad, 2021 378 ISSN 1562-3076. Нелінійні коливання, 2021, т. 24, № 3 **1. Introduction.** Coronavirus infection 2019 (Covid-19) is a communicable respiratory disease. SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus) is a disease caused by a newly discovered virus strain [1]. In Wuhan, China, Covid-19 was first identified in December, 2019, and spread quickly over four months. In a short period, more than 2.9 million inhabitants in 185 nations around the world were infected and 206 thousand person died [2]. On March 11, 2020, "The World Health Organization" announced this coronavirus infections is a pandemic [3]. This disease can spread primarily from small droplets via coughing, sneezing, person-to-person or conversation. By contacting polluted surfaces, prone individuals can also be compromised. The most prevalent signs of this disease are fever, nausea, dry cough, fatigue, breath shortages. All these of signs are parts of Covid-19 [4]. Some patients can have joint pain such as nasal stuffiness, runny nose, sore throat or diarrhea. The symptoms are typically mild, but can slowly occur. In order to prevent infection, hand washing, nose covering or mouth covering while washing sneezing or coughing, avoiding nose, mouth or mouth touch and preventive steps are advised for the eyes and social distances.

Due to the seriousness of the Covid-19 pandemic, many states have done drastic decisions to curb the distribution of Covid-19 infection. In addition, they checked and covered their healthcare systems. Hence, they ruled the cancellation of public events, the closing of public events, schools, public places, borders, restrictions on travel and lockout, etc.. While those measures were helpful, that lockdown lead to socio-economic damage such as bankruptcy of many workplaces, several staff have lost their respective positions and so on. Next, the shutdown has disrupted supply chains and decreased productivity. The shutdown of China's drug-producing plants, which are the shutdown of second largest pharmaceutical product exporter, has been delayed the deliveries of generic drug processing factories [5]. The sectors of tourism, air transport, and oil were visibly influenced. It is also expected that invisible impacts are expected irrespective of the pandemic's duration. According to "The International Monetary Fund" the worldwide economy is expected to shrink by 3% in 2020 [6].

Governments are to save the failure of the economy, thinking of security measures in order to relax the lockdown. Some advanced countries intend to grant a immunity passport, which shows immunity to the illness. However, this technique has been disapproved by "The World Health Organization", since there is a lack of adequate scientific proof that reinfection is aforementioned approach is not possible. A risk balancing strategy was adopted by the South African government to lift the lockout restrictions progressively.

We refer the readers for some scientific works done on infectious diseases to [7-9], in particular, for developed several mathematical models related to Covid-19 to [10-12] and for some recent scientific works on varius fractional mathematical models [10, 13-34].

In this paper, we consider the following Covid-19 model for the existence, stability, and numerical simulations by using Atanga–Baleanu fractional derivative in the Caputo's sense. For the detail, the readers can get benefit from [29, 35];

$${}^{ABC}_{0}D^{\varphi_{1}^{*}}_{t}S = \Lambda_{1} + \gamma Q - \alpha_{1}S - \frac{(1 - \alpha_{1})(\alpha_{2}\eta_{1}I_{A} + \alpha_{3}\eta_{2}I_{S} + \eta_{3}Q)S}{S + S_{P} + I_{A} + I_{S} + Q + R} - \mu S,$$
$${}^{ABC}_{0}D^{\varphi_{2}^{*}}_{t}S_{P} = \alpha_{1}S - \mu S_{P},$$

$${}^{ABC}_{0} D^{\wp_{3}^{*}}_{t} I_{A} = \frac{(1-\alpha_{1})(\alpha_{2}\eta_{1}I_{A} + \alpha_{3}\eta_{2}I_{S} + \eta_{3}Q)S}{S+S_{P} + I_{A} + I_{S} + Q + R} - (\alpha_{2}\rho + \alpha_{2}r_{1} + 1 - \alpha_{2} + \mu)I_{A},$$

$${}^{ABC}_{0} D^{\wp_{*}}_{t} I_{S} = \alpha_{2}\rho I_{A} - (1-\alpha_{3} + \alpha_{3}r_{2} + \mu + \delta)I_{S},$$

$${}^{ABC}_{0} D^{\wp_{5}}_{t} Q = (1-\alpha_{2})I_{A} + (1-\alpha_{3})I_{S} - (\gamma + r_{3} + \mu + \delta)Q,$$

$${}^{ABC}_{0} D^{\wp_{5}}_{t} R = \alpha_{2}r_{1}I_{A} + \alpha_{3}r_{2}I_{S} + R_{3}Q - \mu R.$$

The the population is divided into six compartments. They are: S for the susceptible, S_P are the protected susceptible class, I_A are asymptomatic infected but not quarantined, I_S are the symptomatic infected not quarantined class, the quarantined class is Q and recovered R. The fractional orders $\varphi_i^* \in (0, 1]$. The parameters are: α_1 is fraction of protected susceptible, α_2 is fraction of unidentified asymptomatic infected, α_3 is fraction of unidentified symptomatic infected, η_1 contact rate between S and I_A , η_2 is contact rate between S and I_A , η_3 is contact rate between S and Q, ρ is disease progression rate from I_A to I_S , r_1 is the recovery rate of I_A , r_2 is recovery rate of I_S , r_3 is recovery rate of Q, δ is the death rate due to Covid-19 disease, γ is proportion of non-effected quarantine, μ is natural mortality rate. About the ABC-fractional calculus, we highlight the following useful literature.

Definition 1. The ABC-fractional differential operator on $\psi \in H^*(a, b)$, b > a, for $\wp_1^* \in [0, 1]$ is

$${}^{ABC}{}_{a}\mathcal{D}_{\tau}^{\wp_{1}^{*}}\psi(\tau) = \frac{B(\wp_{1}^{*})}{1-\wp_{1}^{*}}\int_{a}^{\tau}\psi'(s)E_{\wp^{*}}\left[\frac{-\wp^{*}(\tau-s)^{\wp^{*}}}{1-\wp^{*}}\right]ds,\tag{1}$$

where $B(\wp^*)$ is satisfied the property B(0) = B(1) = 1.

Definition 2. For $\psi \in H^*(a, b)$, b > a, $\wp^* \in [0, 1]$, the ABR-fractional derivative is

$${}^{ABR}{}_{a}\mathcal{D}^{\wp^*}_{\tau}\psi(\tau) = \frac{B(\wp^*)}{1-\wp^*} \frac{d}{d\tau} \int_{a}^{\tau} \psi(s) E_{\wp^*} \left[\frac{-\wp^*(\tau-s)^{\wp^*}}{1-\wp^*}\right] ds.$$

Definition 3. The AB-integral of $\psi \in H^*(a, b)$, b > a, $0 < \wp_1^* < 1$ is given by

$${}^{AB}{}_{a}\mathcal{I}_{\tau}^{\varphi_{1}^{*}}\psi(\tau) = \frac{1-\varphi_{1}^{*}}{B(\varphi_{1}^{*})}\psi(\tau) + \frac{\varphi_{1}^{*}}{B(\varphi_{1}^{*})\Gamma(\varphi_{1}^{*})}\int_{a}^{\tau}\psi(s)(\tau-s)^{\varphi_{1}^{*}-1}ds.$$

Lemma 1. The AB fractional derivative and AB fractional integral of the function ψ , satisfy the Newton – Leibniz formula

$${}^{AB}{}_{a}\mathcal{I}^{\varphi_{1}^{*}}_{\tau}\left({}^{ABC}{}_{a}\mathcal{D}^{\varphi_{1}^{*}}_{\tau}\psi(\tau)\right)=\psi(\tau)-\psi(a).$$

ISSN 1562-3076. Нелінійні коливання, 2021, т. 24, № 3

2. Existence criteria. By the AB-fractional integral and Covid-19 model (1), we have

$$\begin{split} S(t) - S(0) &= \frac{1 - \wp_1^*}{\beta(\wp_1^*)} \left[A_1 + \gamma Q - \alpha_1 S - \frac{(1 - \alpha_1)(\alpha_2 \eta_1 I_A + \alpha_3 \eta_2 I_S + \eta_3 Q)S}{S + S_P + I_A + I_S + Q + R} - \mu S \right] + \\ &+ \frac{\wp_1^*}{\beta(\wp_1^*)} [\sum_{\wp_1^*} \int_0^t (t - s)^{\wp_1^* - 1} \left[A_1 + \gamma Q - \alpha_1 S - \\ &- \frac{(1 - \alpha_1)(\alpha_2 \eta_1 I_A + \alpha_3 \eta_2 I_S + \eta_3 Q)S}{S + S_P + I_A + I_S + Q + R} - \mu S \right] ds, \\ S_p(t) - S_p(0) &= \frac{1 - \wp_2^*}{\beta(\wp_2^*)} [\alpha_1 S - \mu S_P] + \frac{\wp_1^*}{\beta(\wp_2^*)} \sum_{0}^t \int_0^t (t - s)^{\wp_2^* - 1} [\alpha_1 S - \mu S_P] ds, \\ I_A(t) - I_A(0) &= \frac{1 - \wp_3^*}{\beta(\wp_3^*)} \left[\frac{(1 - \alpha_1)(\alpha_2 \eta_1 I_A + \alpha_3 \eta_2 I_S + \eta_3 Q)S}{S + S_P + I_A + I_S + Q + R} - \\ &- (\alpha_2 \rho + \alpha_2 r_1 + 1 - \alpha_2 + \mu) I_A \right] + \frac{\wp_1^*}{\beta(\wp_3^*)} \sum_{0}^t \int_0^t (t - s)^{\wp_3^* - 1} \times \\ &\times \left[\frac{(1 - \alpha_1)(\alpha_2 \eta_1 I_A + \alpha_3 \eta_2 I_S + \eta_3 Q)S}{S + S_P + I_A + I_S + Q + R} - \\ &- (\alpha_2 \rho + \alpha_2 r_1 + 1 - \alpha_2 + \mu) I_A \right] + \frac{\wp_1^*}{\beta(\wp_3^*)} \sum_{0}^t \int_0^t (t - s)^{\wp_3^* - 1} \times \\ &\times \left[\frac{(1 - \alpha_1)(\alpha_2 \eta_1 I_A + \alpha_3 \eta_2 I_S + \eta_3 Q)S}{S + (\alpha_2 \rho + \alpha_2 r_1 + 1 - \alpha_2 + \mu) I_A} \right] ds, \\ I_s(t) - I_s(0) &= \frac{1 - \wp_1^*}{\beta(\wp_4^*)} \left[\alpha_2 \rho I_A - (1 - \alpha_3 + \alpha_3 r_2 + \mu + \delta) I_S \alpha_2 \rho I_A - \\ &- (1 - \alpha_3 + \alpha_3 r_2 + \mu + \delta) I_S \right] + \frac{\wp_4^*}{\beta(\wp_4^*)(\Gamma \wp_4^*)} \times \\ &\times \int_0^t (t - s)^{\wp_4^* - 1} \left[(\alpha_2 \rho I_A - (1 - \alpha_3 + \alpha_3 r_2 + \mu + \delta) I_S \right] ds, \\ Q(t) - Q(0) &= \frac{1 - \wp_5^*}{\beta(\wp_5^*)} \left[(1 - \alpha_2) I_A + (1 - \alpha_3) I_S - (\gamma + r_3 + \mu + \delta) Q \right] + \\ &+ \frac{\wp_6^*}{\beta(\wp_5^*)} \left[(\rho_{\wp_5^*}) \int_0^t (t - s)^{\wp_5^* - 1} \left[(\alpha_2 r_1 I_A + \alpha_3 r_2 I_S + R_3 Q - \mu R \right] ds. \\ R(t) - R(0) &= \frac{1 - \wp_5^*}{\beta(\wp_5^*)} \int_0^t (t - s)^{\wp_5^* - 1} \left[(\alpha_2 r_1 I_A + \alpha_3 r_2 I_S + R_3 Q - \mu R \right] ds. \end{split}$$

Assume the functions Y_i , i = 1, ..., 6, are given below:

$$\begin{split} Y_1(t,S) &= \Lambda_1 + \gamma Q - \alpha_1 S - \frac{(1-\alpha_1)(\alpha_2\eta_1 I_A + \alpha_3\eta_2 I_S + \eta_3 Q)S}{S + S_P + I_A + I_S + Q + R} - \mu S, \\ Y_2(t,S_p) &= \alpha_1 S - \mu S_P, \\ Y_3(t,I_A) &= \frac{(1-\alpha_1)(\alpha_2\eta_1 I_A + \alpha_3\eta_2 I_S + \eta_3 Q)S}{S + S_P + I_A + I_S + Q + R} - (\alpha_2 \rho + \alpha_2 r_1 + 1 - \alpha_2 + \mu)I_A, \\ Y_4(t,I_s) &= \alpha_2 \rho I_A - (1-\alpha_3 + \alpha_3 r_2 + \mu + \delta)I_S, \\ Y_5(t,Q) &= (1-\alpha_2)I_A + (1-\alpha_3)I_S - (\gamma + r_3 + \mu + \delta)Q, \\ Y_6(t,R) &= \alpha_2 r_1 I_A + \alpha_3 r_2 I_S + R_3 Q - \mu R, \\ \begin{cases} \psi_1 &= \alpha_1 + k_1 + \mu, \\ \psi_2 &= \mu, \\ \psi_3 &= k_2 + (\alpha_2 \rho + \alpha_2 r_1 + 1 - \alpha_2 + \mu), \\ \psi_4 &= \mu_c, \\ \psi_5 &= 1 - \alpha_3 + \alpha_3 r_2 + \mu + \delta, \\ \psi_6 &= \mu. \end{split}$$

Assumption (B). We assume that, for S(t), $S^*(t)$, $S_p(t)$, $S_p^*(t)$, $I_A(t)$, $I_A^*(t)$, $I_s(t)$, $I_s^*(t)$, Q(t), $Q^*(t)$, R(t), $R^*(t) \in L[0,1]$, there exists constants $\kappa_i > 0$, i = 1, ..., 6, such that $||S(t)|| \le \kappa_1$, $||S_p(t)|| \le \kappa_2$, $||I_A(t)|| \le \kappa_3$, $||I_s(t)|| \le \kappa_4$, $||Q(t)|| \le \kappa_5$, $||R(t)|| \le \kappa_6$, and ξ_1 , $\xi_2 > 0$, and

$$||S(t) + I_A(t) + Q(t)|| \le \xi_1,$$
$$||I_s(t) + R(t)|| \le \xi_2.$$

Theorem 1. The Y_i , $i \in N_1^6$, satisfy Lipschitz condition provided that Assumption (B) is obeyed.

Proof. Consider for Y_1 , below

$$\begin{split} \left\| Y_{1}(t,S) - Y_{1}(t,S^{*}) \right\| &= \left\| \Lambda_{1} + \gamma Q - \alpha_{1}S - \frac{(1 - \alpha_{1})(\alpha_{2}\eta_{1}I_{A} + \alpha_{3}\eta_{2}I_{S} + \eta_{3}Q)S}{S + S_{P} + I_{A} + I_{S} + Q + R} - \mu S - \left(\Lambda_{1} + \gamma Q - \alpha_{1}S^{*} - \frac{(1 - \alpha_{1})(\alpha_{2}\eta_{1}I_{A} + \alpha_{3}\eta_{2}I_{S} + \eta_{3}Q)S^{*}}{S^{*} + S_{P} + I_{A} + I_{S} + Q + R} - \mu S^{*} \right) \right\| \leq \\ &\leq \left\| \alpha_{1} + \frac{(1 - \alpha_{1})(\alpha_{2}\eta_{1}I_{A} + \alpha_{3}\eta_{2}I_{S} + \eta_{3}Q)S^{*}}{S^{*} + S_{P} + I_{A} + I_{S} + Q + R} + \mu \right\| \left\| S - S^{*} \right\| \leq \\ \end{split}$$

$$\leq [\alpha_1 + k_1 + \mu] \|S_c - S_c^*\| = \psi_1 \|S - S^*\|.$$
⁽²⁾

For the $Y_2(t, E_c)$, we have

$$\|Y_{2}(t, S_{p}) - Y_{2}(t, S_{p}^{*})\| = \|(\alpha_{1}S - \mu S_{P}) - (\alpha_{1}S - \mu S_{P}^{*},)\| \le \le [\mu] \|S_{c} - E_{c}^{*}\| \le \psi_{2} \|E_{c} - E_{c}^{*}\|.$$
(3)

The $Y_3(t, I_A^*)$ implies

$$\|Y_{3}(t, I_{A}) - Y_{3}(t, I_{A}^{*})\| = \left\| \left(\frac{(1 - \alpha_{1})(\alpha_{2}\eta_{1}I_{A} + \alpha_{3}\eta_{2}I_{S} + \eta_{3}Q)S}{S + S_{P} + I_{A} + I_{S} + Q + R} - (\alpha_{2}\rho + \alpha_{2}r_{1} + 1 - \alpha_{2} + \mu)I_{A} \right) - (\alpha_{2}\rho + \alpha_{2}r_{1} + 1 - \alpha_{2} + \mu)I_{A} \right) - (\alpha_{2}\rho + \alpha_{2}r_{1} + 1 - \alpha_{2} + \mu)I_{A}^{*} \right) \right\| \leq (1 - \alpha_{1})(\alpha_{2}\eta_{1}I_{A} + \alpha_{3}\eta_{2}I_{S} + \eta_{3}Q)S) + (\alpha_{2}\rho + \alpha_{2}r_{1} + 1 - \alpha_{2} + \mu)I_{A}^{*}) \| \leq (1 - \alpha_{1})(\alpha_{2}\eta_{1}I_{A} + \alpha_{3}\eta_{2}I_{S} + \eta_{3}Q)S) + (\alpha_{2}\rho + \alpha_{2}r_{1} + 1 - \alpha_{2} + \mu)I_{A}^{*}) \| \leq (1 - \alpha_{2})(\alpha_{2}\rho + \alpha_{2}r_{1} + 1 - \alpha_{2} + \mu)\| \|I_{A} - I_{A}^{*}\| \leq (k_{2} + (\alpha_{2}\rho + \alpha_{2}r_{1} + 1 - \alpha_{2} + \mu))\| \|I_{c} - I_{c}^{*}\| = \psi_{3} \|I_{A} - I_{A}^{*}\|$$

$$(4)$$

for $Y_4(t, I)$. We obtain

$$\|Y_{4}(t, I_{s}) - Y_{4}(t, I_{s}^{*})\| = \left\| (\alpha_{2}\rho I_{A} - (1 - \alpha_{3} + \alpha_{3}r_{2} + \mu + \delta)I_{S}) - (\alpha_{2}\rho I_{A} - (1 - \alpha_{3} + \alpha_{3}r_{2} + \mu + \delta)I_{S}^{*}) \right\| \leq \\ \leq \|1 - \alpha_{3} + \alpha_{3}r_{2} + \mu + \delta\| \|I_{s} - I_{s}^{*}\| \leq \\ \leq \left[1 - \alpha_{3} + \alpha_{3}r_{2} + \mu + \delta \right] \|I_{s} - I_{s}^{*}\| \leq \psi_{4} \|I_{s} - I_{s}^{*}\|$$
(5)

for $Y_5(t, Q)$.

We get

$$\|Y_5(t,Q) - Y_5(t,Q^*)\| = \left\| \left((1-\alpha_2)I_A + (1-\alpha_3)I_S - (\gamma+r_3+\mu+\delta)Q \right) - ((1-\alpha_2)I_A + (1-\alpha_3)I_S - (\gamma+r_3+\mu+\delta)Q^*) \right\| \le C_1 + C_2 +$$

ISSN 1562-3076. Нелінійні коливання, 2021, т. 24, № 3

H. KHAN, M. IBRAHIM, A. KHAN, O. TUNÇ, TH. ABDELJAWAD

$$\leq \|\gamma + r_3 + \mu + \delta\| \|Q - Q^*\| = \psi_5 \|Q - Q^*\|$$
(6)

for $Y_6(t, R)$.

Further, we have

$$\|Y_{6}(t,R) - Y_{6}(t,R^{*})\| = \left\| (\alpha_{2}r_{1}I_{A} + \alpha_{3}r_{2}I_{S} + r_{3}Q - \mu R,) - (\alpha_{2}r_{1}I_{A} + \alpha_{3}r_{2}I_{S} + r_{3}Q - \mu R^{*}) \right\| \leq \\ \leq \|\mu\| \|R - R^{*}\| = \psi_{6} \|R - R^{*}\|.$$
(7)

Thus, from (2) to (7), we have that the Y_i , i = 1, ..., 6, satisfy the Lipschitz condition. And this completes the proof.

Assuming that $S(0) = S_p(0) = I_A(0) = I_S(0) = Q(0) = R(0) = 0$, then we have

$$S(t) = \frac{1 - \wp_1^*}{\beta(\wp_1^*)} \mathcal{Y}_1(t, S(t)) + \frac{\wp_1^*}{\beta(\wp_1^*)\Gamma(\wp_1^*)} \int_0^t (t - s)^{\wp_1^* - 1} \mathcal{Y}_1(s, S(s)) ds,$$
(8)

$$S_{p}(t) = \frac{1 - \wp_{2}^{*}}{\beta(\wp_{2}^{*})} \mathcal{Y}_{2}(t, S_{p}(t)) + \frac{\wp_{2}^{*}}{\beta(\wp_{2}^{*})\Gamma(\wp_{2}^{*})} \int_{0}^{t} (t - s)^{\wp_{2}^{*} - 1} \mathcal{Y}_{2}(s, S_{p}(s)) ds,$$
(9)

$$I_A(t) = \frac{1 - \wp_3^*}{\beta(\wp_3^*)} \mathcal{Y}_3(t, I_A(t)) + \frac{\wp_3^*}{\beta(\wp_3^*) \Gamma(\wp_3^*)} \int_0^t (t - s)^{\wp_3^* - 1} \mathcal{Y}_3(s, I_A(s)) ds,$$
(10)

$$I_{s}(t) = \frac{1 - \wp_{4}^{*}}{\beta(\wp_{4}^{*})} \mathcal{Y}_{4}(t, I_{s}(t)) + \frac{\wp_{4}^{*}}{\beta(\wp_{4}^{*})\Gamma(\wp_{4}^{*})} \int_{0}^{t} (t - s)^{\wp_{4}^{*} - 1} \mathcal{Y}_{4}(s, I_{s}(s)) ds,$$
(11)

$$Q(t) = \frac{1 - \wp_5^*}{\beta(\wp_5^*)} \mathcal{Y}_5(t, Q(t)) + \frac{\wp_5^*}{\beta(\wp_5^*) \Gamma(\wp_5^*)} \int_0^t (t - s)^{\wp_5^* - 1} \mathcal{Y}_5(s, Q(s)) ds,$$
(12)

$$R(t) = \frac{1 - \wp_6^*}{\beta(\wp_6^*)} \mathcal{Y}_6(t, R(t)) + \frac{\wp_6^*}{\beta(\wp_6^*) \Gamma(\wp_6^*)} \int_0^t (t - s)^{\wp_6^* - 1} \mathcal{Y}_6(s, R(s)) ds.$$
(13)

For the iterative scheme of the model (1), we define

$$S_{n}(t) = \frac{1 - \wp_{1}^{*}}{\beta(\wp_{1}^{*})} \mathcal{Y}_{1}(t, S_{n-1}(t)) + \frac{\wp_{1}^{*}}{\beta(\wp_{1}^{*})\Gamma(\wp_{1}^{*})} \int_{0}^{t} (t - s)^{\wp_{1}^{*} - 1} \mathcal{Y}_{1}(s, S_{n-1}(s)) ds,$$

$$S_{p_{n}}(t) = \frac{1 - \wp_{2}^{*}}{\beta(\wp_{2}^{*})} \mathcal{Y}_{2}(t, S_{p_{n-1}}(t)) + \frac{\wp_{2}^{*}}{\beta(\wp_{2}^{*})\Gamma(\wp_{2}^{*})} \int_{0}^{t} (t - s)^{\wp_{2}^{*} - 1} \mathcal{Y}_{2}(s, S_{p_{n-1}}(s)) ds,$$

$$I_{An}(t) = \frac{1 - \wp_{3}^{*}}{\beta(\wp_{3}^{*})} \mathcal{Y}_{3}(t, I_{An-1}(t)) + \frac{\wp_{3}^{*}}{\beta(\wp_{3}^{*})\Gamma(\wp_{3}^{*})} \int_{0}^{t} (t - s)^{\wp_{3}^{*} - 1} \mathcal{Y}_{3}(s, I_{An-1}(s)) ds,$$

ISSN 1562-3076. Нелінійні коливання, 2021, т. 24, № 3

$$I_{sn}(t) = \frac{1 - \wp_4^*}{\beta(\wp_4^*)} \mathcal{Y}_4(t, I_{sn-1}(t)) + \frac{\wp_4^*}{\beta(\wp_4^*)} \int_0^t (t-s)^{\wp_4^*-1} \mathcal{Y}_4(s, I_{sn-1}(s)) ds,$$
$$Q_n(t) = \frac{1 - \wp_5^*}{\beta(\wp_5^*)} \mathcal{Y}_5(t, Q_{n-1}(t)) + \frac{\wp_5^*}{\beta(\wp_5^*)} \int_0^t (t-s)^{\wp_5^*-1} \mathcal{Y}_5(s, Q_{n-1}(s)) ds,$$
$$R_n(t) = \frac{1 - \wp_6^*}{\beta(\wp_6^*)} \mathcal{Y}_6(t, R_{n-1}(t)) + \frac{\wp_6^*}{\beta(\wp_6^*)} \int_0^t (t-s)^{\wp_6^*-1} \mathcal{Y}_6(s, R_{n-1}(s)) ds.$$

Theorem 2. The fractional order Covid-19 model (1) has a solution if we have

$$\Delta = \max\{\Psi_i\} < 1, \quad i \in N_1^6.$$

Proof. We define the function

$$\begin{aligned} &\mathcal{K}1_n(t) = S_{n+1}(t) - S(t), \quad \mathcal{K}2_n(t) = S_{p_{n+1}}(t) - S_p(t), \quad \mathcal{K}3_n(t) = I_{An+1}(t) - I_A(t), \\ &\mathcal{K}4_n(t) = I_{sn+1}(t) - I_s(t), \quad \mathcal{K}5_n(t) = Q_{n+1}(t) - Q(t), \quad \mathcal{K}6_n(t) = R_{n+1}(t) - R(t). \end{aligned}$$

Then, by using the above equations, we find that

$$\begin{aligned} \|\mathcal{K}\mathbf{1}_{n}\| &\leq \frac{1-\wp_{1}^{*}}{\beta(\wp_{1}^{*})} \|\mathcal{Y}_{1}(t,\mathcal{S}_{n}(t)) - \mathcal{Y}_{1}(t,S_{n-1}(t))\| + \\ &+ \frac{\wp_{1}^{*}}{\beta(\wp_{1}^{*})\Gamma(\wp_{1}^{*})} \int_{0}^{t} (t-s)^{\wp_{1}^{*}-1} \|\mathcal{Y}_{1}(s,S_{n}(s)) - \mathcal{Y}_{1}(t,S_{n-1}(t))\| ds \leq \\ &\leq \left[\frac{1-\wp_{1}^{*}}{\beta(\wp_{1}^{*})} + \frac{1}{\beta(\wp_{1}^{*})\Gamma(\wp_{1}^{*})}\right] \psi_{1}\|S_{n} - S\| \leq \\ &\leq \left[\frac{1-\wp_{1}^{*}}{\beta(\wp_{1}^{*})} + \frac{1}{\beta(\wp_{1}^{*})\Gamma(\wp_{1}^{*})}\right]^{n} \Delta^{n}\|S_{1} - S\| \end{aligned}$$

and

$$\begin{split} \|\mathcal{K}2_{n}\| &\leq \frac{1-\wp_{2}^{*}}{\beta(\wp_{2}^{*})} \|\mathcal{Y}_{2}(t, S_{p_{n}}(t)) - \mathcal{Y}_{2}(t, S_{p_{n-1}}(t))\| + \\ &+ \frac{\wp_{2}^{*}}{\beta(\wp_{2}^{*})\Gamma(\wp_{2}^{*})} \int_{0}^{t} (t-s)^{\wp_{2}^{*}-1} \|\mathcal{Y}_{2}(s, S_{p_{n}}(s)) - \mathcal{Y}_{2}(t, S_{p_{n-1}}(t))\| ds \leq \\ &\leq \left[\frac{1-\wp_{2}^{*}}{\beta(\wp_{2}^{*})} + \frac{1}{\beta(\wp_{2}^{*})\Gamma(\wp_{2}^{*})}\right] \psi_{2} \|S_{p_{n}} - S_{p}\| \leq \\ &\leq \left[\frac{1-\wp_{2}^{*}}{\beta(\wp_{2}^{*})} + \frac{1}{\beta(\wp_{2}^{*})\Gamma(\wp_{2}^{*})}\right]^{n} \Delta^{n} \|S_{p_{1}} - S_{p}\|. \end{split}$$

Similarly,

$$\begin{split} \|\mathcal{K}3_{n}\| &\leq \frac{1-\varphi_{3}^{*}}{\beta(\varphi_{3}^{*})} \|\mathcal{Y}_{3}(t,I_{An}(t)) - \mathcal{Y}_{3}(t,I_{An-1}(t))\| + \\ &+ \frac{\varphi_{3}^{*}}{\beta(\varphi_{3}^{*})} \sum_{0}^{t} (t-s)^{\varphi_{3}^{*}-1} \|\mathcal{Y}_{3}(s,I_{An}(s)) - \mathcal{Y}_{3}(t,I_{An-1}(t))\| ds \leq \\ &\leq \left[\frac{1-\varphi_{3}^{*}}{\beta(\varphi_{3}^{*})} + \frac{1}{\beta(\varphi_{1}^{*})\Gamma(\varphi_{3}^{*})}\right]^{n} \Delta^{n} \|I_{An} - I_{A}\| \leq \\ &\leq \left[\frac{1-\varphi_{3}^{*}}{\beta(\varphi_{3}^{*})} + \frac{1}{\beta(\varphi_{1}^{*})\Gamma(\varphi_{3}^{*})}\right]^{n} \Delta^{n} \|I_{An} - I_{A}\|, \\ \|\mathcal{K}4_{n}\| &\leq \frac{1-\varphi_{4}^{*}}{\beta(\varphi_{4}^{*})} \|\mathcal{Y}_{4}(t,I_{sn}(t)) - \mathcal{Y}_{4}(t,I_{sn-1}(t))\| + \\ &+ \frac{\varphi_{4}^{*}}{\beta(\varphi_{4}^{*})\Gamma(\varphi_{4}^{*})} \int_{0}^{t} (t-s)^{\varphi_{4}^{*}-1} \|\mathcal{Y}_{4}(s,I_{sn}(s)) - \mathcal{Y}_{4}(t,I_{sn-1}(t))\| ds \leq \\ &\leq \left[\frac{1-\varphi_{4}^{*}}{\beta(\varphi_{4}^{*})} + \frac{1}{\beta(\varphi_{4}^{*})\Gamma(\varphi_{4}^{*})}\right]^{n} \Delta^{n} \|I_{sn} - I_{s}\| \leq \\ &\leq \left[\frac{1-\varphi_{4}^{*}}{\beta(\varphi_{4}^{*})} + \frac{1}{\beta(\varphi_{4}^{*})\Gamma(\varphi_{4}^{*})}\right]^{n} \Delta^{n} \|I_{sn} - I_{s}\|, \\ \|\mathcal{K}5_{n}\| &\leq \frac{1-\varphi_{5}^{*}}{\beta(\varphi_{5}^{*})} \|\mathcal{Y}_{5}(t,Q_{n}(t)) - \mathcal{Y}_{5}(t,Q_{n-1}(t))\| + \\ &+ \frac{\varphi_{5}^{*}}{\beta(\varphi_{1}^{*})\Gamma(\varphi_{5}^{*})} \int_{0}^{t} (t-s)^{\varphi_{5}^{*}-1} \|\mathcal{Y}_{5}(s,Q_{n}(s)) - \mathcal{Y}_{5}(t,Q_{n-1}(t))\| ds \leq \\ &\leq \left[\frac{1-\varphi_{5}^{*}}{\beta(\varphi_{5}^{*})} + \frac{1}{\beta(\varphi_{5}^{*})\Gamma(\varphi_{5}^{*})}\right]^{n} \Delta^{n} \|Q_{1} - Q\| \leq \\ &\leq \left[\frac{1-\varphi_{5}^{*}}{\beta(\varphi_{5}^{*})} \|\mathcal{Y}_{6}(t,R_{n}(t)) - \mathcal{Y}_{6}(t,R_{n-1}(t))\| + \\ &+ \frac{\varphi_{6}^{*}}{\beta(\varphi_{6}^{*})} \|\mathcal{Y}_{6}(t,R_{n}(t)) - \mathcal{Y}_{6}(t,R_{n-1}(t))\| ds \leq \\ &\leq \left[\frac{1-\varphi_{5}^{*}}{\beta(\varphi_{6}^{*})} + \frac{1}{\beta(\varphi_{6}^{*})} \Gamma(\varphi_{6}^{*})}\right]^{t} \psi_{6}\|R_{n} - R\| \leq \\ \end{aligned}$$

ISSN 1562-3076. Нелінійні коливання, 2021, т. 24, № 3

$$\leq \left[\frac{1-\wp_6^*}{\beta(\wp_6^*)} + \frac{1}{\beta(\wp_6^*)\Gamma(\wp_6^*)}\right]^n \Delta^n \|R_1 - R\|.$$

Thus, we have $\mathcal{K}(t)_n \to 0, \ i \in 1, \dots, 6$, as $n \to \infty$ for $\Delta < 1$, which is the required proof.

3. Uniqueness solution. For our suggested model (1), we study the analysis of the uniqueness of solution.

Theorem 3. The Covid-19 model (1) has unique solution if

$$\left[\frac{1-\varphi_i}{\beta(\varphi_i)} + \frac{1}{\beta(\varphi_i)\Gamma(\varphi_i)}\right]\psi_i \le 1, \quad i \in \mathcal{N}_1^6.$$
(14)

Proof. Let there exist another solution $\overline{S}(t)$, $\overline{S_c}(t)$, $\overline{I_A}(t)$, $\overline{i_s}(t)$, $\overline{Q}(t)$, $\overline{R}(t)$ such that

$$\begin{split} \overline{S}(t) &= \frac{1 - \varphi_1^*}{\beta(\varphi_1^*)} \,\mathcal{Y}_1(t, \overline{S}(t)) + \frac{\varphi_1^*}{\beta(\varphi_1^*) \Gamma(\varphi_1^*)} \int_0^t (t-s)^{\varphi_1^*-1} \mathcal{Y}_1(s, \overline{S}(s)) ds, \\ \overline{S_p}(t) &= \frac{1 - \varphi_2^*}{\beta(\varphi_2^*)} \,\mathcal{Y}_2(t, \overline{S_p}(t)) + \frac{\varphi_2^*}{\beta(\varphi_2^*) \Gamma(\varphi_2^*)} \int_0^t (t-s)^{\varphi_2^*-1} \mathcal{Y}_2(s, \overline{S_p}(s)) ds, \\ \overline{I_A}(t) &= \frac{1 - \varphi_3^*}{\beta(\varphi_3^*)} \,\mathcal{Y}_3(t, \overline{I_A}(t)) + \frac{\varphi_3^*}{\beta(\varphi_3^*) \Gamma(\varphi_3^*)} \int_0^t (t-s)^{\varphi_3^*-1} \mathcal{Y}_3(s, \overline{I_A}(s)) ds, \\ \overline{I_s}(t) &= \frac{1 - \varphi_4^*}{\beta(\varphi_4^*)} \,\mathcal{Y}_4(t, \overline{I_s}(t)) + \frac{\varphi_4^*}{\beta(\varphi_4^*) \Gamma(\varphi_4^*)} \int_0^t (t-s)^{\varphi_4^*-1} \mathcal{Y}_4(s, \overline{I_s}(s)) ds, \\ \overline{Q}(t) &= \frac{1 - \varphi_5^*}{\beta(\varphi_5^*)} \,\mathcal{Y}_5(t, \overline{Q}(t)) + \frac{\varphi_5^*}{\beta(\varphi_5^*) \Gamma(\varphi_5^*)} \int_0^t (t-s)^{\varphi_5^*-1} \mathcal{Y}_5(s, \overline{Q}(s)) ds, \\ \overline{R}(t) &= \frac{1 - \varphi_6^*}{\beta(\varphi_6^*)} \,\mathcal{Y}_6(t, \overline{R}(t)) + \frac{\varphi_6^*}{\beta(\varphi_6^*) \Gamma(\varphi_6^*)} \int_0^t (t-s)^{\varphi_6^*-1} \mathcal{Y}_6(s, \overline{R}(s)) ds. \end{split}$$

Then,

$$\begin{split} \|S(t) - \overline{S}(t)\| &\leq \frac{1 - \wp_1^*}{\beta(\wp_1^*)} \|\mathcal{Y}_1(t, S(t)) - Y_1(t, \overline{S}(t))\| + \\ &+ \frac{\wp_1^*}{\beta(\wp_1^*)\Gamma(\wp_1^*)} \int_0^t (t - s)^{\wp_1^* - 1} \|Y_1(s, S(s)) - Y_1(t, \overline{S}(t))\| ds \leq \\ &\leq \left[\frac{1 - \wp_1^*}{\beta(\wp_1^*)} + \frac{1}{\beta(\wp_1^*)\Gamma(\wp_1^*)}\right] \psi_1 \|S - \overline{S}\|, \end{split}$$

which implies

$$\left[\frac{1-\wp_1^*}{\beta(\wp_1^*)}\psi_1 + \frac{\psi_1}{\beta(\wp_1^*)\Gamma(\wp_1^*)} - 1\right] \|S_-\overline{S}\| \ge 0.$$
(15)

By (14), the (15) is true if $||S - \overline{S}|| = 0$, which implies $S(t) = \overline{S}(t)$. Similarly, we have

$$\begin{split} \left\| S_{p}(t) - \overline{S_{p}}(t) \right\| &\leq \frac{1 - \varphi_{2}^{*}}{\beta(\varphi_{2}^{*})} \left\| \mathcal{Y}_{2}(t, S_{p}(t)) - Y_{2}\left(t, \overline{S_{p}}(t)\right) \right\| + \\ &+ \frac{\varphi_{2}^{*}}{\beta(\varphi_{2}^{*})\Gamma(\varphi_{2}^{*})} \int_{0}^{t} (t - s)^{\varphi_{2}^{*} - 1} \left\| Y_{2}(s, S_{p}(s)) - Y_{2}(t, \overline{S_{p}}(t)) \right\| ds \leq \\ &\leq \left[\frac{1 - \varphi_{2}^{*}}{\beta(\varphi_{2}^{*})} + \frac{1}{\beta(\varphi_{2}^{*})\Gamma(\varphi_{2}^{*})} \right] \psi_{2} \left\| S_{P} - \overline{S_{p}} \right\|, \\ &\left[\frac{1 - \varphi_{2}^{*}}{\beta(\varphi_{2}^{*})} \psi_{2} + \frac{\psi_{2}}{\beta(\varphi_{2}^{*})\Gamma(\varphi_{2}^{*})} - 1 \right] \left\| S_{P} - \overline{S_{p}} \right\| \geq 0, \end{split}$$
(16)

which follows. By (14), the (16) is true if $||S_p - \overline{S_p}|| = 0$, which implies $S_p(t) = \overline{S_p}(t)$. Now, for I_A , we have

$$\begin{split} \left\| I_A(t) - \overline{I_A}(t) \right\| &\leq \frac{1 - \wp_3^*}{\beta(\wp_3^*)} \left\| \mathcal{Y}_3(t, I_A(t)) - Y_3(t, \overline{I_A}(t)) \right\| + \\ &+ \frac{\wp_3^*}{\beta(\wp_3^*) \Gamma(\wp_3^*)} \int_0^t (t - s)^{\wp_3^* - 1} \left\| Y_3(s, I_A(s)) - Y_3(t, \overline{I_A}(t)) \right\| ds \leq \\ &\leq \left[\frac{1 - \wp_3^*}{\beta(\wp_3^*)} + \frac{1}{\beta(\wp_3^*) \Gamma(\wp_3^*)} \right] \psi_3 \left\| I_A - \overline{I_A} \right\|, \end{split}$$

which implies

$$\left[\frac{1-\wp_3^*}{\beta(\wp_3^*)}\psi_3 + \frac{\psi_3}{\beta(\wp_3^*)\Gamma(\wp_3^*)} - 1\right] \left\|I_A - \overline{I_A}\right\| \ge 0,\tag{17}$$

which implies by (14), the (17) is true if $||I_A - \overline{I_A}|| = 0$, which implies $I_A(t) = \overline{I_A}(t)$;

$$\begin{split} \left\| I_s(t) - \overline{I_s}(t) \right\| &\leq \frac{1 - \wp_4^*}{\beta(\wp_4^*)} \left\| \mathcal{Y}_4(t, I_s(t)) - Y_4(t, \overline{I_s}(t)) \right\| + \\ &+ \frac{\wp_4^*}{\beta(\wp_4^*) \Gamma(\wp_4^*)} \int_0^t (t - s)^{\wp_4^* - 1} \left\| Y_4(s, I_s(s)) - Y_4(t, \overline{I_s}(t)) \right\| ds \leq \end{split}$$

ISSN 1562-3076. Нелінійні коливання, 2021, т. 24, № 3

$$\leq \left[\frac{1-\wp_4^*}{\beta(\wp_4^*)} + \frac{1}{\beta(\wp_4^*)\Gamma(\wp_4^*)}\right]\psi_4 \left\|I_s - \overline{I_s}\right\|,$$

which implies

$$\left[\frac{1-\wp_4^*}{\beta(\wp_4^*)}\psi_4 + \frac{\psi_4}{\beta(\wp_4^*)\Gamma(\wp_4^*)} - 1\right] \left\|I_{s-}\overline{I_s}\right\| \ge 0.$$
(18)

By (14), the (18) is true if $||I_s - \overline{I_s}|| = 0$, which implies $I_s(t) = \overline{I_s}(t)$. Now, for Q, we have

$$\begin{split} \left\|Q(t) - \overline{Q}(t)\right\| &\leq \frac{1 - \wp_5^*}{\beta(\wp_5^*)} \left\|\mathcal{Y}_5(t, Q(t)) - Y_5(t, \overline{Q}(t))\right\| + \\ &+ \frac{\wp_5^*}{\beta(\wp_5^*)\Gamma(\wp_5^*)} \int_0^t (t-s)^{\wp_5^*-1} \left\|Y_5(s, Q(s)) - Y_5\left(t, \overline{Q}(t)\right)\right\| ds \leq \\ &\leq \left[\frac{1 - \wp_5^*}{\beta(\wp_5^*)} + \frac{1}{\beta(\wp_5^*)\Gamma(\wp_5^*)}\right] \psi_5 \left\|Q - \overline{Q}\right\|, \end{split}$$

which implies

$$\begin{split} \left[\frac{1 - \varphi_5^*}{\beta(\varphi_5^*)} \psi_5 + \frac{\psi_5}{\beta(\varphi_5^*) \Gamma(\varphi_5^*)} - 1 \right] \|Q - \overline{Q}\| \ge 0, \\ \|R(t) - \overline{R}(t)\| \le \frac{1 - \varphi_6^*}{\beta(\varphi_6^*)} \left\| \mathcal{Y}_6(t, R(t)) - Y_6(t, \overline{R}(t)) \right\| + \\ &+ \frac{\varphi_6^*}{\beta(\varphi_6^*) \Gamma(\varphi_6^*)} \int_0^t (t - s)^{\varphi_6^* - 1} \left\| Y_6(s, R(s)) - Y_6(t, \overline{R}(t)) \right\| ds \le \\ &\le \left[\frac{1 - \varphi_6^*}{\beta(\varphi_6^*)} + \frac{1}{\beta(\varphi_6^*) \Gamma(\varphi_6^*)} \right] \psi_6 \left\| R - \overline{R} \right\|, \end{split}$$

which implies

$$\left[\frac{1-\wp_6^*}{\beta(\wp_6^*)}\psi_6 + \frac{\psi_6}{\beta(\wp_6^*)\Gamma(\wp_6^*)} - 1\right] \left\|R - \overline{R}\right\| \ge 0,\tag{19}$$

which implies by (14), the (19) is true if $||R - \overline{R}|| = 0$, which implies $R(t) = \overline{R}(t)$. Thus the (1) has unique solution.

4. Hyers – Ulams stability.

Definition 4. The integral system (8)–(13) is Hyers–Ulam stable if for $\Delta_i > 0$, $i \in \mathcal{N}_1^6$, and $\gamma_i > 0$, $i \in \mathcal{N}_1^6$, such that

$$\left| S(t) - \frac{1 - \wp_1^*}{\beta(\wp_1^*)} \mathcal{Y}_1(t, S(t)) - \frac{\wp_1^*}{\beta(\wp_1^*) \Gamma(\wp_1^*)} \int_0^t (t - s)^{\wp_1^* - 1} \mathcal{Y}_1(s, S(s)) ds \right| \le \gamma_1,$$

ISSN 1562-3076. Нелінійні коливання, 2021, т. 24, № 3

$$\begin{aligned} \left| S_{p}(t) - \frac{1 - \wp_{2}^{*}}{\beta\left(\wp_{2}^{*}\right)} \mathcal{Y}_{2}(t, S_{p}(t)) - \frac{\wp_{2}^{*}}{\beta\left(\wp_{2}^{*}\right)} \int_{0}^{t} (t - s)^{\wp_{2}^{*} - 1} \mathcal{Y}_{2}(s, S_{p}(s)) ds \right| &\leq \gamma_{2}, \\ \left| I_{A}(t) - \frac{1 - \wp_{3}^{*}}{\beta\left(\wp_{3}^{*}\right)} \mathcal{Y}_{3}(t, I_{A}(t)) - \frac{\wp_{3}^{*}}{\beta\left(\wp_{3}^{*}\right)} \int_{0}^{t} (t - s)^{\wp_{3}^{*} - 1} \mathcal{Y}_{3}(s, I_{A}(s)) ds \right| &\leq \gamma_{3}, \\ \left| I_{s}(t) - \frac{1 - \wp_{4}^{*}}{\beta\left(\wp_{4}^{*}\right)} \mathcal{Y}_{4}(t, I_{s}(t)) - \frac{\wp_{4}^{*}}{\beta\left(\wp_{4}^{*}\right)} \int_{0}^{t} (t - s)^{\wp_{4}^{*} - 1} \mathcal{Y}_{4}(s, I_{s}(s)) ds \right| &\leq \gamma_{4}, \\ \left| Q(t) - \frac{1 - \wp_{5}^{*}}{\beta\left(\wp_{5}^{*}\right)} \mathcal{Y}_{5}(t, Q(t)) - \frac{\wp_{5}^{*}}{\beta\left(\wp_{5}^{*}\right)} \int_{0}^{t} (v_{5}^{*}) \int_{0}^{t} (t - s)^{\wp_{5}^{*} - 1} \mathcal{Y}_{5}(s, Q(s)) ds \right| &\leq \gamma_{5}, \\ \left| R(t) - \frac{1 - \wp_{5}^{*}}{\beta\left(\wp_{5}^{*}\right)} \mathcal{Y}_{6}(t, Q(t)) - \frac{\wp_{5}^{*}}{\beta\left(\wp_{5}^{*}\right)} \int_{0}^{t} (t - s)^{\wp_{5}^{*} - 1} \mathcal{Y}_{6}(s, R(s)) ds \right| &\leq \gamma_{5}. \end{aligned} \right|$$

We have $\dot{S}(t)$, $\dot{S}_p(t)$, $\dot{I}_A(t)$, $\dot{I}_s(t)$, $\dot{Q}(t)$, $\dot{R}(t)$, which implies

$$\begin{split} \dot{S}(t) &= \frac{1 - \wp_1^*}{\beta(\wp_1^*)} \,\mathcal{Y}_1(t, \dot{S}(t)) + \frac{\wp_1^*}{\beta(\wp_1^*)} \int_0^t (t-s)^{\wp_1^*-1} \mathcal{Y}_1(s, \dot{S}(s)) ds, \\ \dot{S}_p(t) &= \frac{1 - \wp_2^*}{\beta(\wp_2^*)} \mathcal{Y}_2(t, \dot{S}_p(t)) + \frac{\wp_2^*}{\beta(\wp_2^*)} \int_0^t (t-s)^{\wp_2^*-1} \mathcal{Y}_2(s, \dot{S}_p(s)) ds, \\ \dot{I}_A(t) &= \frac{1 - \wp_3^*}{\beta(\wp_3^*)} \,\mathcal{Y}_3(t, \dot{I}_A(t)) + \frac{\wp_3^*}{\beta(\wp_3^*)} \int_0^t (t-s)^{\wp_3^*-1} \mathcal{Y}_3\left(s, \dot{I}_A(s)\right) ds, \\ \dot{I}_s(t) &= \frac{1 - \wp_4^*}{\beta(\wp_4^*)} \,\mathcal{Y}_4\left(t, \dot{I}_s(t)\right) + \frac{\wp_4^*}{\beta(\wp_4^*)} \int_0^t (t-s)^{\wp_4^*-1} \mathcal{Y}_4\left(s, \dot{I}_s(s)\right) ds, \\ \dot{Q}(t) &= \frac{1 - \wp_4^*}{\beta(\wp_5^*)} \,\mathcal{Y}_5(t, \dot{Q}(t)) + \frac{\wp_5^*}{\beta(\wp_5^*)} \int_0^t (t-s)^{\wp_5^*-1} \mathcal{Y}_5\left(s, \dot{Q}(s)\right) ds, \\ \dot{R}(t) &= \frac{1 - \wp_6^*}{\beta(\wp_6^*)} \,\mathcal{Y}_6\left(t, \dot{R}(t)\right) + \frac{\wp_6^*}{\beta(\wp_6^*)} \int_0^t (t-s)^{\wp_6^*-1} \mathcal{Y}_6\left(s, \dot{R}(s)\right) ds \end{split}$$

such that

$$S(t) - \dot{S}(t) \Big| \le \delta_1 \gamma_1,$$

$$\begin{split} \left| S_p(t) - \dot{S}_p(t) \right| &\leq \delta_2 \gamma_2, \\ \left| I_A(t) - \dot{I}_A(t) \right| &\leq \delta_3 \gamma_3, \\ \left| I_s(t) - \dot{I}_s(t) \right| &\leq \delta_4 \gamma_4, \\ \left| Q(t) - \dot{Q}(t) \right| &\leq \delta_5 \gamma_5, \\ \left| R(t) - \dot{R}(t) \right| &\leq \delta_6 \gamma_6. \end{split}$$

Theorem 4. If Assumption (B) is satisfied, then (1) is Hyers – Ulam-stable.

Proof. By Theorem 3, the Covid-19 model (1) has a unique solution, say, S(t), $S_p(t)$, $I_A(t)$, $I_s(t)$, Q(t), R(t). Let $(\dot{S}(t), \dot{S}_p(t), \dot{I}_A(t), \dot{Q}(t), \dot{R}(t))$ be an approximate solution of (1) satisfying (8)–(13). Then, we have

$$\begin{split} \left\| S(t) - \dot{S}(t) \right\| &\leq \frac{1 - \wp_1^*}{\beta(\wp_1^*)} \left\| Y_1(t, S(t)) - Y_1\left(t, \dot{S}(t)\right) \right\| + \\ &+ \frac{\wp_1^*}{\beta(\wp_1^*) \Gamma(\wp_1^*)} \int_0^t (t - s)^{\wp_1^* - 1} \left\| Y_1(s, S(s)) - Y_1\left(t, \dot{S}(t)\right) \right\| ds \leq \\ &\leq \left[\frac{1 - \wp_1^*}{\beta(\wp_1^*)} + \frac{1}{\beta(\wp_1^*) \Gamma(\wp_1^*)} \right] \psi_1 \| S - \dot{S} \|. \end{split}$$

Taking $\gamma_1 = \psi_1$, $\Delta = \frac{1 - \wp_1^*}{\beta(\wp_1^*)} + \frac{\wp_1^*}{\beta(\wp_1^*)\Gamma(\wp_1^*)}$, this implies $\left\| S(t) - \dot{S}(t) \right\| \le \gamma_1 \Delta_1.$

Similarly, for $S_p(t)$, $\dot{S}_p(t)$, $I_A(t)$, $\dot{I}_A(t)$, $I_s(t)$, $\dot{I}_s(t)$, Q(t), $\dot{Q}(t)$, R(t), $\dot{R}(t)$ we have

$$\begin{split} \left\| S_p(t) - \dot{S}_p(t) \right\| &\leq \gamma_2 \Delta_2, \\ \left\| I_A(t) - \dot{I}_A(t) \right\| &\leq \gamma_3 \Delta_3, \\ \left\| I_s(t) - \dot{I}_s(t) \right\| &\leq \gamma_4 \Delta_4, \\ \left\| Q(t) - \dot{Q}(t) \right\| &\leq \gamma_5 \Delta_5, \\ \left\| R(t) - \dot{R}(t) \right\| &\leq \gamma_6 \Delta_6. \end{split}$$

This implies, the system (1) is Hyers – Ulam stable which ultimately, ensures the stability of (1). This completes the proof.

5. Numerical scheme. With the help of (2)-(7), we produce the following numerical scheme:

$$\begin{cases} {}^{ABC}_{0} \mathcal{D}^{\varphi_{1}^{*}}_{t} S(t) = Y_{1}(t, S), \\ {}^{ABC}_{0} \mathcal{D}^{\varphi_{2}^{*}}_{t} S_{p}(t) = Y_{2}(t, S_{p}), \\ {}^{ABC}_{0} \mathcal{D}^{\varphi_{3}^{*}}_{t} I_{A}(t) = Y_{3}(t, I_{A}), \\ {}^{ABC}_{0} \mathcal{D}^{\varphi_{4}^{*}}_{t} I_{s}(t) = Y_{4}(t, I_{s}), \\ {}^{ABC}_{0} \mathcal{D}^{\varphi_{5}^{*}}_{t} Q(t) = Y_{5}(t, Q), \\ {}^{ABC}_{0} \mathcal{D}^{\varphi_{5}^{*}}_{t} R(t) = Y_{6}(t, R). \end{cases}$$
(20)

With the help of fractional AB-integral operator, (20) takes the following form:

$$\begin{split} S(t) - S(0) &= \frac{1 - \varphi_1^*}{\beta(\varphi_1^*)} \,\mathcal{Y}_1(t, S) + \frac{\varphi_1^*}{\beta(\varphi_1^*)} \int_0^t (t-s)^{\varphi_1^*-1} \mathcal{Y}_1(s, S) ds, \\ S_p(t) - S_p(0) &= \frac{1 - \varphi_2^*}{\beta(\varphi_2^*)} \,\mathcal{Y}_2(t, I_s) + \frac{\varphi_2^*}{\beta(\varphi_2^*)} \int_0^t (t-s)^{\varphi_2^*-1} \mathcal{Y}_2(s, S_p) ds, \\ I_A(t) - I_A(0) &= \frac{1 - \varphi_3^*}{\beta(\varphi_3^*)} \,\mathcal{Y}_2(t, I_A) + \frac{\varphi_3^*}{\beta(\varphi_3^*)} \int_0^t (t-s)^{\varphi_3^*-1} \mathcal{Y}_3(s, I_A) ds, \\ I_s(t) - I_s(0) &= \frac{1 - \varphi_4^*}{\beta(\varphi_4^*)} \,\mathcal{Y}_4(t, I_s) + \frac{\varphi_4^*}{\beta(\varphi_3^*)} \int_0^t (t-s)^{\varphi_4^*-1} \mathcal{Y}_4(s, I_s) ds, \\ Q(t) - Q(0) &= \frac{1 - \varphi_5^*}{\beta(\varphi_5^*)} \,\mathcal{Y}_5(t, Q) + \frac{\varphi_5^*}{\beta(\varphi_5^*)} \int_0^t (t-s)^{\varphi_5^*-1} \mathcal{Y}_5(s, Q) ds, \\ R(t) - R(0) &= \frac{1 - \varphi_6^*}{\beta(\varphi_6^*)} \,\mathcal{Y}_6(t, R) + \frac{\varphi_6^*}{\beta(\varphi_6^*)} \Gamma(\varphi_6^*)} \int_0^t (t-s)^{\varphi_6^*-1} \mathcal{Y}_6(s, R) ds. \end{split}$$

By dividing the assumed interval [0, t] into subintervals by the help of point t_{m+1} , for m = 0, 1, 2..., we obtain

$$S(t_{m+1}) - S(0) = \frac{1 - \wp_1^*}{\beta(\wp_1^*)} \mathcal{Y}_1(t_m, S) + \frac{\wp_1^*}{\beta(\wp_1^*)} \sum_{k=0}^n \int_{t_k}^{t_{k+1}} (t_{m+1} - s)^{\wp_1^* - 1} \mathcal{Y}_1(s, S) ds,$$

$$S_p(t_{m+1}) - S_p(0) = \frac{1 - \wp_2^*}{\beta(\wp_2^*)} \mathcal{Y}_2(t_m, S_p) + \frac{\wp_2^*}{\beta(\wp_2^*)} \sum_{k=0}^n \int_{t_k}^{t_{k+1}} (t_{m+1} - s)^{\wp_2^* - 1} \mathcal{Y}_2(s, S_p) ds,$$

$$I_A(t_{m+1}) - I_c(0) = \frac{1 - \wp_3^*}{\beta(\wp_3^*)} \mathcal{Y}_3(t_m, I_A) + \frac{\wp_3^*}{\beta(\wp_3^*)} \sum_{k=0}^n \int_{t_k}^{t_{k+1}} (t_{m+1} - s)^{\wp_3^* - 1} \mathcal{Y}_3(s, I_A) ds,$$

$$I_s(t_{m+1}) - I_s(0) = \frac{1 - \wp_4^*}{\beta(\wp_4^*)} \mathcal{Y}_4(t_m, I_s) + \frac{\wp_4^*}{\beta(\wp_4^*)\Gamma(\wp_4^*)} \sum_{k=0}^n \int_{t_k}^{t_{k+1}} (t_{m+1} - s)^{\wp_4^* - 1} \mathcal{Y}_4(s, I_s) ds,$$

$$Q(t_{m+1}) - Q(0) = \frac{1 - \wp_5^*}{\beta(\wp_5^*)} \mathcal{Y}_5(t_m, Q) + \frac{\wp_5^*}{\beta(\wp_5^*)\Gamma(\wp_5^*)} \sum_{k=0}^n \int_{t_k}^{t_{k+1}} (t_{m+1} - s)^{\wp_5^* - 1} \mathcal{Y}_5(s, Q) ds,$$

$$R(t_{m+1}) - R(0) = \frac{1 - \wp_6^*}{\beta(\wp_6^*)} \mathcal{Y}_6(t_m, R) + \frac{\wp_6^*}{\beta(\wp_6^*) \Gamma(\wp_6^*)} \sum_{k=0}^n \int_{t_k}^{t_{k+1}} (t_{m+1} - s)^{\wp_6^* - 1} \mathcal{Y}_6(s, R) ds.$$
(21)

Now, by using Lagrange's interpolation, we get

$$\begin{split} S(t_{m+1}) &= S(0) + \frac{1 - \wp_1^*}{\beta(\wp_1^*)} \mathcal{Y}_1(t_k, S) + \frac{\wp_1^*}{\mathcal{B}(\wp_1^*)} \sum_{k=0}^n \left[\frac{h^{\wp_1^*} \mathcal{Y}_1(t_k, S)}{\Gamma(\wp_1^* + 2)} \times \right. \\ & \times \left((m+1-k)^{\wp_1^*} (m-k+2+\wp_1^*) - (m-k)^{\wp_1^*} (m-k+2+2\wp_1^*) \right) - \right. \\ & - \frac{h^{\wp_1^*} \mathcal{Y}_1(t_{k-1}, S)}{\Gamma(\wp_1^* + 2)} \left((m+1-k)^{\wp_1^*} - (m-k)^{\wp_1^*} (m+1-k+\wp_1^*) \right) \right], \\ S_p(t_{m+1}) &= S_p(0) + \frac{1 - \wp_2^*}{\beta(\wp_2^*)} \mathcal{Y}_2(t_k, S_p) + \frac{\wp_2^*}{\mathcal{B}(\wp_2^*)} \sum_{k=0}^n \left[\frac{h^{\wp_2^*} \mathcal{Y}_2(t_k, S_p)}{\Gamma(\wp_2^* + 2)} \times \right. \\ & \times \left((m+1-k)^{\wp_2^*} (m-k+2+\wp_2^*) - (m-k)^{\wp_2^*} (m-k+2+2\wp_2^*) \right) - \right. \\ & - \frac{h^{\wp_2^*} \mathcal{Y}_2(t_{k-1}, S_p)}{\Gamma(\wp_2^* + 2)} \left((m+1-k)^{\wp_2^*} - (m-k)^{\wp_2^*} (m+1-k+\wp_2^*) \right) \right], \\ I_A(t_{m+1}) &= I_A(0) + \frac{1 - \wp_3^*}{\beta(\wp_3^*)} \mathcal{Y}_3(t_k, I_A) + \frac{\wp_3^*}{\mathcal{B}(\wp_3^*)} \sum_{k=0}^n \left[\frac{h^{\wp_3^*} \mathcal{Y}_3(t_k, I_A)}{\Gamma(\wp_3^* + 2)} \times \right. \\ & \times \left((m+1-k)^{\wp_3^*} (m-k+2+\wp_3^*) - (m-k)^{\wp_3^*} (m-k+2+2\wp_3^*) \right) - \right. \\ & - \frac{h^{\wp_3^*} \mathcal{Y}_3(t_{k-1}, I_A)}{\Gamma(\wp_3^* + 2)} \left((m+1-k)^{\wp_3^*} - (m-k)^{\wp_3^*} (m-k+2+2\wp_3^*) \right) \right], \\ I_s(t_{m+1}) &= I_s(0) + \frac{1 - \wp_4^*}{\beta(\wp_4^*)} \mathcal{Y}_4(t_k, I_s) + \frac{\wp_4^*}{\mathcal{B}(\wp_4^*)} \sum_{k=0}^n \left[\frac{h^{\wp_4^*} \mathcal{Y}_4(t_k, I_s)}{\Gamma(\wp_4^* + 2)} \times \right] \end{split}$$

$$\times \left((m+1-k)^{\wp_4^*} (m-k+2+\wp_4^*) (m-k)^{\wp_4^*} (m-k+2+2\wp_4^*) \right) - \\ - \frac{h^{\wp_4^*} \mathcal{Y}_4(t_{k-1}, I_s)}{\Gamma(\wp_4^*+2)} \left((m+1-k)^{\wp_4^*} - (m-k)^{\wp_4^*} (m+1-k+\wp_4^*) \right) \right], \\ Q(t_{m+1}) = Q(0) + \frac{1-\wp_5^*}{\beta(\wp_5^*)} \mathcal{Y}_5(t_k, Q) + \frac{\wp_5^*}{\mathcal{B}(\wp_5^*)} \sum_{k=0}^n \left[\frac{h^{\wp_5^*} \mathcal{Y}_5(t_k, Q)}{\Gamma(\wp_5^*+2)} \times \\ \times \left((m+1-k)^{\wp_5^*} (m-k+2+\wp_5^*) (m-k)^{\wp_5^*} (m-k+2+2\wp_5^*) \right) - \\ - \frac{h^{\wp_5^*} \mathcal{Y}_5(t_{k-1}, Q)}{\Gamma(\wp_5^*+2)} \left((m+1-k)^{\wp_5^*} - (m-k)^{\wp_5^*} (m+1-k+\wp_5^*) \right) \right], \\ R(t_{m+1}) = R(0) + \frac{1-\varphi_6^*}{\beta(\wp_6^*)} \mathcal{Y}_6(t_k, R) + \frac{\varphi_6^*}{\mathcal{B}(\wp_6^*)} \sum_{k=0}^n \left[\frac{h^{\wp_6^*} \mathcal{Y}_6(t_k, R)}{\Gamma(\wp_6^*+2)} \times \\ \times \left((m+1-k)^{\wp_6^*} (m-k+2+\wp_6^*) (m-k)^{\wp_6^*} (m-k+2+2\wp_6^*) \right) - \\ - \frac{h^{\wp_6^*} \mathcal{Y}_6(t_{k-1}, R)}{\Gamma(\wp_6^*+2)} \left((m+1-k)^{\wp_6^*} - (m-k)^{\wp_6^*} (m+1-k+\varphi_6^*) \right) \right].$$

This numerical scheme will help us to predict the role of protected susceptible which has been practically exercised in various nations as a control strategy. Although, this strategy has a worst effect on the economy of a nation but it is essential to curtail the spread of the infection of lethal Covid-19. The sensitivity analysis has been given in [36] which shows that the role of such a strategy is very much effective in the curtail of the spread.

5.1. Numerical results. In this section, we are providing a detail of numerical results related to the model with the available data in literature. The parameters and initial data was carried out from the available literature. The initial values are: S(0) = 59300000, $S_P(0) = 0$, $I_S(0) = 0$, Q(0) = 0, $I_A(0) = 2079$, R(0) = 903, and the parametric values are $\alpha_1 = 0.0008$, $\alpha_2 = 0.1$, $eta_1 = 0.25$, $\rho = 0.0001$, $\eta_2 = 0$, $\eta_3 = 0.385$, $r_1 = 0.2976$, $r_2 = 0$, $\gamma = 0$, $r_3 = 0.2976$, $\mu = 0.00236/90$, $\delta = 0.017/90$, $\alpha_3 = 1$, $\Lambda_1 = 296425.875/90$ [36].

In Fig. 1, we have a joint comparative simulation for the two classes S(t) and $S_P(t)$ for the orders 1.0, 0.99, 0.98, 0.97. The second figure that is Fig. 2, represents a graphical study of the S(t) class for various orders 1.0, 0.99, 0.98, 0.97 for a long time of 5000 days. There is a decrease in the population of the class. Also, as much the order is decreasing a comparative large decrease is observed in the population while the behavior of the class remains similar. Figure 3 shows a comparative analysis of the $S_P(t)$ for the mentioned orders and a gradual increase can be seen in the graph.

The Fig. 4 is for the infection population which shows an increase up to the 300 days while a decrease is observed after 300 to 600. Figure 5 shows a numerical representation of the $I_S(t)$ class for the various orders and the Fig. 6 is for the R(t) class.



Fig. 1. Comparative analysis for the S(t) and $S_P(t)$ for the orders 1.0, 0.99, 0.98, 0.97.



Fig. 2. Comparative analysis for the S(t) for the orders 1.0, 0.99, 0.98, 0.97.

ISSN 1562-3076. Нелінійні коливання, 2021, т. 24, № 3







Fig. 4. Comparative analysis for the I(t) for the orders 1.0, 0.99, 0.98, 0.97.



Fig. 6. Comparative analysis for the R(t) for the orders 1.0, 0.99, 0.98, 0.97.

ISSN 1562-3076. Нелінійні коливання, 2021, т. 24, № 3

6. Conclusion. In this article, we have focused on the theoretical as well computational studies of the fractional order Covid-19 model in the ABC-sense of derivative. The existence, uniqueness results were carried out with the help of iterative sequential approach with the limiting point as the solution of the suggested model (1). We also estimated the Hyers – Ulam stability and a numerical scheme based on the Lagrange's interpolation was obtained. The numerical scheme was then tested and very similar results like the integer order was obtained. The numerical results were interpreted via six graphs. The details are: in Fig. 1, we have a joint comparative simulation for the two classes S(t) and $S_P(t)$ for the orders 1.0, 0.99, 0.98, 0.97. The second figure that is Fig. 2, represents a graphical study of the S(t) class for various orders 1.0, 0.99, 0.98, 0.97 for a long time of 5000 days. There is a decrease in the population of the class. Also, as much the order is decreasing a comparative large decrease is observed in the population while the behavior of the class remains similar. Figure 3, shows a comparative analysis of the $S_P(t)$ for the mentioned orders and a gradual increase can be seen in the graph. The Fig. 4 is for the infection population which shows an increase upto the 300 days while a decrease is observed after 300 to 600. Figure 5 shows a numerical representation of the $I_S(t)$ class for the various orders and the Fig. 6 is for the R(t) class. The reader of the paper can work on the comparative analysis of different fractional operators for more accuracy and better results.

Acknowledgement. The authors A. Khan and T. Abdeljawad would like to thank Prince Sultan University for the support through the TAS research lab. Also, the authors would like to thank prof. T. Abdeljawad for his support to improve this paper.

References

- 1. A. E. Gorbalenya, S. C. Baker, R. S. Baric, R. J. de Groot, C. Drosten, A. A. Gulyaeva, B. L. Haagmans, C. Lauber, A. M. Leontovich, B. W. Neuman, D. Penzar, *Coronaviridae study group of the international committee on taxonomy of viruses. The species severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2*, Nat. Microbiol., **5**, № 4, 536–544 (2020).
- S. A. Morrison, J. Gregor, S. Gregor, Responding to a global pandemic: Republic of Slovenia on maintaining physical activity during self-isolation, Scand. J. Med. Sci. Sports, 30, № 8 (2020).
- 3. N. Kokudo, H. Sugiyama, *Call for international cooperation and collaboration to effectively tackle the Covid-19 pandemic*, Global Health Med., **30**, 2(2), 2–60 (2020).
- S. Omer, S. Ali, Preventive measures and management of Covid-19 in pregnancy, Drugs Therapy Perspect., 36, № 6 (2020).
- 5. K. F. Owusu, E. F. Goufo, S. Mugisha, *Modelling intracellular delay and therapy interruptions within Ghanaian HIV population*, Adv. Difference Equ., **1**, 1–9 (2020).
- 6. M. Anderson, M. Mckee, E. Mossialos, *Developing a sustainable exit strategy for Covid-19: health, economic and public policy implications*, J. R. Soc. Med., **113(5)**, 8–176 (2020).
- 7. E. D. Goufo, R. Maritz, A note on ebolas outbreak and human migration dynamic, J. Human Ecology, **51**, № 3, 257–263 (2015).
- 8. P. T. Djomegni, A.Tekle, M. Y. Dawed, *Pre-exposure prophylaxis HIV/AIDS mathematical model with non classical isolation*, Jpn. J. Ind. Appl. Math., **37**, 781–801 (2020).
- 9. K. F. Owusu, E. F. Goufo, S. Mugisha, *Modelling intracellular delay and therapy interruptions within Ghanaian HIV population*, Adv. Difference Equ., **1**, 1–9 (2020).
- 10. Z. A. Khan, F. Jarad, A. Khan, H. Khan, *Nonlinear discrete fractional sum inequalities related to the theory of discrete fractional calculus with applications*, Adv. Difference Equ., **1**, 1–3 (2021).

- 11. A. Khan, H. M. Alshehri, T. Abdeljawad, Q. M. Al-Madlal, H. Khan, *Stability analysis of fractional nabla difference Covid-19 model*, Results Phys., **4**, 103–888 (2021).
- 12. A. Shah, R. A. Khan, A. Khan, H. Khan, J. F. Gomez-Aguilar, *Investigation of a system of nonlinear fractional* order hybrid differential equations under usual boundary conditions for existence of solution, Math. Methods Appl. Sci., **30**, № 2, 1628–38 (2021).
- 13. T. Abdeljawad, A Lyapunov type inequality for fractional operators with non singular Mittag-Leffler kernel, J. Inequal Appl., **130**, (2017); DOI: 10.1186/s13660-017-1400-5.
- 14. T. Abdeljawad, *Fractional operators with generalized Mittag-Leffler kernels and their iterated differintegrals*, Chaos, **29**, № 2, 023102 (2019).
- 15. T. Abdeljawad, D. Baleanu, *Discrete fractional differences with nonsingular discrete Mittag-Leffler kernels*, Adv. Difference Equ., Paper No. 232 (2016).
- 16. T. Abdeljawad, Q. M. Al-Mdallal, F. Jarad, *Fractional logistic models in the frame of fractional operators generated by conformable derivatives*, Chaos Solitons Fractals, **119**, 94–101 (2019).
- 17. T. Abdeljawad, M. A. Hajji, Q. M. Al-Mdallal, F. Jarad, *Analysis of some generalized ABC-fractional logistic* models, Alexandria Eng. J., **59**, № 4, 8–2141 (2020).
- B. Acay, E. Bas, T. Abdeljawad, Fractional economic models based on market equilibrium in the frame of different type kernels, Chaos Solitons Fractals, 130, 109438 (2020).
- 19. A. Atangana, S. I. Araz, *Mathematical model of Covid-19 spread in Turkey and South Africa: theory, methods and applications*, Adv. Difference Equ., Paper No. 659 (2020).
- 20. A. Atangana, S. I. Araz, Nonlinear equations with global differential and integral operators: existence, uniqueness with application to epidemiology, Results Phys., **20**, 103593 (2021).
- 21. A. Atangana, D. Baleanu, New fractional derivatives with nonlocal and non-singular kernel: theory and application to heat transfer model, Thermal Sci., 20(00), (2016); DOI: 10.2298/TSCI160111018A.
- 22. I. V. Atamas', V. I. Slinko, *Stability of the fixed points of a class of quasilinear cascades in the space* conv \mathbb{R}^n , (Russian. English, Ukrainian summary) Ukr. Mat. Zh., **69**, Nº 8, 1166–1179 (2017).
- 23. M. Arfan, K. Shah, T. Abdeljawad, N. Mlaiki, A. Ullah, *A Caputo power law model predicting the spread of the Covid-19 outbreak in Pakistan*, Alexandria Eng. J., **1**, 60(1), 56–447 (2021).
- 24. R. Begum, O. Tunç, H. Khan, H. Gulzar, A. Khan, A fractional order Zika virus model with Mittag-Leffler kernel, Chaos Solitons Fractals, 146, № 110 898 (2021).
- 25. M. Bohner, O. Tunç, C. Tunç, *Qualitative analysis of Caputo fractional integro-differential equations with constant delays*, Comput. Appl. Math., 40, № 6, Paper № 214 (2021).
- A. I. Dvirnyi, V. I. Slyn'ko, Stability of solutions of pseudolinear differential equations with impulse action, Math. Notes, 93, № 5-6, 691-703 (2013).
- 27. J. F. Gomez, L. Torres, R. F. Escobar, *Fractional derivatives with Mittag-Leffler kernel. Trends and applications in science and engineering*, Studies in Systems, Decision and Control, **194**, Springer, Cham (2019).
- 28. F. Jarad, T. Abdeljawad, Z. Hammouch, On a class of ordinary differential equations in the frame of Atangana-Baleanu fractional derivative, Chaos Solitons Fractals, 1, № 117, 16–20 (2018).
- 29. K. M. Owolabi, A. Atangana, On the formulation of Adams-Bashforth scheme with Atangana-Baleanu-Caputo fractional derivative to model chaotic problems, Chaos, 29, No 2, 023111 (2019).
- 30. K. Shah, M. A. Alqudah, F. Jarad, T. Abdeljawad, *Semi-analytical study of pine wilt disease model with convex rate under Caputo Febrizio fractional order derivative*, Chaos Solitons Fractals, **135**, 109754 (2020).
- 31. O. Tunç, Ö. Atan, C. Tunç, J. C. Yao, *Qualitative analyses of integro-fractional differential equations with Caputo derivatives and retardations via the Lyapunov–Razumikhin method*, Axioms, **10**, № 2 (2021); DOI: https://doi.org/10.3390/axioms10020058.

- 32. E. Tunç, O. Tunç, *On the oscillation of a class of damped fractional differential equations*, Miskolc Math. Notes, **17**, № 1, 647–656 (2016).
- 33. M. Yavuz, T. Abdeljawad, *Nonlinear regularized long-wave models with a new integral transformation applied* to the fractional derivative with power and Mittag-Leffler kernel, Adv. Difference Equ., Paper No. 367 (2020).
- 34. F. B. Yousef, A. Yousef, T. Abdeljawad, A. Kalinli, *Mathematical modeling of breast cancer in a mixed immune-chemotherapy treatment considering the effect of ketogenic diet*, Eur. Phys. J. Plus., **135**, № 12, 1–23 (2020).
- 35. M. A. Khan, A. Atangana, E. Alzahrani, *The dynamics of Covid-19 with quarantined and isolation*, Adv. Difference Equ., **2020**, № 1, 1–22 (2020).
- 36. P. M. Djomegni, M. D. Haggar, W. T. Adigo, *Mathematical model for Covid-19 with "protected susceptible" in the post-lockdown era*, Alexandria Eng. J., **1**, 60(1), 35–527 (2021).

Received 07.09.21, *after revision* — 24.11.21