## Density of periodic orbit measures for piecewise monotonic interval maps

## Peter Raith

Let  $T : [0,1] \to [0,1]$  be a piecewise monotonic map, this means there exists a partition  $\mathcal{Z}$  of [0,1] into finitely many pairwise disjoint open intervals with  $\bigcup_{Z \in \mathcal{Z}} \overline{Z} = [0,1]$ , such that for every  $Z \in \mathcal{Z}$  the map  $T|_Z$  is continuous and strictly monotonic. (Note that T need not be continuous at the endpoints of the intervals in  $\mathcal{Z}$ .) Denote the set of all T-invariant Borel probability measures on [0,1] by  $\mathcal{M}(T)$ , and let  $\mathcal{P}(T)$  be the set of all T-invariant Borel probability measures concentrated on a periodic orbit of T. The set  $\mathcal{M}(T)$  is endowed with the weak star-topology.

**Problem.** Suppose that T is topologically transitive and  $h_{top}(T) > 0$ . Is  $\mathcal{P}(T)$  dense in  $\mathcal{M}(T)$ ?

It is known that the above problem has a positive answer in the following cases:

- The map T is continuous (Alexander Blokh, see e.g. [1]). (*Remark*: In this case T need not be piecewise monotonic.)
- The map T is a monotonic mod one transformation (also called Lorenz map), this means there exists a strictly increasing and continuous function  $f : [0,1] \to \mathbb{R}$ , such that  $Tx = f(x) \pmod{1}$  (Franz Hofbauer in [2]).
- The map T has two intervals of monotonicity, this means card  $\mathcal{Z} = 2$ , where  $\mathcal{Z}$  is as above ([3]).

## References

- A. Blokh, The 'spectral' decomposition for one-dimensional maps, Dynamics Reported, vol. 4 (C. K. R. T. Jones, V. Kirchgraber, H. O. Walther, eds.), Springer, Berlin, 1995, pp. 1–59.
- F. Hofbauer, Generic properties of invariant measures for simple piecewise monotonic transformations, Israel J. Math. 59 (1987), 64–80.
- [3] F. Hofbauer, P. Raith, Density of periodic orbit measures for transformations on the interval with two monotonic pieces, Fund. Math. 157 (1998), 221–234.