

Recursion Operators for Vacuum Einstein Equations with Symmetries

Michal MARVAN

Mathematical Institute, Silesian University in Opava, Na Rybníčku 1, Czech Republic

E-mail: Michal.Marvan@math.slu.cz

Direct and inverse recursion operator is derived for the vacuum Einstein equations for metrics with two commuting Killing vectors that are orthogonal to a foliation by 2-dimensional leaves.

1 Introduction

In the past decade, inverse recursion operators became the subject of a number of papers [7, 8, 14, 15]. In particular, the work of Guthrie [7] opened a new perspective on recursion operators by essentially identifying them with auto-Bäcklund transformations for linearized equation [18].

It is known for a long time that recursion operators of integrable systems are obtainable from their Lax pairs (see [6] and references therein) and ZCR's (see [10, 24]). However, recently it became clear that zero curvature representations are related much closer to inverse recursion operators than to their 'direct' counterparts [19, 20]. Examples that have been already published elsewhere include the Korteweg–de Vries and Tzitzéica equation [19] and the stationary Nizhnik–Novikov–Veselov equation [20]. In the present paper, the methods of [19, 20] are applied to equations of General Relativity.

2 Recursion operators

Let $\mathcal{E} = \{F^l = 0\}$ be a system of PDE's in unknown functions u^k of two independent variables x, y . We assume that F^l are functions of x, y, u^k and a finite number of the derivatives

$$u_{ij}^k = \frac{\partial^{i+j} u^k}{\partial x^i \partial y^j} \quad (u_{00}^k = u^k).$$

Consider the infinite-dimensional jet space J^∞ with local coordinates x, y, u_{ij}^k along with the commuting vector fields

$$D_x = \frac{\partial}{\partial x} + \sum_{ij} u_{i+1,j}^k \frac{\partial}{\partial u_{ij}^k}, \quad D_y = \frac{\partial}{\partial y} + \sum_{ij} u_{i,j+1}^k \frac{\partial}{\partial u_{ij}^k},$$

called *total derivatives*. The submanifold E determined by equations $F^l = 0$ and their differential consequences $D_x F^l = 0, D_y F^l = 0, D_x^2 F^l = 0, D_x D_y F^l = 0, D_y^2 F^l = 0, \dots$, is called the *equation manifold* (and is an underlying space of the diffeity structure [12] employed in [19]). In this context, infinitesimal symmetries (more precisely, their generating functions) are functions U^k defined on E such that

$$\sum_{k,i,j} \frac{\partial F^l}{\partial u_{ij}^k} D_x^i D_y^j U^k = 0.$$

It is then natural to consider the jet space with coordinates x, y, u_{ij}^k, U_{ij}^k , and denote

$$LF^l = \sum_{k,i,j} \frac{\partial F^l}{\partial u_{ij}^k} U_{ij}^k. \quad (1)$$

The system $L\mathcal{E} := \{F^l = 0, LF^l = 0\}$ on unknowns u^k, U^k will be called the *linearized equation*. Now, Guthrie's recursion operators [7] may be interpreted as auto-Bäcklund transformations of the linearized equation $L\mathcal{E}$ that keep variables u^k unchanged [19].

In now standard formalism [21], recursion operators are pseudodifferential operators, characterized by the occurrence of inverses of total derivatives D_x^{-1} . Under Guthrie's approach, $p = D_x^{-1}f$ is introduced as an auxiliary nonlocal variable satisfying

$$p_x = f, \quad p_y = g, \quad (2)$$

provided such a g exists, and it actually does without known exception; see Sergyeyev [23] for a proof in the case of evolution systems. Thus, p is a potential of a conservation law $f dx + g dy$ of the linearized equation $L\mathcal{E}$.

For the inverse recursion operators, the nonlocalities tend to be genuinely non-Abelian pseudopotentials related to a zero curvature representation of the system in question. Let \mathfrak{g} be a matrix Lie algebra. Let $\alpha = A dx + B dy$ be a \mathfrak{g} -valued zero curvature representation (ZCR) for the system \mathcal{E} . This means that A, B are \mathfrak{g} -valued functions on the equation submanifold E and $D_y A - D_x B + [A, B] = 0$ holds on E . Let us introduce the associated pseudopotential P as a \mathfrak{g} -valued solution of the compatible system

$$P_x = [A, P] + LA, \quad P_y = [B, P] + LB. \quad (3)$$

A recursion operator R is then a linear operator in U^k and P such that $U' = R(U, P)$ solves the linearized system $L\mathcal{E}$ whenever U does and P satisfies (3) (see [19,20]). In this way, the inverse recursion operator can be found without previous knowledge of the direct recursion operator. A remarkable aspect of this approach is that $R(U, P)$ tends to be a very simple expression.

For the above scheme to work, it is not necessary that the ZCR α a priori depends on the "spectral parameter". However, if R is a recursion operator related to the ZCR α as above, then $(R^{-1} + \mu \text{Id})^{-1}$ is another recursion operator, associated with a ZCR α_μ which depends on μ .

3 The results

We consider vacuum Einstein equations for a space-time with two commuting Killing vectors that are orthogonal to a foliation by 2-dimensional surfaces [4,5]. Our presentation will be restricted to the case when both Killing vectors are space-like. The case when one of the Killing vectors is time-like is equivalent to ours via an appropriate complex transformation of coordinates.

As is well known, there exist coordinates x, y, z^1, z^2 such that the metric in question can be written in the form $ds^2 = 2f(x, y) dx dy + g_{ij}(x, y) dz^i dz^j$ (the Lewis [13] metric). The vacuum Einstein equations essentially reduce to

$$(\sqrt{\det g} g_x g^{-1})_y + (\sqrt{\det g} g_y g^{-1})_x = 0, \quad (4)$$

while f can be obtained by quadrature. Using the standard normalization $\det g = (x + y)^2$ compatible with equation (4), we parametrize g as follows: $g_{11} = (x + y)/u$, $g_{12} = (x + y)v/u$, $g_{22} = (x + y)(u^2 + v^2)/u$. Equation (4) then becomes

$$u_{xy} = \frac{u_x u_y - v_x v_y}{u} - \frac{1}{2} \frac{u_x + u_y}{x + y}, \quad v_{xy} = \frac{v_x u_y + u_x v_y}{u} - \frac{1}{2} \frac{v_x + v_y}{x + y}. \quad (5)$$

As is well known, equation (5) has a ZCR and a Bäcklund transformation [1–3, 9, 16, 17, 22]. The ZCR reads

$$A = \frac{1}{2} \begin{pmatrix} -(\theta+1)u_x/u & (\theta+1)v_x/u^2 \\ (\theta-1)v_x & (\theta+1)u_x/u \end{pmatrix}, \quad B = \frac{1}{2\theta} \begin{pmatrix} -(\theta+1)u_y/u & (\theta+1)v_y/u^2 \\ (-\theta+1)v_y & (\theta+1)u_y/u \end{pmatrix},$$

where $\theta = \sqrt{(\mu+y)/(\mu-x)}$, μ being the spectral parameter.

The main result of this paper, obtained by the methods of [19, 20], is as follows: If nonlocal variables p_{11} , p_{12} , p_{21} satisfy

$$\begin{aligned} p_{11,x} &= -\frac{\theta-1}{2}v_x p_{12} + \frac{\theta+1}{2}\frac{v_x}{u^2}p_{21} - \frac{\theta+1}{2}\frac{1}{u}U_x + \frac{\theta+1}{2}\frac{u_x}{u^2}U, \\ p_{12,x} &= -(\theta+1)\frac{v_x}{u^2}p_{11} - (\theta+1)\frac{u_x}{u}p_{12} - (\theta+1)\frac{v_x}{u^3}U + \frac{\theta+1}{2}\frac{1}{u^2}V_x, \\ p_{21,x} &= (\theta-1)v_x p_{11} + (\theta+1)\frac{u_x}{u}p_{21} + \frac{\theta-1}{2}V_x, \\ p_{11,y} &= \frac{\theta-1}{2\theta}v_y p_{12} + \frac{\theta+1}{2\theta}\frac{v_y}{u^2}p_{21} + \frac{\theta+1}{2\theta}\frac{u_y}{u^2}U - \frac{\theta+1}{2\theta}\frac{1}{u}U_y, \\ p_{12,y} &= -\frac{\theta+1}{\theta}\frac{v_y}{u^2}p_{11} - \frac{\theta+1}{\theta}\frac{u_y}{u}p_{12} - \frac{\theta+1}{\theta}\frac{v_y}{u^3}U + \frac{\theta+1}{2\theta}\frac{1}{u^2}V_y, \\ p_{21,y} &= -\frac{\theta-1}{\theta}v_y p_{11} + \frac{\theta+1}{\theta}\frac{u_y}{u}p_{21} - \frac{\theta-1}{2\theta}V_y, \end{aligned} \quad (6)$$

then

$$\begin{aligned} U' &= 2\frac{u}{\sqrt{(\mu-x)(\mu+y)}}p_{11} + \frac{1}{\sqrt{(\mu-x)(\mu+y)}}U, \\ V' &= -\frac{u^2}{\sqrt{(\mu-x)(\mu+y)}}p_{12} - \frac{1}{\sqrt{(\mu-x)(\mu+y)}}p_{21} \end{aligned} \quad (7)$$

is a recursion operator for equation (5), namely, it sends symmetries to symmetries if the latter are viewed as solutions of the linearized system

$$\begin{aligned} U_{xy} &= \left(\frac{u_y}{u} - \frac{1}{2(x+y)}\right)U_x + \left(\frac{u_x}{u} - \frac{1}{2(x+y)}\right)U_y - \frac{u_x u_y - v_x v_y}{u^2}U - \frac{v_y}{u}V_x - \frac{v_x}{u}V_y, \\ V_{xy} &= \frac{v_y}{u}U_x + \frac{v_x}{u}U_y - \frac{v_x u_y + u_x v_y}{u^2}U + \left(\frac{u_y}{u} - \frac{1}{2(x+y)}\right)V_x + \left(\frac{u_x}{u} - \frac{1}{2(x+y)}\right)V_y. \end{aligned}$$

The ‘direct’ recursion operator for this equation seems to be missing in the literature; we can obtain it by inverting the operator (7), the result being

$$\begin{aligned} U' &= uvp_1 - up_2 + (y-x)U, \\ V' &= -\frac{1}{2}(u^2 - v^2)p_1 - vp_2 - \frac{1}{2}p_3 + (y-x)V, \end{aligned}$$

where p_1 , p_2 , p_3 satisfy

$$\begin{aligned} p_{1,x} &= (x+y) \left(-2\frac{v_x}{u^3}U + \frac{1}{u^2}V_x\right), \\ p_{2,x} &= (x+y) \left(-\frac{uu_x + 2vv_x}{u^3}U + \frac{1}{u}U_x + \frac{v_x}{u^2}V + \frac{v}{u^2}V_x\right), \\ p_{3,x} &= (x+y) \left(2\frac{(uu_x + vv_x)v}{u^3}U - 2\frac{v}{u}U_x - 2\frac{uu_x + vv_x}{u^2}V + \frac{u^2 - v^2}{u^2}V_x\right), \end{aligned}$$

$$\begin{aligned}
p_{1,y} &= (x+y) \left(2 \frac{v_y}{u^3} U - \frac{1}{u^2} V_y \right), \\
p_{2,y} &= (x+y) \left(\frac{uu_y + 2vv_y}{u^3} U - \frac{1}{u} U_y - \frac{v_y}{u^2} V - \frac{v}{u^2} V_y \right), \\
p_{3,y} &= (x+y) \left(-2 \frac{(uu_y + vv_y)v}{u^3} U + 2 \frac{v}{u} U_y + 2 \frac{uu_y + vv_y}{u^2} V - \frac{u^2 - v^2}{u^2} V_y \right).
\end{aligned}$$

It is readily seen that p_i are potentials of the linearizations [18] of the three obvious conservation laws of equation (4).

Quite unusually, neither of the recursion operators found generates an infinite series of local symmetries (and no such series is known). The action of our operators on the infinite-dimensional Geroch group of nonlocal symmetries [5, 11] remains to be investigated.

It is convenient to rewrite system (6) in triangular form. To achieve this, we introduce the Riccati pseudopotential q by

$$q_x = \frac{\theta - 1}{2} v_x q^2 - (\theta + 1) \frac{u_x}{u} q - \frac{\theta + 1}{2} \frac{v_x}{u^2}, \quad q_y = -\frac{\theta - 1}{2\theta} v_y q^2 - \frac{\theta + 1}{2\theta} \frac{u_y}{u} q - \frac{\theta + 1}{2\theta} \frac{v_y}{u^2}$$

and a nonlocal potential r by

$$r_x = (\theta - 1) v_x q - (\theta + 1) \frac{u_x}{u}, \quad r_y = -\frac{\theta - 1}{\theta} v_y q - \frac{\theta + 1}{\theta} \frac{u_y}{u}.$$

Then the inverse recursion operator assumes the form

$$\begin{aligned}
U' &= \frac{1}{\sqrt{(\mu - x)(\mu + y)}} \left(2uQ - 2 \frac{uq}{e^r} R + U \right), \\
V' &= \frac{1}{\sqrt{(\mu - x)(\mu + y)}} \left(-u^2 e^r P - 2u^2 qQ + \frac{u^2 q^2 - 1}{e^r} R \right),
\end{aligned}$$

where P, Q, R are supposed to satisfy

$$\begin{aligned}
P_x &= (\theta + 1) \frac{q}{u} e^{-r} U_x + \left(\frac{\theta + 1}{2} \frac{1}{u^2} - \frac{\theta - 1}{2} q^2 \right) e^{-r} V_x - (\theta + 1) \left(\frac{qu_x}{u^2} + \frac{v_x}{u^3} \right) e^{-r} U, \\
Q_x &= -\frac{\theta + 1}{2} \frac{1}{u} U_x + \frac{\theta - 1}{2} q V_x + \frac{\theta + 1}{2} \frac{u_x}{u^2} U - \frac{\theta - 1}{2} v_x e^r P, \\
R_x &= \frac{\theta - 1}{2} e^r V_x + (\theta - 1) v_x e^r Q, \\
P_y &= \frac{\theta + 1}{\theta} \frac{q}{u} e^{-r} U_y + \left(\frac{\theta + 1}{2\theta} \frac{1}{u^2} + \frac{\theta - 1}{2\theta} q^2 \right) e^{-r} V_y - \frac{\theta + 1}{\theta} \left(\frac{qu_y}{u^2} + \frac{v_y}{u^3} \right) e^{-r} U, \\
Q_y &= -\frac{\theta + 1}{2\theta} \frac{1}{u} U_y - \frac{\theta - 1}{2\theta} q V_y + \frac{\theta + 1}{2\theta} \frac{u_y}{u^2} U + \frac{\theta - 1}{2\theta} v_y e^r P, \\
R_y &= -\frac{\theta - 1}{2\theta} e^r V_y - \frac{\theta - 1}{\theta} v_y e^r Q.
\end{aligned}$$

This form of the inverse recursion operator is better adapted to generation of symmetries, which is, however, beyond the scope of this paper.

Acknowledgements

I would like to thank G. Vilasi for drawing my attention to the problem. The support from the grant MSM:J10/98:192400002 is gratefully acknowledged.

- [1] Belinskii V.A. and Zakharov V.E., Integration of the Einstein equations by means of the inverse scattering problem, *Soviet Phys. JETP*, 1978, V.75, N 6, 1955–1971.
- [2] Chinea F.J., Bäcklund transformations in general relativity, in Proc. Conf. “Nonlinear Phenomena” (1982, Oaxtepec, México), *Lecture Notes in Phys.*, Vol. 189, Berlin, Springer, 1983, 342–353.
- [3] Dodd R.K., Kinoulty J. and Morris H.C., Bäcklund transformation for the Ernst equation of general relativity, in *Advanced Nonlinear Waves, Res. Notes in Math.*, Vol. 95, Boston, Pitman, 1984, V.1, 254–272.
- [4] Geroch R., A method for generating solutions of Einstein’s equations, *J. Math. Phys.*, 1971, V.12, 918–924.
- [5] Geroch R., A method for generating new solutions of Einstein’s equations. II, *J. Math. Phys.*, 1972, V.13, 394–404.
- [6] Gürses M., Karasu A. and Sokolov V.V., On construction of recursion operators from Lax representation, *J. Math. Phys.*, 1999, V.40, 6473–6490.
- [7] Guthrie G.A., Recursion operators and non-local symmetries, *Proc. R. Soc. London A*, 1994, V.446, 107–114.
- [8] Guthrie G.A. and Hickman M.S., Nonlocal symmetries of the KdV equation, *J. Math. Phys.*, 1993, V.34, 193–205.
- [9] Harrison B.K., Bäcklund transformation for the Ernst equation of general relativity, *Phys. Rev. Lett.*, 1978, V.41, N 18, 1197–1200.
- [10] Karasu (Kalkanli) A., Karasu A. and Sakovich S.Yu., A strange recursion operator for a new integrable system of coupled Korteweg–de Vries equations, *Acta Appl. Math.*, to appear; nlin.SI/0203036.
- [11] Kinnersley W., Symmetries of the stationary Einstein–Maxwell field equations, *J. Math. Phys.*, 1977, V.18, 1529–1537.
- [12] Krasil’shchik I.S. and Vinogradov A.M. (Editors), Symmetries and conservation laws for differential equations of mathematical physics, *Translations of Mathematical Monographs*, Vol. 182, Providence, American Mathematical Society, 1999.
- [13] Lewis T., Some special solutions of the equations of axially symmetric gravitational fields, *Proc. Roy. Soc. London A*, 1932, V.136, 176–192.
- [14] Lou S., Recursion operator and symmetry structure of the Kawamoto-type equation, *Phys. Lett. A*, 1993, V.181, 13–16.
- [15] Lou S. and Chen W., Inverse recursion operator of the AKNS hierarchy, *Phys. Lett. A*, 1993, V.179, 271–274.
- [16] Maison D., Are the stationary, axially symmetric Einstein equations completely integrable?, *Phys. Rev. Lett.*, 1978, V.41, N 8, 521–522.
- [17] Maison D., On the complete integrability of the stationary, axially symmetric Einstein equations, *J. Math. Phys.*, 1979, V.20, N 5, 871–877.
- [18] Marvan M., Another look on recursion operators, in Proc. Conf. “Differential Geometry and Applications” (1995, Brno), Brno, Masaryk University, 1997, 393–402; www.emis.de/proceedings/6ICDGA/IV.
- [19] Marvan M., Reducibility of zero curvature representations with application to recursion operators, *Acta Appl. Math.*, to appear; nlin.SI/0306006.
- [20] Marvan M. and Sergyeyev A., Recursion operator for the Nizhnik–Veselov–Novikov equation, *J. Phys. A: Math. Gen.*, 2003, V.36, L87–L92; nlin.SI/0210028.
- [21] Olver P.J., Evolution equations possessing infinitely many symmetries, *J. Math. Phys.*, 1977, V.18, 1212–1215.
- [22] Omote M. and Wadati M., Bäcklund transformations for the Einstein equation, in *Advanced Nonlinear Waves, Res. Notes in Math.*, Vol. 95, Boston, Pitman, 1984, V.1, 242–253.
- [23] Sergyeyev A., On recursion operators and nonlocal symmetries of evolution equations, in *Proceedings of the Seminar on Differential Geometry (2000, Opava)*, Editor D. Krupka, Opava, Silesian University in Opava, 2000, 159–173; nlin.SI/0012011.
- [24] Sakovich S.Yu., Cyclic bases of zero-curvature representations: five illustrations to one concept, *Acta Appl. Math.*, to appear; nlin.SI/0212019.