

S. A. PLAKSA and V. S. SHPAKIVSKYI

On limiting values of Cauchy type integral in a harmonic algebra with two-dimensional radical

ABSTRACT. We consider a certain analog of Cauchy type integral taking values in a three-dimensional harmonic algebra with two-dimensional radical. We establish sufficient conditions for an existence of limiting values of this integral on the curve of integration.

1. Introduction. Let Γ be a closed Jordan rectifiable curve in the complex plane \mathbb{C} . By D^+ and D^- we denote, respectively, the interior and the exterior domains bounded by the curve Γ .

N. Davydov [1] established sufficient conditions for an existence of limiting values of the Cauchy type integral

$$(1) \quad \frac{1}{2\pi i} \int_{\Gamma} \frac{g(t)}{t - \xi} dt, \quad \xi \in \mathbb{C} \setminus \Gamma,$$

on Γ from the domains D^+ and D^- . This result stimulated a development of the theory of Cauchy type integral on curves which are not piecewise-smooth.

In particular, using the mentioned result of the paper [1], the following result was proved: if the curve Γ satisfies the condition (see [2])

$$(2) \quad \theta(\varepsilon) := \sup_{\xi \in \Gamma} \theta_{\xi}(\varepsilon) = O(\varepsilon), \quad \varepsilon \rightarrow 0$$

2010 *Mathematics Subject Classification.* 30G35, 30E25.

Key words and phrases. Three-dimensional harmonic algebra, Cauchy type integral, limiting values, closed Jordan rectifiable curve.

(here $\theta_\xi(\varepsilon) := \text{mes} \{t \in \Gamma : |t - \xi| \leq \varepsilon\}$, where mes denotes the linear Lebesgue measure on Γ), and the modulus of continuity

$$\omega_g(\varepsilon) := \sup_{t_1, t_2 \in \Gamma, |t_1 - t_2| \leq \varepsilon} |g(t_1) - g(t_2)|$$

of a function $g : \Gamma \rightarrow \mathbb{C}$ satisfies the Dini condition

$$(3) \quad \int_0^1 \frac{\omega_g(\eta)}{\eta} d\eta < \infty,$$

then the integral (1) has limiting values in every point of Γ from the domains D^+ and D^- (see [3]). The condition (2) means that the measure of a part of the curve Γ in every disk centered at a point of the curve is commensurable with the radius of the disk.

In this paper we consider a certain analogue of Cauchy type integral taking values in a three-dimensional harmonic algebra with two-dimensional radical and study the question about an existence of its limiting values on the curve of integration.

2. A three-dimensional harmonic algebra with a two-dimensional radical.

Let \mathbb{A}_3 be a three-dimensional commutative associative Banach algebra with unit 1 over the field of complex numbers \mathbb{C} . Let $\{1, \rho_1, \rho_2\}$ be a basis of algebra \mathbb{A}_3 with the multiplication table: $\rho_1\rho_2 = \rho_2^2 = 0, \rho_1^2 = \rho_2$.

\mathbb{A}_3 is a *harmonic* algebra, i.e. there exists a *harmonic* basis $\{e_1, e_2, e_3\} \subset \mathbb{A}_3$ satisfying the following conditions (see [5], [6], [7], [8], [9]):

$$(4) \quad e_1^2 + e_2^2 + e_3^2 = 0, \quad e_j^2 \neq 0 \text{ for } j = 1, 2, 3.$$

P. Ketchum [5] discovered that every function $\Phi(\zeta)$ analytic with respect to the variable $\zeta := xe_1 + ye_2 + ze_3$ with real x, y, z satisfies the equalities

$$(5) \quad \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right) \Phi(\zeta) = \Phi''(\zeta) (e_1^2 + e_2^2 + e_3^2) = 0$$

owing to the equality (4). I. Mel'nichenko [7] noticed that doubly differentiable in the sense of Gateaux functions form the largest class of functions Φ satisfying the equalities (5).

All harmonic bases in \mathbb{A}_3 are constructed by I. Mel'nichenko in [9].

Consider a harmonic basis

$$e_1 = 1, \quad e_2 = i + \frac{1}{2} i\rho_2, \quad e_3 = -\rho_1 - \frac{\sqrt{3}}{2} i\rho_2$$

in \mathbb{A}_3 and the linear envelope $E_3 := \{\zeta = x + ye_2 + ze_3 : x, y, z \in \mathbb{R}\}$ over the field of real numbers \mathbb{R} , that is generated by the vectors $1, e_2, e_3$. Associate with a domain $\Omega \subset \mathbb{R}^3$ the domain $\Omega_\zeta := \{\zeta = x + ye_2 + ze_3 : (x, y, z) \in \Omega\}$ in E_3 .

The algebra \mathbb{A}_3 have the unique maximal ideal $\{\lambda_1\rho_1 + \lambda_2\rho_2 : \lambda_1, \lambda_2 \in \mathbb{C}\}$ which is also the radical of \mathbb{A}_3 . Thus, it is obvious that the straight line $\{ze_3 : z \in \mathbb{R}\}$ is contained in the radical of algebra \mathbb{A}_3 .

\mathbb{A}_3 is a Banach algebra with the Euclidean norm

$$\|a\| := \sqrt{|\xi_1|^2 + |\xi_2|^2 + |\xi_3|^2},$$

where $a = \xi_1 + \xi_2e_2 + \xi_3e_3$ and $\xi_1, \xi_2, \xi_3 \in \mathbb{C}$.

We say that a continuous function $\Phi : \Omega_\zeta \rightarrow \mathbb{A}_3$ is *monogenic* in a domain $\Omega_\zeta \subset E_3$ if Φ is differentiable in the sense of Gateaux in every point of Ω_ζ , i. e. if for every $\zeta \in \Omega_\zeta$ there exists $\Phi'(\zeta) \in \mathbb{A}_3$ such that

$$\lim_{\varepsilon \rightarrow 0+0} (\Phi(\zeta + \varepsilon h) - \Phi(\zeta))\varepsilon^{-1} = h\Phi'(\zeta) \quad \forall h \in E_3.$$

For monogenic functions $\Phi : \Omega_\zeta \rightarrow \mathbb{A}_3$ we established basic properties analogous to properties of analytic functions of the complex variable: the Cauchy integral theorem, the Cauchy integral formula, the Morera theorem, the Taylor expansion (see [11]).

3. On existence of limiting values of a hypercomplex analogue of the Cauchy type integral. In what follows, $t_1, t_2, x, y, z \in \mathbb{R}$ and the variables x, y, z with subscripts are real. For example, x_0 and x_1 are real, etc.

Let $\Gamma_\zeta := \{\tau = t_1 + t_2e_2 : t_1 + it_2 \in \Gamma\}$ be the curve congruent to the curve $\Gamma \subset \mathbb{C}$. Consider the domain $\Pi_\zeta^\pm := \{\zeta = x + ye_2 + ze_3 : x + iy \in D^\pm, z \in \mathbb{R}\}$ in E_3 . By Σ_ζ we denote the common boundary of domains Π_ζ^+ and Π_ζ^- .

Consider the integral

$$(6) \quad \Phi(\zeta) = \frac{1}{2\pi i} \int_{\Gamma_\zeta} \varphi(\tau)(\tau - \zeta)^{-1} d\tau$$

with a continuous density $\varphi : \Gamma_\zeta \rightarrow \mathbb{R}$. The function (6) is monogenic in the domains Π_ζ^+ and Π_ζ^- , but the integral (6) is not defined for $\zeta \in \Sigma_\zeta$.

For the function $\varphi : \Gamma_\zeta \rightarrow \mathbb{R}$ consider the modulus of continuity

$$\omega_\varphi(\varepsilon) := \sup_{\tau_1, \tau_2 \in \Gamma_\zeta, \|\tau_1 - \tau_2\| \leq \varepsilon} |\varphi(\tau_1) - \varphi(\tau_2)|,$$

and a singular integral

$$\int_{\Gamma_\zeta} (\varphi(\tau) - \varphi(\zeta_0))(\tau - \zeta_0)^{-1} d\tau := \lim_{\varepsilon \rightarrow 0} \int_{\Gamma_\zeta \setminus \Gamma_\zeta^\varepsilon(\zeta_0)} (\varphi(\tau) - \varphi(\zeta_0))(\tau - \zeta_0)^{-1} d\tau,$$

where $\zeta_0 \in \Gamma_\zeta$ and $\Gamma_\zeta^\varepsilon(\zeta_0) := \{\tau \in \Gamma_\zeta : \|\tau - \zeta_0\| \leq \varepsilon\}$.

Below, in Theorem 1 in the case where the curve Γ satisfies the condition (2) and the modulus of continuity of the function φ satisfies a condition of the type (3), we establish the existence of certain limiting values of the integral (6) in points $\zeta_0 \in \Gamma_\zeta$ when ζ tends to ζ_0 from Π_ζ^+ or Π_ζ^- along

a curve that is not tangential to the surface Σ_ζ outside of the plane of curve Γ_ζ .

For the Euclidean norm in \mathbb{A}_3 the following inequalities are fulfilled:

$$(7) \quad \|ab\| \leq 2\sqrt{14}\|a\|\|b\| \quad \forall a, b \in \mathbb{A}_3,$$

$$(8) \quad \left\| \int_{\Gamma'_\zeta} \psi(\tau) d\tau \right\| \leq 9M \int_{\Gamma'_\zeta} \|\psi(\tau)\| \|d\tau\|$$

with the constant $M := \max\{1, \|e_2^2\|, \|e_2e_3\|, \|e_3^2\|\}$ for any measurable set $\Gamma'_\zeta \subset \Gamma_\zeta$ and all continuous functions $\psi : \Gamma'_\zeta \rightarrow \mathbb{A}_3$.

Lemma 1. *Let Γ be a closed Jordan rectifiable curve satisfying the condition (2) and the modulus of continuity of a function $\varphi : \Gamma_\zeta \rightarrow \mathbb{R}$ satisfies the condition of the type (3). If a point ζ tends to $\zeta_0 \in \Gamma_\zeta$ along a curve γ_ζ for which there exists a constant $m < 1$ such that the inequality*

$$(9) \quad |z| \leq m\|\zeta - \zeta_0\|$$

is fulfilled for all $\zeta = x + ye_2 + ze_3 \in \gamma_\zeta$, then

$$\lim_{\zeta \rightarrow \zeta_0, \zeta \in \gamma_\zeta} \int_{\Gamma_\zeta} (\varphi(\tau) - \varphi(\zeta_0))(\tau - \zeta)^{-1} d\tau = \int_{\Gamma_\zeta} (\varphi(\tau) - \varphi(\zeta_0))(\tau - \zeta_0)^{-1} d\tau.$$

Proof. Let $\varepsilon := \|\zeta - \zeta_0\|$. Consider the difference

$$\begin{aligned} & \int_{\Gamma_\zeta} (\varphi(\tau) - \varphi(\zeta_0))(\tau - \zeta)^{-1} d\tau - \int_{\Gamma_\zeta} (\varphi(\tau) - \varphi(\zeta_0))(\tau - \zeta_0)^{-1} d\tau \\ &= \int_{\Gamma_\zeta^{2\varepsilon}(\zeta_0)} (\varphi(\tau) - \varphi(\zeta_0))(\tau - \zeta)^{-1} d\tau - \int_{\Gamma_\zeta^{2\varepsilon}(\zeta_0)} (\varphi(\tau) - \varphi(\zeta_0))(\tau - \zeta_0)^{-1} d\tau \\ &+ (\zeta - \zeta_0) \int_{\Gamma_\zeta \setminus \Gamma_\zeta^{2\varepsilon}(\zeta_0)} (\varphi(\tau) - \varphi(\zeta_0))(\tau - \zeta)^{-1} (\tau - \zeta_0)^{-1} d\tau =: I_1 - I_2 + I_3. \end{aligned}$$

To estimate I_1 we choose a point $\zeta_1 = x_1 + y_1e_2$ on Γ_ζ such that $\|\zeta - \zeta_1\| = \min_{\tau \in \Gamma_\zeta} \|\tau - \zeta\|$. Using the inequalities (7) and (8), we obtain

$$\begin{aligned} \|I_1\| &= \left\| \int_{\Gamma_\zeta^{2\varepsilon}(\zeta_0)} (\varphi(\tau) - \varphi(\zeta_1))(\tau - \zeta)^{-1} d\tau + (\varphi(\zeta_1) - \varphi(\zeta_0)) \int_{\Gamma_\zeta^{2\varepsilon}(\zeta_0)} (\tau - \zeta)^{-1} d\tau \right\| \\ &\leq 18\sqrt{14}M \int_{\Gamma_\zeta^{2\varepsilon}(\zeta_0)} \|\varphi(\tau) - \varphi(\zeta_1)\| \|(\tau - \zeta)^{-1}\| \|d\tau\| \end{aligned}$$

$$+|\varphi(\zeta_1) - \varphi(\zeta_0)| \left\| \int_{\Gamma_\zeta^{2\varepsilon}(\zeta_0)} (\tau - \zeta)^{-1} d\tau \right\| =: I'_1 + I''_1.$$

It follows from Lemma 1.1 [9] that

$$(10) \quad (\tau - \zeta)^{-1} = \frac{1}{t - \xi} - \frac{z}{(t - \xi)^2} \rho_1 + \left(\frac{i}{2} \frac{y - t_2 - \sqrt{3}z}{(t - \xi)^2} + \frac{z^2}{(t - \xi)^3} \right) \rho_2$$

for all $\zeta = x + ye_2 + ze_3 \in \Pi_\zeta^\pm$ and $\tau = t_1 + t_2e_2 \in \Gamma_\zeta$, where $\xi := x + iy$ and $t := t_1 + it_2$. The following inequality follows from the relations (9) and (10):

$$(11) \quad \|(\tau - \zeta)^{-1}\| \leq c(m) \frac{1}{|t - \xi|},$$

where the constant $c(m)$ depends only on m .

Using the inequality $|t - \xi| \geq |t - \xi_1|/2$ with $\xi_1 := x_1 + iy_1$ and the inequality (11), we obtain:

$$\begin{aligned} \|I'_1\| &\leq 18\sqrt{14} Mc(m) \int_{\Gamma_\zeta^{2\varepsilon}(\zeta_0)} \frac{|\varphi(\tau) - \varphi(\zeta_1)|}{|t - \xi|} \|d\tau\| \\ &\leq 36\sqrt{14} Mc(m) \int_{\Gamma_\zeta^{2\varepsilon}(\zeta_0)} \frac{|\varphi(\tau) - \varphi(\zeta_1)|}{|t - \xi_1|} \|d\tau\| \\ &\leq 36\sqrt{14} Mc(m) \int_{[0,4\varepsilon]} \frac{\omega_\varphi(\eta)}{\eta} d\theta_{\xi_1}(\eta), \end{aligned}$$

where the last integral is understood as a Lebesgue–Stieltjes integral.

To estimate the last integral we use Proposition 1 [10] (see also the proof of Theorem 1 [4]) and the condition (2). So, we have

$$\int_{[0,4\varepsilon]} \frac{\omega_\varphi(\eta)}{\eta} d\theta_{\xi_1}(\eta) \leq \int_0^{8\varepsilon} \frac{\theta_{\xi_1}(\eta)\omega_\varphi(\eta)}{\eta^2} d\eta \leq c \int_0^{8\varepsilon} \frac{\omega_\varphi(\eta)}{\eta} d\eta \rightarrow 0, \quad \varepsilon \rightarrow 0,$$

where the constant c does not depend on ε .

To estimate I''_1 we introduce the domain $D_\zeta^{2\varepsilon}(\zeta_0) := \{\tau = t_1 + t_2e_2 : t_1 + it_2 \in D^+, \|\tau - \zeta_0\| \leq 2\varepsilon\}$ and its boundary $\partial D_\zeta^{2\varepsilon}(\zeta_0)$. Using the inequalities (8) and (11), we obtain:

$$\begin{aligned}
\|I_1''\| &\leq \omega_\varphi(\|\zeta_1 - \zeta_0\|) \left\| \int_{\Gamma_\zeta^{2\varepsilon}(\zeta_0)} (\tau - \zeta)^{-1} d\tau \right\| \\
&= \omega_\varphi(\|\zeta_1 - \zeta_0\|) \left\| \int_{\partial D_\zeta^{2\varepsilon}(\zeta_0)} (\tau - \zeta)^{-1} d\tau - \int_{\partial D_\zeta^{2\varepsilon}(\zeta_0) \setminus \Gamma_\zeta^{2\varepsilon}(\zeta_0)} (\tau - \zeta)^{-1} d\tau \right\| \\
&\leq \omega_\varphi(\|\zeta_1 - \zeta_0\|) \left(2\pi + 9Mc(m) \int_{\partial D_\zeta^{2\varepsilon}(\zeta_0) \setminus \Gamma_\zeta^{2\varepsilon}(\zeta_0)} \frac{\|d\tau\|}{|t - \xi|} \right) \\
&\leq \omega_\varphi(2\varepsilon) \left(2\pi + 9Mc(m) \frac{1}{\varepsilon} 2\pi 2\varepsilon \right) \rightarrow 0, \quad \varepsilon \rightarrow 0.
\end{aligned}$$

Estimating I_2 , by analogy with the estimation of I_1' , we obtain:

$$\|I_2\| \leq c \int_0^{4\varepsilon} \frac{\omega_\varphi(\eta)}{\eta} d\eta \rightarrow 0, \quad \varepsilon \rightarrow 0,$$

where the constant c does not depend on ε .

Using the inequality $|t - \xi| \geq |t - \xi_0|/2$, where the point $\xi_0 := x_0 + iy_0$ corresponds to the point $\zeta_0 = x_0 + y_0 e_2$, and using the relations (7), (8), (11) and (2), by analogy with the estimation of I_1' , we obtain:

$$\begin{aligned}
\|I_3\| &\leq 9M(2\sqrt{14})^2 \varepsilon \int_{\Gamma_\zeta \setminus \Gamma_\zeta^{2\varepsilon}(\zeta_0)} |\varphi(\tau) - \varphi(\zeta_0)| \|(\tau - \zeta)^{-1}\| \|(\tau - \zeta_0)^{-1}\| \|d\tau\| \\
&\leq c\varepsilon \int_{\Gamma_\zeta \setminus \Gamma_\zeta^{2\varepsilon}(\zeta_0)} \frac{|\varphi(\tau) - \varphi(\zeta_0)|}{|t - \xi| |t - \xi_0|} \|d\tau\| \leq c\varepsilon \int_{\Gamma_\zeta \setminus \Gamma_\zeta^{2\varepsilon}(\zeta_0)} \frac{|\varphi(\tau) - \varphi(\zeta_0)|}{|t - \xi_0|^2} \|d\tau\| \\
&\leq c\varepsilon \int_{[2\varepsilon, d]} \frac{\omega_\varphi(\eta)}{\eta^2} d\theta_{\xi_0}(\eta) \leq c\varepsilon \int_{2\varepsilon}^{2d} \frac{\theta_{\xi_0}(\eta) \omega_\varphi(\eta)}{\eta^3} d\eta \\
&\leq c\varepsilon \int_{2\varepsilon}^{2d} \frac{\omega_\varphi(\eta)}{\eta^2} d\eta \rightarrow 0, \quad \varepsilon \rightarrow 0,
\end{aligned}$$

where $d := \max_{\xi_1, \xi_2 \in \Gamma} |\xi_1 - \xi_2|$ is the diameter of Γ and c denotes different constants which do not depend on ε . The lemma is proved. \square

Let $\widehat{\Phi}^\pm(\zeta_0)$ be the boundary value of function (6) when ζ tends to $\zeta_0 \in \Gamma_\zeta$ along a curve γ_ζ for which there exists a constant $m < 1$ such that the inequality (9) is fulfilled for all $\zeta = x + ye_2 + ze_3 \in \gamma_\zeta$.

Theorem 1. Let Γ be a closed Jordan rectifiable curve satisfying the condition (2) and the modulus of continuity of a function $\varphi : \Gamma_\zeta \rightarrow \mathbb{R}$ satisfies the condition of the type (3). Then the integral (6) has boundary values $\widehat{\Phi}^\pm(\zeta_0)$ for all $\zeta_0 \in \Gamma_\zeta$ that are expressed by the formulas:

$$\widehat{\Phi}^+(\zeta_0) = \frac{1}{2\pi i} \int_{\Gamma_\zeta} (\varphi(\tau) - \varphi(\zeta_0))(\tau - \zeta_0)^{-1} d\tau + \varphi(\zeta_0)$$

$$\widehat{\Phi}^-(\zeta_0) = \frac{1}{2\pi i} \int_{\Gamma_\zeta} (\varphi(\tau) - \varphi(\zeta_0))(\tau - \zeta_0)^{-1} d\tau.$$

The theorem follows from the Lemma 1 and the equalities

$$\frac{1}{2\pi i} \int_{\Gamma_\zeta} \varphi(\tau)(\tau - \zeta)^{-1} d\tau = \frac{1}{2\pi i} \int_{\Gamma_\zeta} (\varphi(\tau) - \varphi(\zeta_0))(\tau - \zeta)^{-1} d\tau + \varphi(\zeta_0) \quad \forall \zeta \in \Pi_\zeta^+,$$

$$\frac{1}{2\pi i} \int_{\Gamma_\zeta} \varphi(\tau)(\tau - \zeta)^{-1} d\tau = \frac{1}{2\pi i} \int_{\Gamma_\zeta} (\varphi(\tau) - \varphi(\zeta_0))(\tau - \zeta)^{-1} d\tau \quad \forall \zeta \in \Pi_\zeta^-.$$

In comparison with Theorem 1, note that additional assumptions about the function φ are required for an existence of limiting values of the function (6) from Π_ζ^+ or Π_ζ^- on the boundary Σ_ζ . We are going to state these results in next papers.

REFERENCES

- [1] Davydov, N. A., *The continuity of an integral of Cauchy type in a closed domain*, Dokl. Akad. Nauk SSSR **64**, no. 6 (1949), 759–762 (Russian).
- [2] Salaev, V. V., *Direct and inverse estimates for a singular Cauchy integral along a closed curve*, Mat. Zametki **19**, no. 3 (1976), 365–380 (Russian).
- [3] Gerus, O. F., *Finite-dimensional smoothness of Cauchy-type integrals*, Ukrainian Math. J. **29**, no. 5 (1977), 490–493.
- [4] Gerus, O. F., *Some estimates of moduli of smoothness of integrals of the Cauchy type*, Ukrainian Math. J. **30**, no. 5 (1978), 594–601.
- [5] Ketchum, P. W., *Analytic functions of hypercomplex variables*, Trans. Amer. Math. Soc. **30** (1928), 641–667.
- [6] Kunz, K. S., *Application of an algebraic technique to the solution of Laplace's equation in three dimensions*, SIAM J. Appl. Math. **21**, no. 3 (1971), 425–441.
- [7] Mel'nichenko, I. P., *The representation of harmonic mappings by monogenic functions*, Ukrainian Math. J. **27**, no. 5 (1975), 499–505.
- [8] Mel'nichenko, I. P., *Algebras of functionally invariant solutions of the three-dimensional Laplace equation*, Ukrainian Math. J. **55**, no. 9 (2003), 1551–1559.
- [9] Mel'nichenko, I. P., Plaksa, S. A., *Commutative algebras and spatial potential fields*, Inst. Math. NAS Ukraine, Kiev, 2008 (Russian).
- [10] Plaksa, S. A., *Riemann boundary-value problem with infinite index of logarithmic order on a spiral contour. I*, Ukrainian Math. J. **42**, no. 11 (1990), 1509–1517.

- [11] Shpakivskyi, V. S., Plaksa, S. A., *Integral theorems and a Cauchy formula in a commutative three-dimensional harmonic algebra*, Bull. Soc. Sci. Lett. Łódź Sér. Rech. Déform. **60** (2010), 47–54.

S. A. Plaksa

Department of Complex Analysis and Potential Theory
Institute of Mathematics of the National Academy of Sciences of Ukraine
Tereshchenkivska St. 3
01601 Kiev-4
Ukraine
e-mail: plaksa@imath.kiev.ua

V. S. Shpakivskyi

Department of Complex Analysis and Potential Theory
Institute of Mathematics of the National Academy of Sciences of Ukraine
Tereshchenkivska St. 3
01601 Kiev-4
Ukraine
e-mail: shpakivskyi@mail.ru

Received September 30, 2011