

# On supersymmetries in nonrelativistic quantum mechanics

J. Beckers and N. Debergh<sup>a)</sup>

*Theoretical and Mathematical Physics, Institute of Physics, B.5, University of Liège, B-4000 Liège 1, Belgium*

A. G. Nikitin

*Institute of Mathematics, Academy of Sciences of the Ukrainian S.S.R., Repin Street, 3, Kiev 4, USSR*

(Received 22 August 1991; accepted for publication 30 August 1991)

One-dimensional nonrelativistic systems are studied when time-independent potential interactions are involved. Their supersymmetries are determined and their closed subsets generating kinematical invariance Lie superalgebras are pointed out. The study of even supersymmetries is particularly enlightened through the already known symmetries of the corresponding Schrödinger equation. Three tables collect the even, odd, and total supersymmetries as well as the invariance (super)algebras.

## I. INTRODUCTION

In the seventies, systematic studies of *symmetries* in nonrelativistic quantum mechanics (NRQM) had already been realized when the Schrödinger equation involves time-independent potential interactions.<sup>1-4</sup> Some of them have considered one-dimensional,<sup>2,4</sup> three-dimensional,<sup>1</sup> or  $n$ -dimensional<sup>3</sup> (space) systems.

Recently<sup>5</sup> we have added new information by visiting higher-order symmetries of such one-dimensional Schrödinger equations and by considering the third-order context as an effective case. (Remember that the usual well-known symmetries of the Schrödinger equation correspond to first- and second-order operators.)

Since the event of *supersymmetry* in theoretical particle physics<sup>6</sup> and its implications<sup>7</sup> in supersymmetric (nonrelativistic) quantum mechanics (SSQM), we are dealing with supersymmetric wave equations, including time-independent superpotentials [hereafter denoted  $W(x)$  in one-dimensional problems]. Such supersymmetric equations are usually put in the following matrix form:

$$i \partial_t \chi(x,t) = \left[ -\frac{1}{2} \partial_x^2 + \frac{1}{2} W^2(x) + \frac{1}{2} W'(x) \sigma_3 \right] \chi(x,t), \quad (1.1)$$

where  $\partial_t = \partial/\partial t$ , the prime refers to the space derivative  $\partial_x \equiv \partial/\partial x$ ,  $\sigma_3$  is the Pauli  $2 \times 2$  matrix  $\text{diag}(1, -1)$ , and correlatively  $\chi(x,t)$  is a two-component wave function. For simplicity we consider systems with unit masses and choose  $\hbar = 1$  in this quantum context.

In Hamiltonian form, Eq. (1.1) contains, as expected, the bosonic part  $H_b$  and the fermionic part  $H_f$  given by

$$H_b = -\frac{1}{2} \partial_x^2 + \frac{1}{2} W^2(x), \quad H_f = \frac{1}{2} W'(x) \sigma_3,$$

$$H_{ss} = H_b + H_f, \quad (1.2)$$

showing that superpotentials  $W(x)$  are simply related to usual potentials  $U(x)$  by

$$U(x) = \frac{1}{2} W^2(x). \quad (1.3)$$

Such supersymmetric equations (1.1) and characteristics (1.2) have already suggested new methods and results<sup>8,9</sup> by considering even and odd symmetries in connection with the so-called Lie extended method developed in quantum physics.<sup>10</sup>

As a particular context that has been recently visited,<sup>8</sup> let us mention the case of the one-dimensional supersymmetric harmonic oscillator corresponding to

$$W(x) = \omega x \Leftrightarrow U(x) = \frac{1}{2} \omega^2 x^2. \quad (1.4)$$

Here we want to study systematically the supersymmetries as well as their superstructures subtended by the SSQM-Schrödinger equation (1.1) for *arbitrary* superpotentials  $W(x)$ . Such a program will appear in the following sections where we will get a complete classification of all solvable (admissible) interactions admitting nontrivial supersymmetries.

The contents of this paper are distributed as follows. In Sec. II we come back on the Boyer results,<sup>3</sup> but for one-dimensional systems. We present the corresponding study in a particularly convenient way for our purpose in the supersymmetric context developed in Sec. III. There we first determine the even supersymmetries (Sec. III A), then second the odd ones (Sec. III B), and finally we superpose both contexts (Sec. III C) and get only four privileged classes of superpotentials. All the results are

<sup>a)</sup> Chercheur Institut Interuniversitaire des Sciences Nucléaires, Bruxelles.

quoted in three respective tables, where we mention the largest invariance Lie (super)algebras generated by the corresponding operators. Remember that these extended (super)symmetries do not necessarily close. Section IV finally contains some comments on isomorphic structures and some conclusions.

**II. GOING BACK ON SYMMETRIES IN NRQM**

In fact, Boyer<sup>3</sup> has already solved the problem of determining all the symmetries of the Schrödinger equation in  $n$  dimensions. Applied to the simplest one-dimensional context, this problem consists in the resolution of the following *invariance condition*:

$$[\Delta, Q] = i\lambda\Delta, \quad \lambda \equiv \lambda(x, t). \tag{2.1}$$

Here the Schrödinger equation defines the  $\Delta$  operator on the form

$$\Delta\psi(x, t) \equiv (-i\partial_t - \frac{1}{2}\partial_x^2 + U(x))\psi(x, t) = 0, \tag{2.2}$$

while we are searching for symmetry operators  $Q$  of (at most) second order with respect to space derivatives, i.e.,

$$Q \equiv ia(x, t)\partial_t + ib(x, t)\partial_x + ic(x, t), \tag{2.3}$$

$a$ ,  $b$ , and  $c$  being arbitrary functions. Let us mention that higher-order symmetry operators have recently been considered elsewhere in the same context.<sup>5</sup>

The above problem leads to a set of partial differential equations giving rise to  $x$ -independent functions  $a_0$ ,  $b_0$ , and  $c_0$  and to the following first-order partial differential equation on  $U(x)$ :

$$\begin{aligned} (\frac{1}{2}\dot{a}_0x + b_0)U'(x) + \dot{a}_0U(x) \\ = -\frac{1}{4}\ddot{a}_0x^2 - \ddot{b}_0x - \dot{c}_0 + (i/4)\ddot{a}_0, \end{aligned} \tag{2.4}$$

where the overdots evidently refer to time derivatives. The general solution of Eq. (2.4) appears as the sum of the general solution  $U_0(x)$  of the homogeneous equation and of a particular solution  $U_1(x)$  of the inhomogeneous equation. Due to the at most quadratic dependence on  $x$  in the inhomogeneous part of Eq. (2.4), Boyer has proposed to search for the general solution<sup>3</sup>

$$U(x) = U_0(x) + U_1(x), \tag{2.5}$$

with

$$\begin{aligned} U_1(x) &= \frac{1}{2}\alpha x^2 + \beta x + \gamma, \\ \alpha, \beta, \gamma &= \text{arbitrary constants}, \end{aligned} \tag{2.6}$$

and to distinguish three cases according to  $\alpha = 0$ ,  $\alpha = \omega^2 > 0$ , or  $\alpha = -\omega^2 < 0$ .

If the whole discussion takes place in Boyer's work,<sup>3</sup> let us notice here the following results and properties in order to exploit them in the supersymmetric context.

In the free case, i.e.,  $U = 0$  ( $U_0 = 0 = U_1$ ), as already obtained by Niederer,<sup>1</sup> we get at most six symmetries whose commutation relations lead to the largest invariance algebra seen as a semidirect sum ( $\square$ ) of the so-called "conformal" algebra  $so(2,1)$  and the Heisenberg algebra  $h(2)$ .

In the interacting case but when  $U_0 = 0$ , we want to point out a new result, i.e., the potential

$$U(x) = U_1(x) \tag{2.7}$$

leads to isomorphic structures  $so(2,1)\square h(2)$  as in the free case ( $\alpha = \beta = \gamma = 0$ ) whatever are the constants  $\alpha$ ,  $\beta$ , and  $\gamma$ . As particular cases, the harmonic oscillator corresponds to  $U_0 = 0$ ,  $\alpha = \omega^2$ ,  $\beta = \gamma = 0$  [see Eqs. (1.3) and (2.7)] while the linear potential to  $U_0 = 0$ ,  $\alpha = 0$ , i.e.,

$$U(x) = \beta x + \gamma. \tag{2.8}$$

Consequently, these cases also admit six symmetries and there exist changes of variables connecting all the equations including the potential forms (2.7) for arbitrary  $\alpha$ ,  $\beta$ , and  $\gamma$ , with the free Schrödinger equation. Let us just quote as an example some formulas corresponding to the change of variables between the case  $\alpha = \omega^2 > 0$  and the free case ( $\alpha = \beta = \gamma = 0$ ). It can be shown that the relations

$$\begin{aligned} t_1 &= (1/\omega) \tan^{-1} \omega t_2, \\ x_1 &= (1 + \omega^2 t_2^2)^{-1/2} (x + \beta/\omega^2) - \beta/\omega^2 \end{aligned} \tag{2.9}$$

correspond to a well-defined change of variables between the above interacting case (indices 1) and the free case (indices 2) implying a modification in the corresponding Schrödinger wave functions according to

$$\begin{aligned} \Psi_1(x_1, t_1) &= (1 + \omega^2 t_2^2)^{1/4} \exp \left[ \left( \frac{i\beta^2}{2\omega^3} - \frac{i\gamma}{\omega} \right) \tan^{-1} \omega t_2 \right. \\ &\quad \left. - \frac{i\beta^2}{2\omega^2} \frac{t_2}{1 + \omega^2 t_2^2} \right] \exp \left[ -\frac{it_2 x}{2(1 + \omega^2 t_2^2)} \right. \\ &\quad \left. \times (2\beta + \omega^2 x) \right] \Psi_2(x_2, t_2). \end{aligned} \tag{2.10}$$

As a more particular case included in this comment, i.e., the harmonic oscillator case with  $\alpha = \omega^2$ ,  $\beta = \gamma = 0$ , we immediately recover the Niederer result:<sup>1</sup>

$$\Psi_1(x_1, t_1) = (1 + \omega^2 t_1^2)^{1/4} \exp \left[ -\frac{it_1 \omega^2 x^2}{2(1 + \omega^2 t_1^2)} \right] \times \Psi_2(x_2, t_2). \tag{2.11}$$

When nontrivial potentials  $U_0(x) \neq 0$  are considered, only two cases have to be mentioned: either  $U_0$  is arbitrary, then all the potentials

$$U(x) = U_0 + \frac{1}{2}\alpha x^2 + \beta x + \gamma \tag{2.12}$$

lead to at least two symmetries (the corresponding Hamiltonian and the identity operator); or  $U_0$  takes the nonzero form

$$U_0(x) = \delta / (\mu x + \epsilon)^2, \quad \mu \neq 0, \tag{2.13}$$

then all the potentials

$$U(x) = \delta / (\mu x + \epsilon)^2 + \frac{1}{2}\alpha x^2 + \beta x + \gamma, \tag{2.14}$$

$$[(\mu, \epsilon) = (\alpha, \beta) \text{ if } \alpha \neq 0],$$

lead to four symmetries generating the direct sum  $so(2,1) \oplus I$ .

Such properties are of special interest for the following discussion of the supersymmetries of Eq. (1.1) in particular.

### III. GOING TO SUPERSYMMETRIES IN SSQM

Already considered<sup>8,9</sup> for only one-dimensional supersymmetric harmonic oscillators characterized by in-

teracting terms given in Eq. (1.4), the search for the largest number(s) of one-parameter Lie algebras can be extended to arbitrary (one-dimensional) systems. This asks for considering the problem (2.1) but with an operator  $\Delta_{SS}$  defined by the supersymmetric equation (1.1) and with symmetry operators  $Q$  containing even ( $\bar{0}$ ) and odd ( $\bar{1}$ ) parts<sup>8</sup> according to

$$Q = Q_{\bar{0}} + Q_{\bar{1}}, \tag{3.1}$$

referring to the expected graduation in the supercontext. Consequently, let us decompose our program in three steps: first to study the even supersymmetries (Sec. III A) by exploiting the results contained in Sec. II on symmetries; second to develop the new odd context and to get the corresponding supersymmetries (Sec. III B); and, third, to superpose the two sets of results (Sec. III C) in order to obtain the complete classification of solvable interactions in SSQM.

Let us just notice here that, in connection with Eq. (1.1), we understand that even and odd considerations are directly connected with  $2 \times 2$  Pauli matrices or more correctly with the Clifford algebra  $Cl_2 \equiv \{\sigma_0 \equiv I_2, \sigma_1, \sigma_2, \sigma_3\}$ , where we easily distinguish the even matrices ( $\sigma_0, \sigma_3$ ) and the odd ones ( $\sigma_1, \sigma_2$ ) in the usual fundamental representation.<sup>11</sup> Such a remark directly enlightens Secs. III A and III B.

#### A. Even supersymmetries in SSQM

The invariance condition (2.1) is replaced here by

$$[\Delta_{SS}, Q_{\bar{0}}] = i\lambda_{\bar{0}}\Delta_{SS}, \quad \lambda_{\bar{0}} = \lambda_{\bar{0}}(x, t), \tag{3.2}$$

where

$$\Delta_{SS} = \begin{pmatrix} -i\partial_t - \frac{1}{2}\partial_x^2 + \frac{1}{2}W^2(x) + \frac{1}{2}W'(x) & 0 \\ 0 & -i\partial_t - \frac{1}{2}\partial_x^2 + \frac{1}{2}W^2(x) - \frac{1}{2}W'(x) \end{pmatrix} \tag{3.3}$$

and

$$Q_{\bar{0}} = \begin{pmatrix} i(a_0 + a_3)\partial_t + i(b_0 + b_3)\partial_x + i(c_0 + c_3) & 0 \\ 0 & i(a_0 - a_3)\partial_t + i(b_0 - b_3)\partial_x + i(c_0 - c_3) \end{pmatrix} \tag{3.4}$$

due to the explicit forms of the even matrices  $\sigma_0$  and  $\sigma_3$ . Such a problem is equivalent to a set of two distinct ones in NRQM, as discussed in Sec. II. Indeed we get the two invariance conditions

$$[-i\partial_t - \frac{1}{2}\partial_x^2 + V_1(x), i(A_1\partial_t + B_1\partial_x + C_1)]$$

$$= i\Lambda_1(-i\partial_t - \frac{1}{2}\partial_x^2 + V_1(x)) \tag{3.5a}$$

and

$$[-i\partial_t - \frac{1}{2}\partial_x^2 + V_2(x), i(A_2\partial_t + B_2\partial_x + C_2)] = i\Lambda_2(-i\partial_t - \frac{1}{2}\partial_x^2 + V_2(x)), \tag{3.5b}$$

TABLE I. Even supersymmetries and associated superpotentials.

Classes	$\mathcal{E}$ supersymmetries and their Lie algebras	Number of $\mathcal{E}$ super-symmetries	Explicit forms of associated superpotentials	Characteristics
1	$[\text{so}(2,1) \square \mathfrak{h}(2)] \oplus [\text{so}(2,1) \square \mathfrak{h}(2)]$	12	$W(x) = ax + b$	Free case Linear case Harmonic oscillator <sup>9</sup>
2	$[\text{so}(2,1) \square \mathfrak{h}(2)] \oplus [\text{so}(2,1) \square \mathfrak{gl}(1)]$	10	$W(x) = \pm 1/(x + c)$	Coulomb ( $c=0$ )
3			$W(x) = ax \pm 1/x$	Calogero ( $a=\omega$ )
4	$[\text{so}(2,1) \oplus \mathfrak{gl}(1)] \oplus [\text{so}(2,1) \oplus \mathfrak{gl}(1)]$	8	$W(x) = ax + c/x$	$a \neq 0, c \neq \pm 1$
5			$W(x) = (fx + g)/(cx^2 + dx + h)$	$f \neq 0, c \neq 0, f \neq \pm c$
6	$[\text{so}(2,1) \square \mathfrak{h}(2)] \oplus [\text{so}(2) \oplus \mathfrak{gl}(1)]$	8	$W(x) = ax + b + \frac{ce^{\mp ax^2 \mp 2bx}}{d + cfe^{\mp ax^2 \mp 2bx} dx}$	$c \neq 0$ $(a,b) \neq (0,0)$
7		6	$W(x) = \frac{c}{dx + f} + \frac{g}{h(dx + f)^{\pm 2c/d} \pm [g/(d \mp 2c)](dx + f)}$	$d \neq 0, c \neq 0$ $d \neq \pm 2c$ $g \neq 0$
8	$[\text{so}(2,1) \oplus \mathfrak{gl}(1)] \oplus [\text{so}(2) \oplus \mathfrak{gl}(1)]$	6	$W(x) = \frac{c}{\pm 2cx + d} + \frac{f}{(\pm 2cx + d)(g + (f/2c) \ln  \pm 2cx + d )}$	$c \neq 0$ $f \neq 0$
9		6	$W(x) = ax + b + \frac{c}{dx + f} + \frac{g(dx + f)^{\mp 2c/d} e^{\mp ax^2 \mp 2bx}}{h \pm gfe^{\mp ax^2 \mp 2bx} (dx + f)^{\mp 2c/d} dx}$	$a \neq 0$ $d \neq 0$ $c \neq 0, \pm 1$
10	$[\text{so}(2) \oplus \mathfrak{gl}(1)] \oplus [\text{so}(2) \oplus \mathfrak{gl}(1)]$	4	$W(x) \neq$ above forms	...

where the arbitrary functions  $A_i, B_i, C_i$  and  $\Lambda_i$  ( $i = 1, 2$ ) are evidently simply related to the above  $a_0, a_3, \dots$ , while we have defined

$$V_1(x) = \frac{1}{2}W^2(x) + \frac{1}{2}W'(x) \tag{3.6a}$$

and

$$V_2(x) = \frac{1}{2}W^2(x) - \frac{1}{2}W'(x), \tag{3.6b}$$

these functions being nothing else than the potentials associated with superpartners.<sup>12</sup>

By combining and superposing the potentials (2.6), (2.7), (2.12), and (2.14) in known NRQM, we immediately deduce the cases leading to (at most) 12 even supersymmetries or to 10, 8, 6, and (at least) 4 of them. The (six) associated invariance Lie algebras are also easily determined (see Table I): the largest one is seen as the direct sum  $[\text{so}(2,1) \square \mathfrak{h}(2)] \oplus [\text{so}(2,1) \square \mathfrak{h}(2)]$  and the smallest one as  $[\text{so}(2) \oplus \mathfrak{gl}(1)] \oplus [\text{so}(2) \oplus \mathfrak{gl}(1)]$ . Such direct sums evidently come from the superposition of both projections by  $P_{\pm} = \frac{1}{2}(\sigma_0 \pm \sigma_3)$  of the ordinary symmetries obtained in NRQM. Moreover, by exploiting the explicit forms obtained for  $V_1$  and  $V_2$  in NRQM, it is possible through Eqs. (3.6) to get the corresponding superpoten-

tials  $W(x)$  entering in Eq. (1.1). Let us, for example, treat the case leading to the maximal number of even supersymmetries, i.e., let us consider

$$V_1(x) = \frac{1}{2}\alpha_1 x^2 + \beta_1 x + \gamma_1 \tag{3.7}$$

and

$$V_2(x) = \frac{1}{2}\alpha_2 x^2 + \beta_2 x + \gamma_2.$$

Through Eqs. (3.6) we readily get the linear superpotential

$$W(x) = ax + b, \quad a, b = \text{const}, \tag{3.8}$$

which also corresponds to the free case ( $a = b = 0$ ) as well as to the one-dimensional harmonic oscillator ( $a = \omega, b = 0$ ). Here are associated 12 even supersymmetries, an old result quoted in the even notations,<sup>9</sup>

$$[(H_B, C_{\pm}, I, P_{\pm})\sigma_0] \quad \text{and} \quad [(H_B, C_{\pm}, I, P_{\pm})\sigma_3]. \tag{3.9}$$

The general discussion leads to specific families of superpotentials given in Table I simultaneously with their associated invariance Lie algebras and their dimensions.

Let us point out the Coulomb-like and Calogero-like forms admitting ten even supersymmetries.

**B. Odd supersymmetries in SSQM**

With the operator  $\Delta_{SS} \equiv (3.3)$  we have now to exploit the invariance condition

$$[\Delta_{SS}, Q_{\bar{1}}] = i\lambda_{\bar{1}}\Delta_{SS}, \quad \lambda_{\bar{1}} = \lambda_{\bar{1}}(x, t), \tag{3.10}$$

where

$$Q_{\bar{1}} = \begin{pmatrix} 0 & (ia_1 + a_2)\partial_t + (ib_1 + b_2)\partial_x + ic_1 + c_2 \\ (ia_1 - a_2)\partial_t + (ib_1 - b_2)\partial_x + ic_1 - c_2 & 0 \end{pmatrix} \tag{3.11}$$

due to the explicit forms of the odd matrices  $\sigma_1$  and  $\sigma_2$ . Such a problem leads to a set of two third-order equations, which take the following forms, where the functions  $\alpha_i, \beta_i$  and  $\gamma_i$  are the  $x$ -independent parts of  $a_i, b_i$  and  $c_i$  ( $i = 1, 2$ ), respectively:

$$\begin{aligned} & \frac{i}{8}\alpha_2 W''' - \frac{3i}{4}\alpha_2 W^2 W' - \frac{i}{4}\ddot{\alpha}_2 x^2 W' - i\dot{\beta}_2 x W' \\ & - \frac{1}{4}\dot{\alpha}_2 W' + \gamma_2 W' - \frac{i}{4}\dot{\alpha}_1 W^2 - \frac{i}{2}\ddot{\alpha}_2 x W - i\dot{\beta}_2 W \\ & - \frac{i}{4}\ddot{\alpha}_1 x^2 - i\dot{\beta}_1 x - \frac{1}{4}\ddot{\alpha}_1 + \dot{\gamma}_1 = 0 \end{aligned} \tag{3.12a}$$

and

$$\begin{aligned} & -\frac{i}{8}\alpha_1 W''' + \frac{3i}{4}\alpha_1 W^2 W' + \frac{i}{4}\ddot{\alpha}_1 x^2 W' + i\dot{\beta}_1 x W' \\ & + \frac{1}{4}\dot{\alpha}_1 W' - \gamma_1 W' - \frac{i}{4}\dot{\alpha}_2 W^2 + \frac{i}{2}\ddot{\alpha}_1 x W + i\dot{\beta}_1 W \\ & - \frac{i}{4}\ddot{\alpha}_2 x^2 - i\dot{\beta}_2 x - \frac{1}{4}\ddot{\alpha}_2 + \dot{\gamma}_2 = 0. \end{aligned} \tag{3.12b}$$

These inhomogeneous nonlinear equations simultaneously admit the particular solution (3.8), so that we can in this supersymmetric context search for the general solution

$$W(x) = W_0(x) + W_1(x), \tag{3.13}$$

where we accept

$$W_1(x) = ax + b, \quad a, b = \text{const}, \tag{3.14}$$

and where we have to discuss the two cases  $W_0 = 0$  or  $W_0 \neq 0$ . Such a discussion is analogous to the one developed in NRQM through Eqs. (2.5), (2.7), and (2.12).

Let us (first) consider  $W_0(x) = 0$ , so that

$$W(x) = W_1(x) = ax + b. \tag{3.15}$$

Here again we can distinguish  $a = 0$  and  $a \neq 0$ . Both cases lead to 12 (odd) supersymmetries. In particular, if  $a = \omega, b = 0$ , this context coincides with the supersymmetric harmonic oscillator already visited and characterized by this maximal number of odd supersymmetries.<sup>9</sup>

Then let us (second) take  $W_0 \neq 0$  and insert the value (3.13) in Eqs. (3.12). For arbitrary  $W_0$ , it is once again interesting to distinguish between the two cases  $a = 0$  and  $a \neq 0$ . Such a discussion leads to at least two odd supersymmetries, which appear on the forms

$$Q_{\bar{1}}^{(1)} = i\sigma_1 \partial_x - \sigma_2 W(x) \quad \text{and} \quad Q_{\bar{1}}^{(2)} = i\sigma_2 \partial_x + \sigma_1 W(x). \tag{3.16}$$

We also point out that they generate the simplest  $N = 2$  superalgebra sqm (2) initially introduced by Witten.<sup>7</sup> We effectively have

$$\begin{aligned} \{Q_{\bar{1}}^{(1)}, Q_{\bar{1}}^{(1)}\} &= \{Q_{\bar{1}}^{(2)}, Q_{\bar{1}}^{(2)}\} \\ &= -2\partial_x^2 + 2W^2(x) + 2\sigma_3 W'(x) \\ &= 4H_{SS} = 4i\partial_t, \end{aligned} \tag{3.17}$$

$$\{Q_{\bar{1}}^{(1)}, Q_{\bar{1}}^{(2)}\} = 0, \quad [Q_{\bar{1}}^{(i)}, H_{SS}] = 0, \quad i = 1, 2.$$

In order to complete our classification of admissible interactions characterized by odd supersymmetries, we now want to determine the impact due to each symmetry on the superpotentials  $W_0(x)$  left arbitrary in Eq. (3.13). This can be studied by quoting the general forms of our arbitrary functions  $\alpha_i, \beta_i$  and  $\gamma_i$  (depending only on  $t$ ) appearing in Eq. (3.12). When  $a = 0$ , these functions are obtained in terms of the 12 arbitrary constants  $A_{(0)}, B_{(0)}, C_{(0)}, D_{(0)}, E_{(0)}, F_{(0)}, G_{(0)}, H_{(0)}, K_{(0)}, L_{(0)}, M_{(0)}$ , and  $N_{(0)}$  (the subscript refers to this  $a = 0$  case) as follows:

$$\begin{aligned}
 \alpha_1(t) &= \frac{1}{2}A_{(0)}t^2 + B_{(0)}t + C_{(0)}, \\
 \alpha_2(t) &= \frac{1}{2}D_{(0)}t^2 + E_{(0)}t + F_{(0)}, \\
 \beta_1(t) &= -\frac{1}{4}bD_{(0)}t^2 + G_{(0)}t + H_{(0)}, \\
 \beta_2(t) &= \frac{1}{4}bA_{(0)}t^2 + K_{(0)}t + L_{(0)}, \tag{3.18} \\
 \gamma_1(t) &= \frac{1}{4}A_{(0)}t \left( 1 + \frac{3i}{2}b^2t \right) + \frac{i}{4}B_{(0)}b^2t + iK_{(0)}bt + M_{(0)}, \\
 \gamma_2(t) &= \frac{1}{4}D_{(0)}t \left( 1 + \frac{3i}{2}b^2t \right) + \frac{i}{4}E_{(0)}b^2t - iG_{(0)}bt + N_{(0)},
 \end{aligned}$$

while, when  $a \neq 0$ , they are given by

$$\begin{aligned}
 \alpha_1(t) &= A \exp(iat) + B \exp(-iat) \\
 &+ iC \exp(3iat) - iD \exp(-3iat),
 \end{aligned}$$

$$\begin{aligned}
 \alpha_2(t) &= E \exp(iat) + F \exp(-iat) + C \exp(3iat) \\
 &+ D \exp(-3iat), \\
 \beta_1(t) &= (i/2)b[A \exp(iat) - B \exp(-iat)] \\
 &+ i[K \exp(2iat) - L \exp(-2iat)] \\
 &- \frac{3}{2}b[C \exp(3iat) - D \exp(-3iat)] + M, \\
 \beta_2(t) &= (i/2)b[E \exp(iat) - F \exp(-iat)] \\
 &+ K \exp(2iat) + L \exp(-2iat) + (3i/2)b \\
 &\times [C \exp(3iat) - D \exp(-3iat)] + G, \tag{3.19}
 \end{aligned}$$

---


$$\begin{aligned}
 \gamma_1(t) &= \exp(iat) \left( iP + \frac{1}{4}aE + \frac{1}{4}b^2E + \frac{i}{4}aA + \frac{i}{4}b^2A \right) - \exp(-iat) \left( iQ - \frac{1}{4}aF + \frac{1}{4}b^2F + \frac{i}{4}aB - \frac{i}{4}b^2B \right) \\
 &+ 2ib[K \exp(2iat) + L \exp(-2iat)] - C \exp(3iat) \left( \frac{3}{4}a + \frac{9}{4}b^2 \right) - D \exp(-3iat) \left( \frac{3}{4}a - \frac{9}{4}b^2 \right), \\
 \gamma_2(t) &= P \exp(iat) + Q \exp(-iat) + 2b[K \exp(2iat) - L \exp(-2iat)] + (3i/4)C \exp(3iat) \\
 &\times (3b^2 + a) + (3i/4)D \exp(-3iat)(3b^2 - a),
 \end{aligned}$$

showing once again 12 arbitrary constants.

As an example, let us introduce (3.13) and (3.14) in Eqs. (3.12) when  $a = 0$ , i.e., when  $W(x) = W_0(x) + b$ . By exploiting the relations (3.18) when only the constant  $A_{(0)}$  is nonzero, we finally obtain three time-independent conditions on  $W_0(x)$ , which are

$$\begin{aligned}
 W''_0 - 6W_0(W_0 + 2b)W'_0 &= 0, \\
 2bxW'_0 + W_0(W_0 + 4b) &= 0, \\
 x(xW'_0 + 2W_0) &= 0.
 \end{aligned}$$

They are easily handled for getting the unique solution  $W_0(x) = 0$ . This  $A_{(0)}$ -context leads to the symmetry operator

$$Q_1^{A_{(0)}} = \sigma_1 \left( \frac{i}{2}t^2\partial_t + \frac{i}{2}tx\partial_x + \frac{1}{4}x + \frac{1}{4}bt^2W - \frac{1}{8}t^2W^2 \right)$$

---


$$\begin{aligned}
 &+ \frac{i}{4}t - \frac{3}{8}b^2t^2 \Big) + \sigma_2 \left( \frac{i}{4}bt^2\partial_x + \frac{1}{2}btx + \frac{i}{8}t^2W' \right. \\
 &\left. - \frac{i}{4}t^2W\partial_x - \frac{1}{2}txW \right). \tag{3.20}
 \end{aligned}$$

By collecting all the similar information for the whole set of odd (super)symmetries associated with the case  $a = 0$ , we finally determine besides the maximal (12) and minimal (2) numbers of odd supersymmetries already obtained that only three intermediate cases can occur: either  $W(x) = \pm 1/x$  admits ten (odd) symmetries; or  $W(x) = \pm c/x$  admits four (odd) symmetries if  $c \neq \pm 1$ ; or  $W(x) = b + W_0(x)$  admits four (odd) symmetries if we take account of an extradependence in terms of Legendre functions.

When  $a \neq 0$ , the developments are more elaborate but the complete results can once again be obtained through the resolution of relatively complicated nonlinear differential equations. Just as an example with

$$W(x) = W_0(x) + ax + b, \quad a \neq 0, \quad (3.21)$$

let us use Eqs. (3.12), (3.13), (3.14) and insert Eq. (3.19) when only the arbitrary constant  $C$  is nonzero. This case is then characterized by the functions

$$\begin{aligned} \alpha_1(t) &= iC \exp(3iat), & \alpha_2(t) &= C \exp(3iat), \\ \beta_1(t) &= -\frac{3}{2}bC \exp(3iat), \\ \beta_2(t) &= (3i/2)bC \exp(3iat), & (3.22) \\ \gamma_1(t) &= -\frac{3}{4}aC \exp(3iat) - \frac{3}{4}b^2C \exp(3iat), \\ \gamma_2(t) &= (9i/4)b^2C \exp(3iat) + (3i/4)aC \exp(3iat). \end{aligned}$$

Thus we get the following third-order (nonlinear) differential equation on  $W_0(x)$ :

$$\begin{aligned} W_0''' &= 6W_0^2W_0' - 12(ax+b)^2W_0' + 12(ax+b)W_0W_0' \\ &\quad - 36a(ax+b)W_0, \end{aligned} \quad (3.23)$$

or, in terms of  $W(x) \equiv (3.21)$ ,

$$\begin{aligned} W''' &= 6W^2W' - 18(ax+b)^2W' - 6aW^2 \\ &\quad - 36a(ax+b)W + 54a(ax+b)^2. \end{aligned} \quad (3.24)$$

Such an equation has already been quoted and solved in the literature.<sup>13</sup> It corresponds to the derivative of a Painlevé IV-type equation and leads to the solution<sup>13</sup>

$$W(x) = \epsilon \frac{dP_4}{dx}(0, \delta; x) + \frac{\epsilon}{\mu} \left( x + \frac{b}{a} \right). \quad (3.25)$$

Now, by collecting all the corresponding information for the whole set of odd (super)symmetries associated with the case  $a \neq 0$ , we see that only five intermediate cases can occur: Either  $W(x) = \pm 1/(ax+b) + ax+b$  admits ten (odd) symmetries; or  $W(x) = c/(ax+b) + ax+b$  admits four (odd) symmetries if  $c \neq \pm 1$ ; or  $W(x) = W_0(x) + ax + b$  admits four (odd) symmetries if  $W_0$  is a solution of Eq. (3.23); or

$$W(x) = \frac{c \exp(\pm ax^2)}{d \mp c \int \exp(\pm ax^2) dx} + ax$$

admits three (odd) symmetries; or

$$W(x) = \frac{c \exp(\frac{1}{2}b^2x^2 \mp 2bx)}{d \pm c \int \exp(\frac{1}{2}b^2x^2 \mp 2bx) dx} \mp \frac{1}{2}b^2x + b$$

admits three (odd) symmetries.

Consequently, we can summarize in Table II the specific families of superpotentials admitting odd supersymmetries.

Already mentioned in the even context, we notice once again that the free case, the linear case, and the harmonic oscillator case are also involved within the same class 1' and that their 12 odd supersymmetries can be quoted [in correspondence with (3.9)] in the forms

$$[(H_B, C_{\pm}, I, P_{\pm})\sigma_1] \quad \text{and} \quad [(H_B, C_{\pm}, I, P_{\pm})\sigma_2] \quad (3.26)$$

according to old notations<sup>9</sup> and remembering that  $\sigma_1$  and  $\sigma_2$  are the two odd matrices of  $Cl_2$ .

### C. Supersymmetries and invariance superalgebras in SSQM

We can now superpose the results on even and odd supersymmetries collected in Tables I and II, respectively. If such a superposition is relatively direct for superpotentials belonging to the classes 1, 2, 3, 4 and 1', 2', 3', 4' due to their similar forms, it is evident that the superposition of all other information becomes relatively tedious and that it is not interesting to insist on the corresponding properties.

Besides the information on the maximal number 24 ( $= 12 + 12$ ) of supersymmetries associated with the classes 1 and 1', as well as on the minimal number 6 ( $= 4 + 2$ ) associated with the classes 10 and 10', we can distinguish what are the superstructures generated by some subsets of operators. We have determined that only three (closed) invariance Lie superalgebras can be pointed out. We have summarized these structures and their associated properties in Table III. We evidently recover the expected results<sup>9</sup> for the supersymmetric harmonic oscillator but also find new ones for the Coulomb and the Calogero problems.

We notice that only a few cases are privileged in this supersymmetric context when we require invariance superstructures.

### IV. COMMENTS AND CONCLUSIONS

As already noticed (and expected), the free case leads to the largest number (24) of supersymmetries and to the largest (13-dimensional) kinematical invariance superalgebra. These results are also valid for the linear case and the harmonic oscillator context, which, both, are isomorphic to the free case as it is true in NRQM. In fact, let us point out that for those three cases we are dealing with the superpotential given by (see Table III):

$$W(x) = ax + b, \quad (4.1)$$

so that the corresponding supersymmetric Hamiltonian (1.2) is

TABLE II. Odd supersymmetries and associated superpotentials.

Classes	Number of $\mathcal{O}$ supersymmetries	Explicit forms of associated superpotentials	Characteristics
1'	12	$W(x) = ax + b$	Free case Linear case Harmonic oscillator <sup>9</sup>
2'	10	$W(x) = c/x$	$c = \pm 1$
3'		$W(x) = ax + b + c/(x + b/a)$	$a \neq 0, c = \pm 1$
4'	4	$W(x) = ax + b + c/(x + b/a)$	$a \neq 0, c \neq \pm 1$
5'		$W(x) = c/x$	$c \neq \pm 1$
6'	4	$W(x) = b + W_0(x)$ if $W''_0 - 6W_0^2W'_0 - 12bW_0W'_0 - 6b^2W'_0 = 0$	$a = 0$
7'		$W(x) = ax + b + W_0(x)$ if $W''_0 - 6W_0^2W'_0 - 12(ax + b)W_0W'_0 + 12(ax + b)^2W'_0 + 36a(ax + b)W_0 = 0$	$a \neq 0$
8'	3	$W(x) = ax + \frac{c \exp(\pm ax^2)}{d \mp cf \exp(\pm ax^2) dx}$	$a \neq 0, c \neq 0$
9'		$W(x) = \mp \frac{1}{2} b^2 x + b + \frac{c \exp(\frac{1}{2} b^2 x^2 \mp 2bx)}{d \pm cf \exp(\frac{1}{2} b^2 x^2 \mp 2bx) dx}$	$b \neq 0, c \neq 0$
10'	2	$W(x) = W_0(x) + ax + b, W_0 \neq$ above forms	...

$$H_{SS} = \frac{1}{2}p^2 + \frac{1}{2}(a^2x^2 + 2abx + b^2) + \frac{1}{2}a\sigma_3, \quad (4.2) \quad \Delta_{SS} \equiv -i\partial_t + H_{SS}, \quad (4.4)$$

Then, by using the unitary transformation

$$U = \exp[(i/2)at\sigma_3], \quad (4.3)$$

we immediately get for the operator  $\Delta_{SS} \equiv (3.3)$ , i.e.,

that

$$U\Delta_{SS}U^{-1} = [-i\partial_t - \frac{1}{2}\partial_x^2 + \frac{1}{2}(a^2x^2 + 2abx + b^2)]I_2. \quad (4.5)$$

TABLE III. Total number of supersymmetries and associated Lie superalgebras.

Superpotentials	Number of supersymmetries			Invariance Lie superalgebras		Characteristics
	$\mathcal{E}$	$\mathcal{O}$	$N = \mathcal{E} + \mathcal{O}$	Dimension ( $d$ )	Superstructure	
1 $W(x) = ax + b$	12	12	24	$d = 13$	$osp(2/2) \square sh(2/2)$	Free case Linear case Harmonic oscillator <sup>9</sup>
2 $W(x) = c/x \begin{cases} c = \pm 1, \\ c \neq \pm 1, c \neq 0 \end{cases}$	10 8	10 4	20 12	$d = 7$	$[osp(2/1) \square so(2)] \oplus gl(1)$	Coulomb
3 $W(x) = ax + c/x \begin{cases} c = \pm 1, \\ c \neq \pm 1, c \neq 0 \end{cases}$	10 8	10 4	20 12	$d = 7$	$[osp(2/1) \square so(2)] \oplus gl(1)$	Calogero
4 $W(x) \neq$ above forms	8, 6, or 4	4, 3, or 2	$12 < N < 6$	$d = 5$	$[sqm(2) \square so(2)] \oplus gl(1)$	...



This expression corresponds to the NRQM context characterized by the potential (2.6), but amplified by the identity matrix  $I_2$  belonging to the Clifford algebra  $Cl_2$ . We thus recover the six symmetries obtained by Niederer<sup>1</sup> multiplied here (four times) by the elements of  $Cl_2$  leading to  $12\mathcal{E} + 12\mathcal{O}$ , i.e., to 24 supersymmetries<sup>9</sup> characterized in Eqs. (3.9) and (3.26). Among them only 13 close and lead to the largest kinematical invariance superalgebra  $\text{osp}(2/2) \square \text{sh}(2/2)$ .<sup>14</sup> Associated with these comments are changes of variables easily determined from the formulas (2.9)–(2.11) and directly connected with other results<sup>1,15</sup> when, for example,  $a = \omega$  and  $b = 0$ .

Let us also insist on the specific interest of the simplest Witten superalgebra  $\text{sqm}(2)$  considered in Eqs. (3.17) and recovered as a part of the minimal closed superstructure found in Table III. We immediately notice that

$$\text{osp}(2/2) \supset \text{osp}(2/1) \supset \text{sqm}(2), \quad (4.6)$$

a physical chain of particular interest.

Besides the general conclusions that can be drawn from the tables and more particularly from Table III, let us recall that different types of superpotentials have already been studied in SSQM (see more particularly the reviews of D'Hoker *et al.*<sup>16</sup> and of Lahiri *et al.*<sup>17</sup>). All these superpotentials fall into one of the classes 1–4 contained in Table III, as well as those we have considered as partner potentials in parasupersymmetric quantum mechanics.<sup>18</sup>

#### ACKNOWLEDGMENT

We want to thank Dr. J. Lombet for interesting information concerning nonlinear differential equations.

- <sup>1</sup>U. Niederer, *Helv. Phys. Acta* **45**, 802 (1972); **46**, 191 (1973).
- <sup>2</sup>R. L. Anderson, S. Kumei, and C. E. Wulfman, *Rev. Mex. Fis.* **21**, 1 (1972).
- <sup>3</sup>C. P. Boyer, *Helv. Phys. Acta* **47**, 590 (1974).
- <sup>4</sup>W. Miller, Jr., *Symmetry and Separation of Variables*, Encyclopedia of Mathematics and its Applications, (Addison-Wesley, London, 1977), Vol. 4.
- <sup>5</sup>J. Beckers, N. Debergh, and A. G. Nikitin, *J. Phys. A* **24** (1991).
- <sup>6</sup>P. Fayet and S. Ferrara, *Phys. Rep.* **32**, 250 (1977); *Supersymmetry in Physics* (North-Holland, Amsterdam, 1985).
- <sup>7</sup>E. Witten, *Nucl. Phys. B* **188**, 513 (1981).
- <sup>8</sup>J. Beckers and N. Debergh, *J. Phys. A* **23**, L353 (1990).
- <sup>9</sup>J. Beckers and N. Debergh, *Helv. Phys. Acta* **64**, 24 (1991).
- <sup>10</sup>S. Lie, *Vorlesungen über Differentialgleichungen mit Bekarmten Infinitesimalen Transformationen* (Teubner, Leipzig, 1891); L. V. Ovsiannikov, *Group Analysis of Differential Equations* (Academic, New York, 1982); P. J. Olver, *Applications of Lie Groups to Differential Equations* (Springer-Verlag, New York, 1986); W. I. Fushchich and A. G. Nikitin, *Symmetries of Maxwell's Equations* (Reidel, Dordrecht, 1987).
- <sup>11</sup>D. H. Sattinger and O. L. Weaver, *Lie Groups and Algebras with Applications to Physics, Geometry, and Mechanics* (Springer-Verlag, Berlin, 1986).
- <sup>12</sup>B. Mielnik, *J. Math. Phys.* **25**, 3387 (1984); A. A. Andrianov, N. V. Borisov, and M. V. Ioffe, *Phys. Lett. A* **105**, 19 (1984); M. M. Nieto, *Phys. Lett. B* **145**, 208 (1984); C. V. Sukumar, *J. Phys. A* **18**, 2917 (1985).
- <sup>13</sup>F. J. Bureau, *Ann. Matem. Pura Appl.* **LXIV**, 229 (1964); **LXVI**, 1 (1964).
- <sup>14</sup>J. Beckers and V. Hussin, *Phys. Lett. A* **118**, 319 (1986).
- <sup>15</sup>J. Beckers, D. Dehin, and V. Hussin, *J. Phys. A*, **20**, 1137 (1987); **21**, 651 (1988).
- <sup>16</sup>E. D'Hoker, V. A. Kostelecky, and L. Vinet, *Spectrum Generating Superalgebras in Dynamical Groups and Spectrum Generating Algebras*, edited by A. Barut, A. Bohm, and Y. Ne'eman (World Scientific, Singapore, 1989).
- <sup>17</sup>A. Lahiri, P. K. Roy, and B. Bagchi, *Int. J. Mod. Phys. A* **5**, 1383 (1990).
- <sup>18</sup>J. Beckers and N. Debergh, *Nucl. Phys. B* **340**, 767 (1990); *Z. Phys. C* **51**, 519 (1991).