

INVOLUTIVE SYMMETRIES, SUPERSYMMETRIES  
AND REDUCTIONS OF THE DIRAC EQUATION<sup>1</sup>

J.Niederle

Institute of Physics,  
Academy of Sciences of the Czech Republic,  
Na Slovance 2, Prague 8

and

A.G.Nikitin

Institute of Mathematics,  
National Academy of Sciences of the Ukraine,  
Tereshchenkivs'ka Street 3, Kiev-4, Ukraine

Abstract. A new algebra of involutive symmetries of the Dirac equation is found. This algebra is used to reduce the Dirac equation for a charged particle, interacting with an external field and to describe hidden supersymmetries of this equation. Reducibility of a class of equations of supersymmetric quantum mechanics is established.

PACS numbers: 11.30.Ly, 11.30.Pb, 11.10.Qr, 03.65.Fd, 33.10.Cs

---

<sup>1</sup>This publication is based on work sponsored by the US-Czechoslovakia Science and Technological Joint Fund under Project Number 92067 and by Grant of EU under Project Number CIPA-CT92-3006

## I. INTRODUCTION

It is well-known that symmetries of differential equations form powerful tools for the study of symmetries of these equations. They are used to separate variables [1], to derive conservation laws [2], to construct exact solutions of linear and nonlinear differential equations [3-6], to find spectra of linear differential operators [7,8], etc. etc.

In this paper we investigate special involutive symmetries of the Dirac equation. It is well known that this equation is invariant w.r.t. the extended Poincare group. Pauli, Gursev, Plebanski and Pursey [9] found the additional  $SL(2,C)$  symmetry of the Dirac equation, which is realized by antilinear transformations (i.e., including the complex conjugation). Hidden  $SL(2,C)$  symmetry of this equation (generated by linear non-local integro-differential operators and by first order differential operators with matrix coefficients) was described in papers [10] and [11] (refer also to [8]).

In this paper we present a new symmetry algebra of the Dirac equation. It is specified by the following features:

- (i) All its basis elements are involutions;
- (ii) It includes proper discrete symmetries (like reflections  $P$ ,  $T$  and charge conjugation  $C$ ) as well as finite rotations;
- (iii) It is a finite dimensional Lie algebra whose dimension is much more extended than dimensions of other finite symmetry algebras of the Dirac equation.

We use this symmetry algebra for two purposes. First, to reduce the Dirac equation to two uncoupled subsystems or even to

four uncoupled one-component equations. The necessary and sufficient condition for existence of such a reduction is that the components of the vector-potential  $A_\mu$  (treated as given functions of  $x_0, x_1, x_2, x_3$ ) have definite parities, i.e., are invariant (up to a sign) under reflections of  $x_\mu$ .

The other important application of involutive symmetries is searching for systems with exact supersymmetry (SUSY). Using the former algebra we extend the list of known systems with N=2 SUSY [12,13] and find a class of external potentials for the Dirac equation which generate extended SUSY.

In Section II we describe the involutive symmetry algebra of the Dirac equation. The corresponding reductions for the Dirac equation are discussed in Section III and presented explicitly in the Appendix.

Sections IV and V are devoted to reduction of the Dirac oscillator and to searching for exact SUSY. Section VI includes application of the reduction technique to SUSY quantum mechanics.

## II. INVOLUTIVE SYMMETRIES OF THE DIRAC EQUATION

We start with the free Dirac equation

$$L_0 \psi = (\gamma^\mu p_\mu - m) \psi = 0 \quad (2.1)$$

which is invariant w.r.t. the complete Lorentz group. Here  $\gamma_\mu$  ( $\mu=0,1,2,3$ ) are the Dirac matrices with diagonal  $\gamma_5 = i\gamma_0\gamma_1\gamma_2\gamma_3$ :

$$\gamma_0 = \begin{pmatrix} 0 & I_2 \\ I_2 & 0 \end{pmatrix}, \quad \gamma_a = \begin{pmatrix} 0 & -\sigma_a \\ \sigma_a & 0 \end{pmatrix}, \quad \gamma_5 = i\gamma_0\gamma_1\gamma_2\gamma_3 = \begin{pmatrix} I_2 & 0 \\ 0 & -I_2 \end{pmatrix}, \quad (2.2)$$

## 4

$\sigma_a$ ,  $a=1,2,3$  are the Pauli matrices and  $I_2$  denotes a 2x2 unit matrix.

Let us note that this equation admits also non-Lie symmetries [8, 10, 11].

Here we study the class of involutive symmetries of (2.1). Such symmetries form a subset of the complete Lorentz group, which is defined by reflections of coordinate axes, rotations by the angle  $\pi$  w.r.t. a given axis (each of them can be reduced to a reflection of a pair of axes) and by products of these transformations. There are sixteen of them, and they form a finite group composed of:

- four reflections of coordinates  $x_\mu$ :

$$x_\mu \rightarrow (\theta_\lambda X)_\mu = (1 - 2\delta_{\lambda\mu}) x_\mu, \quad \lambda=0, 1, 2, 3, \quad (2.3a)$$

- six reflections of pairs of coordinates:

$$x_\mu \rightarrow (\theta_{\lambda\sigma} X)_\mu = (1 - 2\delta_{\lambda\mu} - 2\delta_{\sigma\mu}) x_\mu, \quad \lambda \neq \sigma, \quad \lambda, \sigma=0, 1, 2, 3, \quad (2.3b)$$

- four reflections of triplets of coordinates:

$$x_\mu \rightarrow (\theta'_\lambda X)_\mu = (2\delta_{\lambda\mu} - 1) x_\mu, \quad \lambda=0, 1, 2, 3, \quad (2.3c)$$

- a complete reflection of all coordinates:

$$x_\mu \rightarrow (\theta X)_\mu = -x_\mu, \quad (2.4a)$$

and the identity transformation

$$x_\mu \rightarrow (IX_\mu) = x_\mu \quad (2.4b)$$

( $\mu=0,1,2,3$ ; no sums over  $\mu$  in (2.3)). We will use also the following notation for reflections (2.3), (2.4):

$$\theta = \theta_{54}, \quad \theta_v = \theta_{5v}, \quad \theta'_v = \theta_{4v}. \quad (2.5)$$

We see that for  $\lambda, \sigma = 1, 2, 3$  transformation (2.3b) are rotations while (2.3a), (2.3c) and (2.3b) for zero  $\lambda$  or  $\sigma$  are proper reflections.

The corresponding symmetries of the Dirac equation form a projective representation [16] of the 16-dimensional group (2.3), (2.4) and have the following form

$$\psi(x) \rightarrow R_{kl} \psi(x) = \hat{S}_{kl} \hat{\theta}_{kl} \psi(x) \equiv \hat{S}_{kl} \psi(\theta_{kl} x), \quad (2.6)$$

where

$$\begin{aligned} \hat{S}_{\mu\nu} &= \tilde{\gamma}_\mu \tilde{\gamma}_\nu, & \hat{S}_{5\mu} &= \tilde{\gamma}_5 \tilde{\gamma}_\mu, & \hat{S}_{4\mu} &= -\hat{S}_{\mu 4} = \gamma_\mu \\ \tilde{\gamma}_0 &= \gamma_0, & \tilde{\gamma}_5 &= \gamma_5, & \tilde{\gamma}_a &= i\gamma_a, \\ k, l, &= 0, 1, \dots, 5, & m &= 0, 1, 2, 3, & a &= 1, 2, 3 \end{aligned}$$

and the trivial identity transformation (corresponding to (2.4b) is omitted.

It is easy to verify that such defined operators  $R_{kl}$  commute with the operator  $L_0$  and so transform solutions of (2.1) into themselves.

Symmetries (2.6) satisfy the following commutation relations

$$[R_{kl}, R_{mn}] = 2i(\delta_{kn} R_{lm} + \delta_{lm} R_{kn} - \delta_{ln} R_{km} - \delta_{km} R_{ln}), \quad (2.7)$$

(by definition  $R_{ab} = -R_{ba}$ ).

In accordance with (2.7) symmetries (2.6) realize a representation of the algebra  $so(6)$ .

Let us now specify antilinear (i.e., including the complex conjugation) symmetries of equation (2.1) corresponding to reflections (2.3), (2.4). On the set of solutions of the Dirac equation they are reduced to the form

where  $R_{kl}$  are transformations (2.6) and  $C$  is the charge

$$\psi(x) \rightarrow B_{kl} \psi(x) \equiv CR_{kl} \psi(x), \quad (2.8)$$

conjugation transformation

$$C\psi(x) = \gamma_2 C\psi(x) \equiv i\gamma_2 \psi^*(x). \quad (2.9)$$

Using the relations

$$\begin{aligned} [C, R_{\lambda a}] = \{C, R_{\lambda \sigma}\} = \{C, R_{ab}\} = 0, \\ a, b, c = 1, 2, 3, \quad \lambda, \sigma = 0, 4, 5, \end{aligned} \quad (2.10)$$

we conclude that among the transformations (2.7) there are six representatives which satisfy  $(B_{Ac})^2 = -I$  (for  $A, C = 0, 4, 5$  or  $A, C = 1, 2, 3$ ) and nine representatives which satisfy the condition  $(B_{Ac})^2 = I$ , where  $I$  is the identity operator. We have a special interest in such transformations (2.7) whose square is positive (otherwise the corresponding  $B_{\mu\lambda}$  cannot be diagonalized to the real matrix  $\gamma_5$  and so they cannot be used for reductions considered in the following section). The corresponding symmetries are

$$B_{4a} = CR_{4a}, \quad (2.11a)$$

$$B_{5a} = CR_{5a}, \quad (2.11b)$$

$$B_{0a} = CR_{0a}. \quad (2.11c)$$

Using (2.7), (2.10) it is not difficult to specify commutation and anticommutation relations for operators (2.6) and (2.11). We notice that the set of operators  $\{\hat{R}_{kl} = iR_{kl}, B_{\alpha a}, C\}$  forms a basis of the 25-dimensional Lie algebra  $A_{25}$  characterized by commutation relations (2.7) and (2.12):

**Remark:** By including all symmetries (2.8) and products of

$$\begin{aligned}
[B_{\alpha a}, B_{\beta b}] &= -2(\delta_{ab}\hat{R}_{\alpha\beta} + \delta_{\alpha\beta}\hat{R}_{ab}), \\
[B_{\alpha a}, \hat{R}_{\beta\sigma}] &= 2(\delta_{\alpha\beta}B_{\sigma a} - \delta_{\alpha\sigma}B_{\beta a}), \\
[B_{\alpha a}, \hat{R}_{bc}] &= -2(\delta_{ac}B_{\alpha b} - \delta_{ab}B_{\alpha c}), \\
[B_{\alpha a}, \hat{R}_{\beta b}] &= \epsilon_{a\alpha b\beta c\sigma}B_{\sigma c} - 2\delta_{ab}\delta_{\alpha\beta}C, \\
[C, B_{\alpha a}] &= 2\hat{R}_{\alpha a}, \quad [C, \hat{R}_{\alpha a}] = 2B_{\alpha a}, \\
[C, \hat{R}_{ab}] &= [C, \hat{R}_{\alpha\beta}] = 0, \\
a, b, c &= 1, 2, 3, \quad \alpha, \beta, \sigma = 0, 4, 5.
\end{aligned} \tag{2.12}$$

symmetries (2.6), (2.8) and operator of multiplication by  $i=\sqrt{-1}$  the algebra  $A_{25}$  can be extended to the 64-dimensional Lie algebra defined over the field of real numbers. Additional extensions can be made by including the non-Lie involutive symmetries [8, 11].

Thus involutive symmetries of the Dirac equation generate the extended Lie algebra  $A_{25}$ . In the following sections we use it to reduce the Dirac equation for a charged particle interacting with various external fields and to search for supersymmetries of the Dirac equation.

### III. REDUCTION OF THE DIRAC EQUATION

Now we shall apply the results of the previous subsection to reduce the Dirac equation for a charged particle in an external field

$$L\Psi \equiv (\gamma^\mu \pi_\mu - m)\Psi = 0, \tag{3.1}$$

where  $\pi_\mu = p_\mu - eA_\mu$ ,  $p_\mu = i\partial/\partial x^\mu$ ,  $A_\mu = A_\mu(x) = A_\mu(x_0, \mathbf{x})$  is the vector-potential.

Equation (3.1) is invariant under one of the transformations described in (2.4a), (2.3a-c) provided vector-potential  $A_\mu$  satisfies one of the relations

$$A_\mu(-x_0, -\mathbf{x}) = -A_\mu(x_0, \mathbf{x}), \quad (3.2)$$

$$A_\mu(\theta_\lambda X) = (1 - 2\delta_{\mu\lambda}) A_\mu(X) \quad (3.3a)$$

$$A_\mu(\theta_{\lambda\sigma} X) = (1 - 2\delta_{\mu\lambda} - 2\delta_{\mu\sigma}) A_\mu(X) \quad (3.3b)$$

$$A_\mu(\theta'_\lambda X) = (2\delta_{\mu\lambda} - 1) A_\mu(X) \quad (3.3c)$$

respectively with  $\lambda$  and  $\sigma$  being fixed. On the other hand, if we require that (3.1) admits one of the symmetries (2.11a-c), vector-potential  $A_\mu$  has to satisfy one of the corresponding relations

$$A_\mu(\theta'_a X) = -(2\delta_{\mu a} - 1) A_\mu(X), \quad (3.4a)$$

$$A_\mu(\theta_a X) = -(1 - 2\delta_{\mu a}) A_\mu(X), \quad (3.4b)$$

$$A_\mu(\theta_{0a} X) = -(1 - 2\delta_{\mu 0} - 2\delta_{\mu a}) A_\mu(X) \quad (3.4c)$$

respectively.

We notice that relations (3.3) and (3.4) leave the Lorentz gauge  $\partial_\mu A^\mu = 0$  invariant.

To reduce (3.1) we diagonalize the corresponding symmetries (2.6). Let us consider in detail the case (3.2), i.e. when equation (3.1) is invariant under the transformation



$$\hat{R}\psi(x) = \gamma_5 \hat{\theta} \psi(x) = \gamma_5 \psi(-x). \quad (3.5)$$

To diagonalize this symmetry we use the operator

$$W = \frac{1}{\sqrt{2}}(1 + \gamma_5 \gamma_0) \frac{1}{\sqrt{2}}(1 + \gamma_5 \gamma_0 \hat{\theta}) = \hat{\theta}_+ + \gamma_0 \gamma_5 \hat{\theta}_- \quad (3.6)$$

with  $\hat{\theta}_\pm = \frac{1}{2}(1 \pm \hat{\theta})$ , then

$$W \hat{R} W^\dagger = W \gamma_5 \hat{\theta} W^\dagger = \gamma_5 \quad (3.7)$$

Simultaneously operator  $L$  of (3.1) is reduced to the block diagonal form:

$$W L W^\dagger = L' = -\gamma_5 \pi_0 - \frac{i}{2} \epsilon_{abc} \gamma_a \gamma_b \pi_c \hat{\theta} - m. \quad (3.8)$$

Thus the transformed equation

$$L' \psi' = 0, \quad \psi' = U \psi \quad (3.9)$$

has the desired reduced form

$$(-\mu \pi_0 - \sigma \cdot \pi \hat{\theta} - m) \psi'_\mu = 0, \quad \mu = \pm 1, \quad (3.10)$$

where  $\psi'_\mu$  are two-component spinors, i.e., non-zero components of eigenvectors of  $\gamma_5$  satisfying  $\gamma_5 \psi' = \mu \psi'$ .

For  $A_\mu = 0$  equation (3.10) is equivalent to the one considered in [17].

If equations (3.10) again admit a discrete symmetry, say

$$\psi_\mu(x) \rightarrow \hat{R} \psi_\mu = \sigma_3 \hat{\theta}_{12} \psi_\mu(x) = \sigma_3 \psi_\mu(x'), \quad x' = (\hat{\theta}_{12} x) \quad (3.11)$$

(which is the case if  $A_0(x') = A_0(x)$ ,  $A_1(x') = -A_1(x)$ ,  $A_2(x') = -A_2(x)$ ,  $A_3(x') = A_3(x)$ ), then they can further be reduced to one-component uncoupled subsystems. Indeed, by diagonalizing symmetry

$\hat{K}=\sigma_3\hat{\theta}_{12}$  and using transformation  $\psi'_\mu\rightarrow\psi''_\mu=W\psi'_\mu$  (the corresponding transformation operator is  $W=[1-\hat{\theta}_{12}-i\sigma_2(1+\hat{\theta}_{12})]/2$ ,  $W^{-1}=W^\dagger$ ) we change equation (3.10) to the following one

$$[-\mu\pi_0-\lambda\pi_1-\hat{\theta}_{12}(\pi_3+i\pi_2)-m]\psi_{\mu\lambda}=0, \quad (3.12)$$

where both  $\mu$  and  $\lambda$  runs independently over the values  $+, -$  and  $\psi_{\mu\lambda}$  are one-component functions, i.e., nonzero components of eigenvectors of matrix  $\sigma_3$  satisfying  $\sigma_3\psi''_\mu=\lambda\psi''_\mu$ .

We notice that transformations (3.6), (3.7), (3.8) can be used also for reduction of Dirac's equation with the anomalous (Pauli) interaction:

$$\left(\gamma^\mu\pi_\mu-m-\frac{ke}{2m}S_{\mu\nu}F^{\mu\nu}\right)\psi=0, \quad (3.13)$$

where

$$S_{\mu\nu}=\frac{i}{4}[\gamma_\mu, \gamma_\nu], \quad F_{\mu\nu}=i[\pi_\mu, \pi_\nu].$$

For example, if  $A_\mu$  satisfies (2.4a), then (3.13) is reduced to the following two subsystems for two-component spinors:

$$\left(-\mu\pi_0-\boldsymbol{\sigma}\cdot\boldsymbol{\pi}\hat{\theta}+\frac{ek}{m}\boldsymbol{\sigma}\cdot\boldsymbol{H}-\mu\frac{iek}{m}\boldsymbol{\sigma}\cdot\boldsymbol{E}\hat{\theta}-m\right)\psi_\mu=0, \quad \mu=\pm 1,$$

where  $\boldsymbol{H}$  and  $\boldsymbol{E}$  are vectors of the magnetic and electric field strengths:  $E_a=F_{0a}$ ,  $H_a=\epsilon_{abc}F_{bc}/2$ .

In an analogous way it is possible to reduce the Dirac equation (3.1) if vector-potentials satisfy one of relations (3.3) or (3.4). We present the complete list of the corresponding reductions in the Appendix.

## VI. REDUCTION OF THE DIRAC OSCILLATOR

The Dirac oscillator equation [14,15] can be written as

$$(\gamma^\mu P_\mu - i\omega \gamma_0 \gamma_a X_a - m)\psi = 0 \quad (4.1)$$

This equation is  $P$ -invariant, i.e., admits the following involutive symmetry

$$x_0 \rightarrow x'_0 = x_0, \quad \mathbf{x} \rightarrow \mathbf{x}' = -\mathbf{x}, \quad \psi(x_0, \mathbf{x}) \rightarrow \gamma_0 \psi(x_0, -\mathbf{x})$$

and consequently can be reduced to two uncoupled subsystems. Indeed, using the transformation  $\psi \rightarrow \psi' = W\psi$  where  $W = \frac{1}{\sqrt{2}}(1 - i\gamma_5 R_{40})$ , equation (4.1) decouples and can be expressed as

$$P_0 \psi_\pm = [\pm(\boldsymbol{\sigma} \cdot \mathbf{p} + m\hat{\theta}'_0) + i\omega \boldsymbol{\sigma} \cdot \mathbf{x} \hat{\theta}'_0] \psi_\pm. \quad (4.2)$$

Equations (4.2) admit involutive symmetry (3.9) and by means of operator  $W$  for (3.9) can be reduced to four one-component uncoupled equations.

However, there is another involutive symmetry for equation (4.2) which can be written as

$$Q = B\hat{\theta}'_0, \quad Q^2 \equiv 1, \quad (4.3)$$

where  $B$  is the Biedenharn operator [18]:

$$B = \frac{q}{|q|}, \quad q = \boldsymbol{\sigma} \cdot \mathbf{L} + 1, \quad \mathbf{L} = \mathbf{x} \times \mathbf{p}. \quad (4.4)$$

Operator  $B$  anticommutes with  $\boldsymbol{\sigma} \cdot \mathbf{p}$  and  $\boldsymbol{\sigma} \cdot \mathbf{x}$ , thus  $Q$  introduced in (4.3) commutes with the operator in square brackets defined in (4.2). On the set of functions  $\psi_p^\mu$  satisfying

$$Q\Psi_\rho^\mu = \mu\Psi_\rho^\mu, \quad \rho = \pm, \quad \mu = \pm 1 \quad (4.5)$$

equations (4.2) are reduced to the form

$$P_0\Psi_\rho^\mu = (\boldsymbol{\rho}\boldsymbol{\sigma}\mathbf{p} + \mu\rho mB + i\mu\boldsymbol{\sigma}\cdot\mathbf{x}B)\Psi_\rho^\mu. \quad (4.6)$$

In other words, the Dirac oscillator equation is reduced to four uncoupled two-component subsystems.

Setting  $m=0$  in (4.2) and (4.6), we receive the equations which we shall call the Weyl oscillators. Analogously to the Dirac oscillator case they generate oscillator-like spectra and are related to the free (Weyl) equation by changing  $\mathbf{p} \rightarrow \mathbf{p} - i\omega\mathbf{x}\beta$  with  $\beta=P$  or  $\beta=B$  being an operator anticommuting with the differential part of the corresponding Hamiltonian in (4.2) or (4.6) respectively.

The Weyl oscillators will be studied in more detail in the next paper. It appears they have very interesting symmetries and supersymmetries which are preserved if we change  $\mathbf{x} \rightarrow \frac{\partial W}{\partial \mathbf{x}}$  in (4.6), where  $W(\mathbf{x})$  is an arbitrary even superpotential.

## V. EXTENDED SUPERSYMMETRIES

We say that an equation of motion is supersymmetric if it admits specific symmetries (supercharges)  $Q_a$ ,  $a=1,2$ , which form the Witten superalgebra [19] (we choose  $Q_a$  Hermitian):

$$\{Q_a, Q_b\} = 2\delta_{ab}\hat{H}, \quad [Q_a, \hat{H}] = 0 \quad (5.1)$$

$\hat{H}$  is the related Hamiltonian.

To search for SUSY we use the following anticommutative relations for involutive symmetries (2.6)

$$\{R_{kl}, R_{mn}\} = \epsilon_{klmngf} R_{gf} + 2\delta_{km}\delta_{ln}I - 2\delta_{lm}\delta_{kn}I. \quad (5.2)$$

We start with the Dirac equation (3.1) for a charged particle interacting with the time-independent magnetic field. The corresponding vector-potentials have the form

$$A_0 = 0, \quad A_a = A_a(\mathbf{x}). \quad (5.3)$$

Denoting  $\psi_{\pm} = \frac{1}{2}(1 \mp i\gamma_5)\psi$  we have

$$\psi = \psi_{+} + \psi_{-}, \quad \psi_{+} = \begin{pmatrix} \Phi \\ 0 \end{pmatrix}, \quad \psi_{-} = \begin{pmatrix} 0 \\ \xi \end{pmatrix} \quad (5.4)$$

where  $\Phi$  and  $\xi$  are two component spinors, and 0 is the two component zero column. Expressing  $\psi$  via  $\psi_{\pm}$  we come from (3.1), (5.3) to the following equations:

$$(p_0^2 - m^2)\Phi = \hat{H}\Phi, \quad \hat{H} = \boldsymbol{\pi}^2 - e\boldsymbol{\sigma}\cdot\mathbf{H} \quad (5.5)$$

$$\xi = \frac{1}{m}(p_0 - \boldsymbol{\sigma}\cdot\boldsymbol{\pi})\Phi. \quad (5.6)$$

We will search for the SUSY of equation (5.5). The corresponding symmetries for (3.1) can be found using relations (5.4), (5.6).

For the case of arbitrary vector-potential  $A(\mathbf{x})$  equation (5.5) admits the following symmetry operator (supercharge)

$$Q_1 = \boldsymbol{\sigma}\cdot\boldsymbol{\pi}, \quad (5.7)$$

which satisfies the relation  $Q_1^2 = \hat{H}$  and so commutes with the "Hamiltonian"  $\hat{H}$ .

To find additional supercharges we suppose that the vector-

potential  $A_a(\mathbf{x})$  satisfies one of the relations (3.3) where  $\mu, \sigma = 1, 2, 3$ . The corresponding equation (5.5) admits the following symmetries:

$$\Phi(t, \mathbf{x}) \rightarrow r_a \Phi(t, \mathbf{x}) \equiv \sigma_a \Phi(t, \theta_a \mathbf{x}), \quad (5.8a)$$

$$\Phi(t, \mathbf{x}) \rightarrow r_{ab} \Phi(t, \mathbf{x}) \equiv \varepsilon_{abc} \sigma_c \Phi(t, \theta_{ab} \mathbf{x}), \quad (5.8b)$$

$$\Phi(t, \mathbf{x}) \rightarrow r \Phi(t, \mathbf{x}) \equiv \Phi(t, \theta \mathbf{x}). \quad (5.8c)$$

Operators (5.8) satisfy the following relations (compare with (5.2))

$$r_a^2 = r_{ab}^2 = r^2 = 1, \quad \{r_{ab}, r_b\} = 0, \quad \{r_a, r_b\} = 0, \quad a \neq b \quad (5.9)$$

(now sum over  $b$ ) and

$$\{r, \sigma \cdot \pi\} = \{r_a, \sigma \cdot \pi\} = 0, \quad [r_{ab}, \sigma \cdot \pi] = 0 \quad (5.10)$$

which enable us to construct the second supercharges

$$Q_2 = ir_a \sigma \cdot \pi \quad \text{and} \quad Q_2 = ir \sigma \cdot \pi \quad (5.11)$$

for the cases (3.3a) and (3.3c) respectively.

Thus the corresponding equation (3.3) admits N=2 SUSY.

If  $A_a(\mathbf{x})$  satisfy two or more relations (3.3) simultaneously, then equation (5.5) admits extended SUSY. All nonequivalent possibilities are listed in the following formulae:

$$\begin{cases} A_a(\theta_b \mathbf{x}) = (1 - 2\delta_{ab}) A_a(\mathbf{x}), \\ A_a(\theta_c \mathbf{x}) = (1 - 2\delta_{ac}) A_a(\mathbf{x}), \end{cases} \quad (5.12)$$

$$Q_1 = \sigma \cdot \pi, \quad Q_2 = ir_b \sigma \cdot \pi, \quad Q_3 = ir_c \sigma \cdot \pi, \quad c \neq b, \quad b, c = 1, 2, 3;$$

Relations (5.12) define three classes of vector-potentials  $A_\mu$  (corresponding to different fixed values of  $b, c$ ) which

$$\begin{cases} A_a(\theta_{12}\mathbf{x}) = (1-2\delta_{a1}-2\delta_{a2})A(\mathbf{x}), \\ A_a(\theta_{31}\mathbf{x}) = (1-2\delta_{a1}-2\delta_{a3})A(\mathbf{x}), \end{cases} \quad (5.13)$$

$$Q_1 = ir_{23}\sigma\pi, \quad Q_2 = ir_{31}\sigma\pi, \quad Q_3 = ir_{12}\sigma\pi;$$

$$\begin{cases} A_a(r_1\mathbf{x}) = (1-2\delta_{a1})A_a(\mathbf{x}), \\ A_a(r_2\mathbf{x}) = (1-2\delta_{2a})A_a(\mathbf{x}), \\ A_a(r_3\mathbf{x}) = (1-2\delta_{3a})A_a(\mathbf{x}), \end{cases} \quad (5.14)$$

$$Q_1 = ir_1\sigma\pi, \quad Q_2 = ir_2\sigma\pi, \quad Q_3 = ir_3\sigma\pi, \quad Q_4 = \sigma\pi.$$

generate SUSY. Operators  $Q_a$  (5.12), (5.13) realize N=3 extended SUSY, while the corresponding operators (5.14) realize N=4 extended SUSY.

It is necessary to note that the extended SUSY found in the above does not have direct connections with SUSY quantum field theory and cannot be used for a nontrivial extension of the Poincare group. But the symmetries (5.12)-(5.14) have rather nontrivial consequences in the quantum mechanical context, which consists of the specific degeneration of the corresponding energy spectra. Indeed, calculating *commutation* relations for the supercharges (5.12), we obtain

$$[Q_1, Q_2] = -ir_b\hat{H}, \quad [Q_2, Q_3] = -ir_{bc}\hat{H}, \quad [Q_3, Q_1] = ir_c\hat{H}. \quad (5.15)$$

It follows from (5.9), (5.15) that on the subspace of solutions of equation (5.5), corresponding to the nonzero eigenvalue  $E$  of the Hamiltonian  $\hat{H}$ , the operators

$$\hat{R}_{4k} = \frac{1}{\sqrt{E}}Q_k, \quad \hat{R}_{kl} = \frac{i}{2|E|}[Q_k, Q_l] \quad (5.16)$$

satisfy relations (2.7) and (5.17):

$$\hat{R}_{kl}^2 = 1, \quad \frac{1}{4!} \epsilon_{klmn} \hat{R}_{kl} \hat{R}_{mn} = 3 r_{ab} \quad (5.17)$$

Eigenvalues of  $r_{ab}$  are equal to  $\pm 1$ , and so operators (5.16) realize the representation  $D\left(\frac{1}{2}, \frac{1}{2}\right) \oplus D\left(\frac{1}{2}, -\frac{1}{2}\right)$  of the algebra  $so(4)$ .

In an analogous way, choosing the basis (5.16), we conclude that the supercharges (5.13) generate the same representation of the algebra  $so(4)$ , but the supercharges (5.14) generate the representation  $D\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right)$  of the algebra  $so(5)$  (in the last case we have in (5.16)  $k, l = 1, 2, 3, 5$ ). The corresponding commutation relations for operators (5.16) again can be expressed in the form (2.7).

Thus, for any symmetry (5.12) - (5.14) each nonzero eigenvalue of the Hamiltonian  $\hat{H}$  has four-fold degeneracy due to the hidden symmetry  $so(4)$  or  $so(5)$ .

N=2 and N=1 SUSY aspects of the Dirac equation were discussed by a number of authors, refer, e.g., to papers [20], surveys [12, 21] and monograph [22]. We extend the list of problems generating this symmetry and find a class of potentials generating extended SUSY.

## VI. REDUCTION TECHNIQUE IN SUPERSYMMETRIC QUANTUM MECHANICS

The idea of diagonalizing a discrete symmetry in order to reduce the corresponding equation of motion can be applied to many problems in mathematical physics. Continuing the theme of SUSY, we apply this idea to one-dimensional SUSY quantum mechanics [19]. The corresponding equation of motion has the form



$$i \frac{\partial}{\partial x_0} \psi = H \psi, \quad (6.1)$$

where  $H$  is the Hamiltonian with matrix potential

$$H = \frac{1}{2} (\hat{p}^2 + W^2 + W' \sigma_3), \quad \hat{p} = -i \frac{\partial}{\partial x}. \quad (6.2)$$

Equation (6.1) admits specific symmetries (supercharges) of the form

$$Q = \frac{1}{2\sqrt{2}} (\sigma_2 + i\sigma_1) (\hat{p} + iW), \quad \bar{Q} = \frac{1}{2\sqrt{2}} (\sigma_2 - i\sigma_1) (\hat{p} - iW) \quad (6.3)$$

which transform solutions into themselves and generate the following superalgebra (which is isomorphic to (5.1)):

$$\begin{aligned} Q^2 = \bar{Q}^2 = 0, \quad Q\bar{Q} + \bar{Q}Q = H, \\ [Q, H] = [\bar{Q}, H] = 0. \end{aligned} \quad (6.4)$$

Let us demonstrate that this superalgebra is reducible for odd superpotentials  $W(x)$ , i.e., for which  $W(-x) = -W(x)$ . Indeed, for  $W$  odd there exists the invariant operator

$$K = \sigma_3 p \quad (6.5)$$

( $p\psi(x) = \psi(-x)$ ) which commutes with supercharges  $Q$  and  $\bar{Q}$ . In order to diagonalize  $K$  we apply the operator

$$U = p_+ - i\sigma_2 p_-, \quad p_{\pm} = \frac{1}{2} (1 \pm p), \quad (6.6)$$

so that

$$UKU^\dagger = \sigma_3. \quad (6.7)$$

The corresponding supercharges are transformed into the diagonal form

$$\begin{aligned}
UQU^\dagger = Q' &= \frac{i}{2\sqrt{2}} [(1-\sigma_3) p_+ - (1+\sigma_3) p_-] (\hat{p} + iW), \\
U\bar{Q}U^\dagger = \bar{Q}' &= -\frac{i}{2\sqrt{2}} [(1-\sigma_3) p_- - (1+\sigma_3) p_+] (\hat{p} - iW),
\end{aligned}$$

i.e.,

$$Q' = \begin{pmatrix} Q_+ & 0 \\ 0 & Q_- \end{pmatrix}, \quad \bar{Q}' = \begin{pmatrix} \bar{Q}_+ & 0 \\ 0 & \bar{Q}_- \end{pmatrix}, \quad (6.8)$$

where

$$Q_+ = -\frac{i}{\sqrt{2}} (\hat{p} + iW) p_+, \quad \bar{Q}_+ = \frac{i}{\sqrt{2}} (\hat{p} - iW) p_-, \quad (6.9a)$$

$$Q_- = \frac{i}{\sqrt{2}} (\hat{p} + iW) p_-, \quad \bar{Q}_- = -\frac{i}{\sqrt{2}} (\hat{p} - iW) p_+. \quad (6.9b)$$

Thus supercharges (6.3) generate a reducible representation of the algebra (6.4) which is equivalent to a direct sum of representations (6.9a) and (6.9b). The corresponding Hamiltonians are of the form

$$H_+ = \frac{1}{2} (\hat{p}^2 + W^2 + W'p) \quad (6.10a)$$

and

$$H_- = \frac{1}{2} (\hat{p}^2 + W^2 - W'p). \quad (6.10b)$$

Operators (6.9a) and (6.10a) (as well as (6.9b) and (6.10b)) form a one-dimensional realization of SSQM, which has a very unique property: supercharges  $Q_+$  and  $\bar{Q}_+$  are not products of

commutative bosonic and fermionic operators. As a consequence of this fact the spectra of superhamiltonians with familiar potentials differ from the corresponding spectra in standard realization of SSQM. For instance, if  $W=\omega x$  then supercharges (6.9b) and Hamiltonian (6.10b) correspond to a specific version of the supersymmetric oscillator for which differences between eigenvalues are not equal to  $\omega$  (compare with paper [19]) but to  $2\omega$ , whereas supercharges (6.9a) and Hamiltonian (6.10a) present a supersymmetric system with spontaneously broken supersymmetry (i.e., with a degenerated ground state).

In conclusion we notice that the N=2 Wess-Zumino SSQM [23] with a superpotential being an odd complex function is completely reducible too.

## VII. CONCLUSION

We present the extended Lie algebra formed by involutive symmetries of the Dirac equation and apply it to reduction of a number of problems connected with interaction of a spin-1/2 particle with an external field.

Such a reduction technique can be generalized for reduction of systems of ordinary differential equations as well as many other systems of partial differential equations, including nonlinear ones. We plan to outline the results of our investigations of these possibilities elsewhere.

The other interesting application of involutive symmetries is searching for exact SUSY for the Dirac equation. We demonstrate that in addition to the known class of systems with

N=2 SUSY this equation generates also extended supersymmetries. Moreover, the list of supersymmetric problems can be extended by including antilinear involutive symmetries.

The other goal of the present article is to demonstrate that a wide class of realizations of SSQM is completely reducible. We obtain a one-dimensional representations of the Witten superalgebra (6.4) which can be extended to the case of multidimensional space of independent variables.

Finally we note that the Weyl oscillators of Section V are the simplest consistent examples of generalizations of the Dirac oscillator [14,15]. Such generalizations for the cases of arbitrary spin multi-body systems are intensively discussed in literature [24, 25].

## APPENDIX

### EXPLICIT FORM OF REDUCTIONS

Here we present in explicit form reductions of the Dirac equation which are possible if vector-potential  $A_\mu$  satisfies one of the relations (3.3a-c) or (3.4a-c). To reduce (3.1) it is sufficient to diagonalize the corresponding symmetries (2.6) or (2.8). This can be done by using the operators

$$W_{4v} = \frac{1}{\sqrt{2}}(1 - i\gamma_5 R_{4v}), \quad v=0, 1, 2, 3, \quad (\text{A1.a})$$

$$W_{va} = \frac{1}{2}(1 + i\gamma_5 \gamma_a)(1 + \gamma_a R_{va}), \quad a=1, 2, 3, \quad (\text{A1.b})$$

$$W_{5v} = \frac{1}{\sqrt{2}}(1 - i\gamma_5 R_{5v}) \quad (\text{A1.c})$$

or

$$\begin{aligned} \bar{W}_{4a'} &= \frac{1}{2}(1 + i\gamma_5 \gamma_2)(1 - i\gamma_2 B_{4a'}), \quad a'=1, 3, \\ \bar{W}_{42} &= \frac{1}{2}(1 + i\gamma_5 \gamma_0 \theta'_2 c)(1 + \gamma_0), \end{aligned} \quad (\text{A2.a})$$

$$\bar{W}_{5a} = \frac{1}{2}(1 + \gamma_4 \gamma_0)(1 + \gamma_0 B_{5a}), \quad a=1, 2, 3, \quad (\text{A2.b})$$

$$\bar{W}_{0a} = \frac{1}{\sqrt{2}}(1 + \gamma_5 B_{0a}). \quad (\text{A2.c})$$

In all the cases we have  $W_{AB} R_{AB} W_{AB}^\dagger = i\gamma_5$  or  $\bar{W}_{AB} B_{AB} \bar{W}_{AB}^\dagger = \gamma_5$ . The transformations  $\psi \rightarrow W_{AB} \psi$  reduce equation (3.1) to uncoupled subsystems of the following form:

$$\begin{aligned} [\pi_0 \mp (\boldsymbol{\sigma} \cdot \boldsymbol{\pi} - im\boldsymbol{\theta}_0)] \psi_\pm &= 0, \\ [\pi_0 \mp (\boldsymbol{\sigma} \cdot \boldsymbol{\pi} - m\boldsymbol{\theta}_a \sigma_a)] \psi_\pm &= 0; \end{aligned} \quad (\text{A3.a})$$

$$\begin{aligned} [\sigma_a (\mp \pi_0 + m) \boldsymbol{\theta}_{0a} + \boldsymbol{\sigma} \cdot \boldsymbol{\pi}] \psi_\pm &= 0, \\ [\pi_0 \mp \boldsymbol{\sigma} \cdot \boldsymbol{\pi} - \sigma_b m + (i\sigma_a \boldsymbol{\theta}_{ab} \pm \sigma_b) \pi_b] \psi_\pm &= 0; \end{aligned} \quad (\text{A3.b})$$

$$\begin{aligned} [\pi_0 \mp (\boldsymbol{\sigma} \cdot \boldsymbol{\pi} + m\boldsymbol{\theta}'_0)] \psi_\pm &= 0, \\ [\pi_0 \mp \boldsymbol{\sigma} \cdot \boldsymbol{\pi} + im\boldsymbol{\theta}'_a \sigma_a] \psi_\pm &= 0 \end{aligned} \quad (\text{A3.c})$$

or

(no sums over repeated indices). Here  $\psi_\pm$  are two-component wave functions, i.e., nonzero components of eigenvectors of the matrix

$$\begin{aligned} (\hat{\theta}'_2 c\pi_0 - \boldsymbol{\sigma} \cdot \boldsymbol{\pi} + im)\psi_{\pm} &= 0, \\ (\mp\pi_0 - \boldsymbol{\sigma} \cdot \boldsymbol{\pi} + \pi_{a'}(\boldsymbol{\sigma}_{a'} \mp ic\hat{\theta}'_{a'}) - c\hat{\theta}'_{a'}m)\psi_{\pm} &= 0, \quad a'=1,3 \end{aligned} \quad (\text{A4.a})$$

$$\begin{aligned} (\pm\pi_0 + \hat{\theta}_{a'}\boldsymbol{\sigma}_{a'} \cdot \boldsymbol{\pi} - m)\psi_{\pm} &= 0, \\ (\pm\pi_0 + c\hat{\theta}_2 \boldsymbol{\sigma} \cdot \boldsymbol{\pi} - m)\psi_{\pm} &= 0, \end{aligned} \quad (\text{A4.b})$$

$$\begin{aligned} (\pm\pi_0 - \boldsymbol{\sigma} \cdot \boldsymbol{\pi} - ic\hat{\theta}_{03} \boldsymbol{\sigma}_1 m)\psi_{\pm} &= 0, \\ (\pm\pi_0 - \boldsymbol{\sigma} \cdot \boldsymbol{\pi} - ic\hat{\theta}_{01} \boldsymbol{\sigma}_3 m)\psi_{\pm} &= 0, \\ (\pi_0 \mp \boldsymbol{\sigma} \cdot \boldsymbol{\pi} \mp c\hat{\theta}_{02} m)\psi_{\pm} &= 0. \end{aligned} \quad (\text{A4.c})$$

$\gamma_5$ , corresponding to the eigenvalues  $\pm 1$ .

Symmetry (4.8) commutes with operator  $L$  of (3.1) iff  $A_{\mu}=0$  and so can be used only for reduction of the free Dirac equation (2.1). The corresponding operator  $W = \frac{1}{\sqrt{2}}(1 + i\gamma_5 C)$  diagonalizes symmetry (2.9) to the form  $\gamma_5$  and reduces equation (2.1) to the following uncoupled subsystems:

$$(D_0 - \mu \boldsymbol{\sigma} \cdot \boldsymbol{p} + i\sigma_2 cm)\psi_{\mu} = 0, \quad \mu = \pm 1. \quad (\text{A.5})$$

Imposing condition (4.5) on solutions of the first equation (A3.a) and setting  $A_{\mu}=0$  we come to the equations proposed in [18].

If the vector-potential  $A_{\mu}$  has such parities that the corresponding Dirac equation (3.1) admits two commuting symmetries from the set (2.6), (2.8) then we can reduce (3.1) to four uncoupled subsystems. If we find such a pair  $(S_1, S_2)$ , then  $(S_1, S_1 S_2)$  and  $(S_2, S_1 S_2)$  are also sets of commuting symmetries equivalent to the set  $(S_1, S_2)$ . Using (2.7), (2.12), it is not difficult to write down the 51 nonequivalent pairs of commuting symmetries:

In other words, there are 51 possible reductions of the

$$\begin{aligned} & \{R_{0a}, R_{bc}\}, \{R_{4a}, R_{bc}\}, \{R_{0a}, R_{4b}\}, \{R_{04}, R_{ab}\}, \{B_{4a}, R_{0b}\}, \\ & \{B_{0a}, R_{5b}\}, \{B_{5a}, R_{4b}\}, \{B_{5a}, R_{ab}\}, \{B_{4a}, R_{ab}\}, \{B_{4a}, R_{54}\}, \\ & \{B_{0a}, R_{40}\}, \{B_{0a}, R_{50}\}, \{B_{0a}, R_{50}\}, \\ & a, b, c, = 1, 2, 3, \quad a \neq b, \quad a \neq c, \quad b \neq c. \end{aligned}$$

Dirac equation to four uncoupled subsystems by means of linear and antilinear involutive symmetries. The explicit form of these reductions can be found in analogy with the above.

We notice that only two of the involutive symmetries of the Dirac equation commute with any Lorentz transformation, namely  $R$  and  $C$  given in (3.5) and (2.9) respectively. Consequently the corresponding reduced equations (3.9) and (4.19) are Lorentz-invariant.

## REFERENCES

- [1] Vilenkin H Ya and Klimyk A U 1991-1993 Representations of Lie Groups and Special Functions, vol.1-3; 1995 Representations of Lie Groups and Special Functions. Recent Advances (Kluwer, Dordrecht); Miller U 1977 Symmetry and Separation of Variables (Addison-Wesley)
- [2] Gardner C et al 1967 Phys. Rev. Lett. **19** 109; Michelson J and Niederle J 1984 Lett.Math.Phys. **8**, 195
- [3] Olver P 1986 Application of Lie Groups to Differential Equations (N.Y.:Springer); Ibragimov N Ch 1983 Transformation Groups in Mathematical Physics (Nauka, Moscow)

- [4] Taflin E 1983 Pacific J. Math. **108**, 203; M. Flato and Simon J 1980 J.Math.Phys. **21**, 913; 1980 Phys.Lett. **94B**, 518 and references cited therein
- [5] Fushchich W I, Shtelen W M and Serov N I 1993 Symmetry Analysis and Exact Solutions of Equations of Nonlinear Mathematical Physics (Dordrecht: D.Reidel)
- [6] Konopelchenko B G 1987 Nonlinear Integrable Equations (Lecture Notes in Physics **270**, Springer-Verlag, Berlin-Heidelberg)
- [7] Limic N, Niederle J and Raczka R 1966 J. Math. Phys. **7**, 1861, 2026; 1967 *ibid* **8**, 1079
- [8] Fushchich W I and Nikitin A G 1994 Symmetries of Equations of Quantum Mechanics (N.Y.: Allerton Press Inc)
- [9] Plebanski J F and Pursey D L 1960 Bull. Am. Phys. Soc. **5**, 81; Pursey D L and Plebanski J F 1984 Phys. Rev. D **29**, 1848; Gürsey Feza 1958 Nuovo Cimento **8**, 411
- [10] Fushchich W I 1971 Teor. Mat. Fiz. **7**, 3
- [11] Fushchich W I and Nikitin A G Lett. Nuovo Cimento 1977 **19**, 347
- [12] Gendenstein L E and Krive I V 1985 Usp.Fiz.Nauk **146**, 553
- [13] Cooper F, Khare A and Sukhamtme U 1995 Phys. Rep. **211**, 268
- [14] Itô D, Mori K and Carreri E 1967 Nuovo Cim. **A 51**, 119; Cook P A 1971 Lett.Nuovo.Cim. **1**, 419
- [15] Moshinsky M and Szczepaniak A 1989 J.Phys. **A 22**, L817
- [16] Wigner E P 1964 Lect.Istanbul School of Theor.Phys. (Gordon and Breach)
- [17] Srivastava T 1984 Nuovo Cim. **A 80**, 113, 131
- [18] Biedenharn L C and Han M Y 1971 Phys.Lett. **B 36**, 475



- [19] Witten E 1981 Nucl.Phys. **188**, 513
- [20] Quesne C 1991 Int.J.Mod.Phys. **A 6**, 1567; Cooper F, Khare A, Musto R, and Wipf A 1988 Ann.Phys. **187**, 1; Nogamy Y and Toyama F 1993 Phys.Rev. **A 47** 1708
- [21] Cooper F, Khare A, and Sukhatme U 1995 Phys.Rep. **251** 267
- [22] Thaller B 1992 The Dirac Equation (Springer-Verlag)
- [23] Arai A 1989 J.Math.Phys. **30**, 2973
- [24] Moshinsky M, Loyola G and Villegas C J. 1991 Math. Phys. **32** 373; Del Sol Mesa A and Moshinsky M 1994 J.Phys. **A 27**, 4685
- [25] Beckers J, Debergh N and Nikitin A G 1992 J. Math. Phys. **33**, 3387; Debergh N, Ndimubandi J and Strivay D 1992 Z. Phys. C. **56**, 421; Nedjadji M and Barrett R C 1994 J.Phys. **A 27** 4301; Dvoyeglazov V V 1994 Nuovo Cim. **A 107** 1785