

Revisiting the Symmetries of the Quantum Smorodinsky–Winternitz System in D Dimensions^{*}

Christiane QUESNE

Physique Nucléaire Théorique et Physique Mathématique, Université Libre de Bruxelles, Campus de la Plaine CP229, Boulevard du Triomphe, B-1050 Brussels, Belgium

E-mail: cquesne@ulb.ac.be

Received January 17, 2011, in final form March 25, 2011; Published online April 02, 2011

doi:[10.3842/SIGMA.2011.035](https://doi.org/10.3842/SIGMA.2011.035)

Abstract. The D -dimensional Smorodinsky–Winternitz system, proposed some years ago by Evans, is re-examined from an algebraic viewpoint. It is shown to possess a potential algebra, as well as a dynamical potential one, in addition to its known symmetry and dynamical algebras. The first two are obtained in hyperspherical coordinates by introducing D auxiliary continuous variables and by reducing a $2D$ -dimensional harmonic oscillator Hamiltonian. The $\mathfrak{su}(2D)$ symmetry and $\mathfrak{w}(2D) \oplus_s \mathfrak{sp}(4D, \mathbb{R})$ dynamical algebras of this Hamiltonian are then transformed into the searched for potential and dynamical potential algebras of the Smorodinsky–Winternitz system. The action of generators on wavefunctions is given in explicit form for $D = 2$.

Key words: Schrödinger equation; superintegrability; potential algebras; dynamical potential algebras

2010 Mathematics Subject Classification: 20C35; 81R05; 81R12

1 Introduction

In classical mechanics, a Hamiltonian H with D degrees of freedom is said to be completely integrable if it allows D integrals of motion X_μ , $\mu = 1, 2, \dots, D$, that are well-defined functions on phase space, are in involution and are functionally independent (see, e.g., [1]). These include the Hamiltonian, so that we may assume $X_D = H$. The system is superintegrable if there exist k additional integrals of motion Y_ν , $\nu = 1, 2, \dots, k$, $1 \leq k \leq D - 1$, that are also well-defined functions on phase space and are such that the integrals $H, X_1, X_2, \dots, X_{D-1}, Y_1, Y_2, \dots, Y_k$ are functionally independent. The cases $k = 1$ and $k = D - 1$ correspond to minimal and maximal superintegrability, respectively.

Similar definitions apply in quantum mechanics with Poisson brackets replaced by commutators, but H, X_μ , and Y_ν must now be well-defined operators forming an algebraically independent set. Maximally superintegrable quantum systems appear in many domains of physics, such as condensed matter as well as atomic, molecular, and nuclear physics. They have a lot of nice properties: they can be exactly (or quasi-exactly) solved, they are often separable in several coordinate systems and their spectrum presents some “accidental” degeneracies, i.e., degeneracies that do not follow from the geometrical symmetries of the problem.

The most familiar examples of such systems are the Kepler–Coulomb [2, 3, 4] and the oscillator [5, 6] ones. Other well-known instances are those resulting from the first systematic search for superintegrable Hamiltonians on E_2 carried out by Smorodinsky, Winternitz, and collaborators [7, 8, 9] and from its continuation by Evans on E_3 [10]. These studies were restricted to those cases where the integrals of motion are first- or second-order polynomials in the momenta. Later

^{*}This paper is a contribution to the Special Issue “Symmetry, Separation, Super-integrability and Special Functions (S⁴)”. The full collection is available at <http://www.emis.de/journals/SIGMA/S4.html>

on, many efforts have been devoted to arriving at a complete classification of these so-called second-order superintegrable systems (see, e.g., [11, 12, 13, 14, 15, 16, 17]).

Only recently, the pioneering work of Drach [18, 19] on two-dimensional Hamiltonian systems with third-order integrals of motion has been continued [20, 21]. Nowadays the search for D -dimensional superintegrable systems with higher-order integrals of motion has become a very active field of research (see, e.g., [22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32]).

In the present paper, we plan to re-examine from an algebraic viewpoint one of the classical examples of D -dimensional superintegrable quantum systems, namely the Smorodinsky–Winternitz (SW) one [7, 8, 9, 10, 33, 34], which may be defined in Cartesian coordinates as

$$H^{(\mathbf{k})} = \sum_{\mu=1}^D \left(-\partial_{x_\mu}^2 + \frac{k_\mu^2}{x_\mu^2} + \omega^2 x_\mu^2 \right). \quad (1.1)$$

Here $\omega, k_1, k_2, \dots, k_D$ are some constants, which we assume to be real and positive.

Several distinct algebraic methods may be used in connection with superintegrable systems. One of them is based on the fact that the integrals of motion generate a nonlinear algebra closing at some order [35, 36, 37]. It has been shown, for instance, that for two-dimensional second-order superintegrable systems with nondegenerate potential and the corresponding three-dimensional conformally flat systems, one gets a quadratic algebra closing at order 6 [11, 12, 13, 14, 15]. Its finite-dimensional unitary representations can be determined [38] by using a deformed parafermion oscillator realization [39, 40], thereby allowing a calculation of the energy spectrum. This procedure can be extended to higher-order integrals of motion and to the corresponding higher-degree nonlinear algebras [25, 26].

Superintegrable systems may also be related [25, 26] to systems studied in supersymmetric quantum mechanics [41, 42] or higher-order supersymmetric quantum mechanics [43, 44, 45, 46, 47, 48, 49, 50, 51], hence can be described in terms of either linear or nonlinear superalgebras. As a consequence, supersymmetry provides a convenient tool for generating superintegrable quantum systems with higher-order integrals of motion [52, 53].

The concept of exact or quasi-exact solvability [54, 55, 56], based on the existence of an infinite flag of functionally linear spaces preserved by the Hamiltonian or only that of one of these spaces, appears to be related to finite-dimensional representations of some Lie algebras of first-order differential operators, such as $\mathfrak{sl}(2, \mathbb{R})$, $\mathfrak{sl}(3, \mathbb{R})$, etc. Although different from the concept of superintegrability, it can be related to the latter for some superintegrable systems (see, e.g., [27, 49, 57]). It is worth noting, however, that some alternative definitions of exact and quasi-exact solvability have been proposed for some specific superintegrable systems in connection with multiseparability of the corresponding Schrödinger equation [58, 59].

The accidental degeneracies appearing in the bound-state spectrum of superintegrable quantum systems may be understood in terms of a symmetry algebra, which is such that for any energy level the wavefunctions corresponding to degenerate states span the carrier space of one of its unitary irreducible representations (unirreps) [60, 61]. The generators of this symmetry algebra, commuting with the Hamiltonian, are integrals of motion, which may assume a rather complicated form in terms of some basic ones due to the fact that linear algebras are often preferred¹ (note, however, that nonlinear algebras may also be considered [62]). A familiar example of this phenomenon is provided by the $\mathfrak{so}(4)$ symmetry algebra of the three-dimensional Kepler–Coulomb problem [2, 3, 4]. Another one corresponds to the $\mathfrak{su}(3)$ symmetry algebra of the three-dimensional SW system [34] (or, in general, $\mathfrak{su}(D)$ for the D -dimensional one).

In some cases, the symmetry algebra can be enlarged to a spectrum generating algebra (also called dynamical algebra) by including some ladder operators, which are not integrals of motion

¹It is worth observing here that this may be seen as the obverse of the approach used in [11, 12, 13, 14, 15, 35, 36, 37, 38], where the generators are the basic integrals of motion but the algebra turns out to be nonlinear.

but act as raising or lowering operators on the bound-state wavefunctions in such a way that all of them carry a single unirrep of the algebra [63, 64, 65]. For the three-dimensional SW system, it has been shown [34] to be given by the semidirect sum Lie algebra $\mathfrak{w}(3) \oplus_s \mathfrak{sp}(6, \mathbb{R})$, where $\mathfrak{w}(3)$ denotes a Weyl algebra (or, in general, by $\mathfrak{w}(D) \oplus_s \mathfrak{sp}(2D, \mathbb{R})$ in D dimensions).

For one-dimensional systems, three other types of Lie algebraic approaches have been extensively studied. All of them rely on an embedding of the system into a higher-dimensional space by introducing some auxiliary continuous variables and on the subsequent reduction of the extended system to the initial one, a procedure also used in discussing superintegrability (see, e.g., [24]). They work for hierarchies of Hamiltonians, whose members correspond to the same potential but different quantized strengths. The simplest ones are the potential algebras [66, 67, 68], whose unirrep carrier spaces are spanned by wavefunctions with the same energy, but different potential strengths. Larger algebras, which also contain some generators connecting wavefunctions with different energies, are called dynamical potential algebras [69, 70, 71]². Finally, a third kind of algebras, termed satellite algebras [74, 75], have the property that there is a conserved quantity different from the energy.

Up to now, only the first one of these Lie algebraic approaches, namely that of potential algebras, has been applied to some D -dimensional superintegrable systems [76, 77, 78, 79, 80, 81, 82, 83].

The purpose of the present paper is threefold: first to apply this technique to the D -dimensional SW Hamiltonian (1.1), second to present for the same the first construction of a dynamical potential algebra in more than one dimension, and third to show very explicitly the action of both the potential and dynamical potential algebra generators on the wavefunctions in the two-dimensional case.

The paper is organized as follows. In Section 2, the solutions, as well as the symmetry and dynamical algebras, of a $2D$ -dimensional harmonic oscillator are obtained in a suitable orthogonal coordinate system. In Section 3, they are transformed into the solutions, as well as the potential and dynamical potential algebras, of the D -dimensional SW system in hyperspherical coordinates. The $D = 2$ case is then dealt with in detail in Section 4. Finally, Section 5 contains the conclusion.

2 $2D$ -dimensional harmonic oscillator

Let us consider a harmonic oscillator Hamiltonian

$$H^{\text{osc}} = \sum_{\mu=1}^{2D} \left(-\partial_{X_\mu}^2 + X_\mu^2 \right)$$

in a $2D$ -dimensional space, whose Cartesian coordinates are denoted by X_μ , $\mu = 1, 2, \dots, 2D$. For our purposes, it is convenient to consider it in a different orthogonal coordinate system, which we will now proceed to introduce.

2.1 Harmonic oscillator in variables $R, \theta_1, \theta_2, \dots, \theta_{D-1}, \lambda_1, \lambda_2, \dots, \lambda_D$

On making the change of variables

$$X_1 = R \sin \theta_1 \sin \theta_2 \cdots \sin \theta_{D-1} \sin \lambda_1, \quad X_2 = R \sin \theta_1 \sin \theta_2 \cdots \sin \theta_{D-1} \cos \lambda_1,$$

²Some authors prefer to use the terminology of dynamical algebra of the hierarchy instead of dynamical potential algebra and to employ discrete variables, related to the quantum numbers characterizing the system, instead of continuous auxiliary variables. In this way, they get discrete-differential realizations of the algebras [72]. Other authors favour the use of nonlinear superalgebras [73].

$$\begin{aligned}
X_{2\nu-1} &= R \sin \theta_1 \sin \theta_2 \cdots \sin \theta_{D-\nu} \cos \theta_{D-\nu+1} \sin \lambda_\nu, & \nu = 2, 3, \dots, D-1, \\
X_{2\nu} &= R \sin \theta_1 \sin \theta_2 \cdots \sin \theta_{D-\nu} \cos \theta_{D-\nu+1} \cos \lambda_\nu, & \nu = 2, 3, \dots, D-1, \\
X_{2D-1} &= R \cos \theta_1 \sin \lambda_D, & X_{2D} &= R \cos \theta_1 \cos \lambda_D,
\end{aligned} \tag{2.1}$$

where $0 \leq R < \infty$, $0 \leq \theta_\nu < \frac{\pi}{2}$, $\nu = 1, 2, \dots, D-1$, and $0 \leq \lambda_\nu < 2\pi$, $\nu = 1, 2, \dots, D$, H^{osc} can be rewritten as

$$\begin{aligned}
H^{\text{osc}} &= -\partial_R^2 - \frac{2D-1}{R} \partial_R - \frac{1}{R^2} \left\{ \partial_{\theta_1}^2 + [(2D-3) \cot \theta_1 - \tan \theta_1] \partial_{\theta_1} \right. \\
&\quad + \sum_{\nu=2}^{D-1} \frac{1}{\sin^2 \theta_1 \sin^2 \theta_2 \cdots \sin^2 \theta_{\nu-1}} \left[\partial_{\theta_\nu}^2 + [(2D-2\nu-1) \cot \theta_\nu - \tan \theta_\nu] \partial_{\theta_\nu} \right] \\
&\quad + \frac{1}{\sin^2 \theta_1 \sin^2 \theta_2 \cdots \sin^2 \theta_{D-1}} \partial_{\lambda_1}^2 + \sum_{\nu=2}^{D-1} \frac{1}{\sin^2 \theta_1 \sin^2 \theta_2 \cdots \sin^2 \theta_{D-\nu} \cos^2 \theta_{D-\nu+1}} \partial_{\lambda_\nu}^2 \\
&\quad \left. + \frac{1}{\cos^2 \theta_1} \partial_{\lambda_D}^2 \right\} + R^2
\end{aligned}$$

and is clearly separable.

In the corresponding Schrödinger equation

$$H^{\text{osc}} \Psi^{\text{osc}}(R, \boldsymbol{\theta}, \boldsymbol{\lambda}) = E^{\text{osc}} \Psi^{\text{osc}}(R, \boldsymbol{\theta}, \boldsymbol{\lambda}) \tag{2.2}$$

with $\boldsymbol{\theta} = \theta_1 \theta_2 \cdots \theta_{D-1}$ and $\boldsymbol{\lambda} = \lambda_1 \lambda_2 \cdots \lambda_D$, we may therefore write

$$\Psi^{\text{osc}}(R, \boldsymbol{\theta}, \boldsymbol{\lambda}) = \mathcal{N}^{\text{osc}} \mathcal{L}(z) \left(\prod_{\nu=1}^{D-1} \Theta_\nu(\theta_\nu) \right) \left(\prod_{\nu=1}^D e^{ip_{D-\nu+1} \lambda_\nu} \right), \quad z = R^2, \tag{2.3}$$

where

$$\partial_{\lambda_\nu}^2 \Psi^{\text{osc}}(R, \boldsymbol{\theta}, \boldsymbol{\lambda}) = -p_{D-\nu+1}^2 \Psi^{\text{osc}}(R, \boldsymbol{\theta}, \boldsymbol{\lambda})$$

and $p_1, p_2, \dots, p_D \in \mathbb{Z}$. The normalization constant \mathcal{N}^{osc} in (2.3) will be determined in such a way that

$$\int dV |\Psi^{\text{osc}}(R, \boldsymbol{\theta}, \boldsymbol{\lambda})|^2 = 1, \tag{2.4}$$

where

$$dV = \prod_{\mu=1}^{2D} dX_\mu = R^{2D-1} dR \left[\prod_{\nu=1}^{D-1} (\sin \theta_\nu)^{2D-2\nu-1} \cos \theta_\nu d\theta_\nu \right] \left(\prod_{\nu=1}^D d\lambda_\nu \right). \tag{2.5}$$

As shown in the appendix, the angular part of wavefunctions (2.3) can be written as

$$\Theta_{\mathbf{n}}^{(\mathbf{p})}(\boldsymbol{\theta}) = \prod_{\nu=1}^{D-1} \Theta_{n_\nu}^{(a_\nu, b_\nu)}(\theta_\nu), \quad \mathbf{n} = n_1 n_2 \cdots n_{D-1}, \quad \mathbf{p} = p_1 p_2 \cdots p_D, \tag{2.6}$$

$$\Theta_{n_\nu}^{(a_\nu, b_\nu)}(\theta_\nu) = (\cos \theta_\nu)^{a_\nu - \frac{1}{2}} (\sin \theta_\nu)^{b_\nu - \frac{1}{2}} P_{n_\nu}^{(a_\nu - \frac{1}{2}, b_\nu + D - \nu - \frac{3}{2})}(-\cos 2\theta_\nu), \tag{2.7}$$

where $n_1, n_2, \dots, n_{D-1} \in \mathbb{N}$,

$$a_\nu = |p_\nu| + \frac{1}{2}, \quad \nu = 1, 2, \dots, D-1,$$

$$\begin{aligned} b_\nu &= 2n_{\nu+1} + 2n_{\nu+2} + \cdots + 2n_{D-1} + |p_{\nu+1}| + |p_{\nu+2}| + \cdots + |p_D| + \frac{1}{2}, \quad \nu = 1, 2, \dots, D-2, \\ b_{D-1} &= |p_D| + \frac{1}{2}, \end{aligned} \quad (2.8)$$

and $P_{n_\nu}^{(a_\nu - \frac{1}{2}, b_\nu + D - \nu - \frac{3}{2})}(-\cos 2\theta_\nu)$ denotes a Jacobi polynomial [84], while the radial part can be expressed as

$$\mathcal{L}_{n_r}^{(j)}(z) = z^j L_{n_r}^{(2j+D-1)}(z) e^{-\frac{1}{2}z}, \quad (2.9)$$

in terms of a Laguerre polynomial [84]. Here $n_r \in \mathbb{N}$, while j is defined by

$$j = n_1 + n_2 + \cdots + n_{D-1} + \frac{1}{2}(|p_1| + |p_2| + \cdots + |p_D|) \quad (2.10)$$

and may take nonnegative integer or half-integer values.

The corresponding energy eigenvalues are given by

$$E_{n_r, j}^{\text{osc}} = 2(2n_r + 2j + D). \quad (2.11)$$

We therefore recover the well-known spectrum of the $2D$ -dimensional harmonic oscillator

$$E_N^{\text{osc}} = 2(N + D), \quad N = 2n_r + 2j = 0, 1, 2, \dots,$$

whose levels, completely characterized by N , have a degeneracy equal to $\binom{N+2D-1}{2D-1}$.

Finally, the normalization constant in equation (2.3) can be easily determined from some well-known properties of Laguerre and Jacobi polynomials [84] and is given by

$$\begin{aligned} \mathcal{N}_{n_r, \text{np}}^{\text{osc}} &= \left(\frac{n_r!}{\pi^D (n_r + 2j + D - 1)!} \right)^{1/2} \\ &\times \prod_{\nu=1}^{D-1} \left(\frac{n_\nu! (2n_\nu + a_\nu + b_\nu + D - \nu - 1) (n_\nu + a_\nu + b_\nu + D - \nu - 2)!}{(n_\nu + a_\nu - \frac{1}{2})! (n_\nu + b_\nu + D - \nu - \frac{3}{2})!} \right)^{1/2}. \end{aligned} \quad (2.12)$$

2.2 Harmonic oscillator symmetry and dynamical algebras

As it is well known [5, 6], to each of the oscillator levels specified by N there corresponds a symmetric unirrep $[N]$ of its $\text{su}(2D)$ symmetry algebra. The generators of the latter

$$\bar{E}_{\mu\nu} = E_{\mu\nu} - \frac{1}{2D} \delta_{\mu,\nu} \sum_{\rho} E_{\rho\rho}, \quad \mu, \nu = 1, 2, \dots, 2D,$$

with

$$[\bar{E}_{\mu\nu}, \bar{E}_{\mu'\nu'}] = \delta_{\nu,\mu'} \bar{E}_{\mu\nu'} - \delta_{\mu,\nu'} \bar{E}_{\mu'\nu}, \quad \bar{E}_{\mu\nu}^\dagger = \bar{E}_{\nu\mu},$$

are most easily constructed in terms of bosonic creation and annihilation operators

$$\alpha_\mu^\dagger = \frac{1}{\sqrt{2}} (X_\mu - \partial_{X_\mu}), \quad \alpha_\mu = \frac{1}{\sqrt{2}} (X_\mu + \partial_{X_\mu}), \quad \mu = 1, 2, \dots, 2D, \quad (2.13)$$

from

$$E_{\mu\nu} = \frac{1}{2} \{ \alpha_\mu^\dagger, \alpha_\nu \} = \alpha_\mu^\dagger \alpha_\nu + \frac{1}{2} \delta_{\mu,\nu}. \quad (2.14)$$

The harmonic oscillator Hamiltonian turns out to be proportional to the first-order Casimir operator \mathcal{C}_1 of $\text{u}(2D)$,

$$H^{\text{osc}} = 2\mathcal{C}_1 = 2 \sum_{\mu} E_{\mu\mu} = 2\mathcal{E}. \quad (2.15)$$

In the coordinates (2.1) chosen to describe the oscillator, the $\mathfrak{so}(2D)$ subalgebra of $\mathfrak{su}(2D)$, generated by

$$L_{\mu\nu} = -i(\bar{E}_{\mu\nu} - \bar{E}_{\nu\mu}) = -i(E_{\mu\nu} - E_{\nu\mu}), \quad (2.16)$$

such that

$$[L_{\mu\nu}, L_{\mu'\nu'}] = i(\delta_{\mu,\mu'}L_{\nu\nu'} - \delta_{\mu,\nu'}L_{\nu\mu'} - \delta_{\nu,\mu'}L_{\mu\nu'} + \delta_{\nu,\nu'}L_{\mu\mu'}), \quad L_{\mu\nu}^\dagger = L_{\mu\nu} = -L_{\nu\mu},$$

is explicitly reduced. Its unirreps are characterized by $2j$, which runs over $N, N-2, \dots, 0$ (or 1) for a given N . The remaining generators of $\mathfrak{su}(2D)$ may be taken as

$$T_{\mu\nu} = \bar{E}_{\mu\nu} + \bar{E}_{\nu\mu}. \quad (2.17)$$

The operators

$$D_{\mu\nu}^\dagger = \alpha_\mu^\dagger \alpha_\nu^\dagger, \quad D_{\mu\nu} = \alpha_\mu \alpha_\nu \quad (2.18)$$

act as raising and lowering operators relating among themselves wavefunctions corresponding to even or odd values of N . Together with $E_{\mu\nu}$, they generate an $\mathfrak{sp}(4D, \mathbb{R})$ Lie algebra, whose (nonvanishing) commutation relations are given by

$$\begin{aligned} [E_{\mu\nu}, E_{\mu'\nu'}] &= \delta_{\nu,\mu'}E_{\mu\nu'} - \delta_{\mu,\nu'}E_{\mu'\nu}, \\ [E_{\mu\nu}, D_{\mu'\nu'}^\dagger] &= \delta_{\nu,\mu'}D_{\mu\nu'}^\dagger + \delta_{\nu,\nu'}D_{\mu\mu'}^\dagger, \\ [E_{\mu\nu}, D_{\mu'\nu'}] &= -\delta_{\mu,\mu'}D_{\nu\nu'} - \delta_{\mu,\nu'}D_{\nu\mu'}, \\ [D_{\mu\nu}, D_{\mu'\nu'}^\dagger] &= \delta_{\mu,\mu'}E_{\nu'\nu} + \delta_{\mu,\nu'}E_{\mu'\nu} + \delta_{\nu,\mu'}E_{\nu'\mu} + \delta_{\nu,\nu'}E_{\mu'\mu}. \end{aligned}$$

To connect the wavefunctions with an even N value to those with an odd one, we have to use the bosonic creation and annihilation operators (2.13), which generate a Weyl algebra $\mathfrak{w}(2D)$, specified by

$$[\alpha_\mu, \alpha_\nu^\dagger] = \delta_{\mu,\nu}I.$$

The whole set of operators $\{E_{\mu\nu}, D_{\mu\nu}^\dagger, D_{\mu\nu}, \alpha_\mu^\dagger, \alpha_\mu, I\}$ then provides us with the harmonic oscillator dynamical algebra, which is the semidirect sum Lie algebra $\mathfrak{w}(2D) \oplus_s \mathfrak{sp}(4D, \mathbb{R})$, as shown by the remaining (nonvanishing) commutation relations

$$\begin{aligned} [E_{\mu\nu}, \alpha_{\mu'}^\dagger] &= \delta_{\nu,\mu'}\alpha_{\mu'}^\dagger, & [E_{\mu\nu}, \alpha_{\mu'}] &= -\delta_{\mu,\mu'}\alpha_{\nu}, \\ [D_{\mu\nu}, \alpha_{\mu'}^\dagger] &= \delta_{\mu,\mu'}\alpha_{\nu} + \delta_{\nu,\mu'}\alpha_{\mu}, & [D_{\mu\nu}^\dagger, \alpha_{\mu'}] &= -\delta_{\mu,\mu'}\alpha_{\nu}^\dagger - \delta_{\nu,\mu'}\alpha_{\mu}^\dagger. \end{aligned}$$

To apply the symmetry and dynamical algebra generators to the oscillator wavefunctions (2.3) written in the variables $R, \boldsymbol{\theta}, \boldsymbol{\lambda}$, we have to express the creation and annihilation operators $\alpha_\mu^\dagger, \alpha_\mu$ in such variables. This implies combining the transformation (2.1) with the corresponding change for the partial differential operators

$$\begin{aligned} \partial_{X_{2\nu-1}} &= \sin \lambda_\nu \partial^{(\nu,1)} + \cos \lambda_\nu \partial^{(\nu,2)}, \\ \partial_{X_{2\nu}} &= \cos \lambda_\nu \partial^{(\nu,1)} - \sin \lambda_\nu \partial^{(\nu,2)}, \quad \nu = 1, 2, \dots, D, \end{aligned} \quad (2.19)$$

where

$$\partial^{(1,1)} = \sin \theta_1 \sin \theta_2 \cdots \sin \theta_{D-1} \partial_R$$

$$\begin{aligned}
& + \frac{1}{R} \sum_{\rho=1}^{D-1} \csc \theta_1 \csc \theta_2 \cdots \csc \theta_{\rho-1} \cos \theta_\rho \sin \theta_{\rho+1} \sin \theta_{\rho+2} \cdots \sin \theta_{D-1} \partial_{\theta_\rho}, \\
\partial^{(\nu,1)} & = \sin \theta_1 \sin \theta_2 \cdots \sin \theta_{D-\nu} \cos \theta_{D-\nu+1} \partial_R \\
& + \frac{1}{R} \sum_{\rho=1}^{D-\nu} \csc \theta_1 \csc \theta_2 \cdots \csc \theta_{\rho-1} \cos \theta_\rho \sin \theta_{\rho+1} \sin \theta_{\rho+2} \cdots \sin \theta_{D-\nu} \cos \theta_{D-\nu+1} \partial_{\theta_\rho} \\
& - \frac{1}{R} \csc \theta_1 \csc \theta_2 \cdots \csc \theta_{D-\nu} \sin \theta_{D-\nu+1} \partial_{\theta_{D-\nu+1}}, \quad \nu = 2, 3, \dots, D-1, \\
\partial^{(D,1)} & = \cos \theta_1 \partial_R - \frac{1}{R} \sin \theta_1 \partial_{\theta_1},
\end{aligned}$$

and

$$\begin{aligned}
\partial^{(1,2)} & = \frac{1}{R} \csc \theta_1 \csc \theta_2 \cdots \csc \theta_{D-1} \partial_{\lambda_1}, \\
\partial^{(\nu,2)} & = \frac{1}{R} \csc \theta_1 \csc \theta_2 \cdots \csc \theta_{D-\nu} \sec \theta_{D-\nu+1} \partial_{\lambda_\nu}, \quad \nu = 2, 3, \dots, D-1, \\
\partial^{(D,2)} & = \frac{1}{R} \sec \theta_1 \partial_{\lambda_D}.
\end{aligned}$$

We shall carry out this transformation explicitly for $D = 2$ in Section 4.

3 Reduction of the $2D$ -dimensional harmonic oscillator to the D -dimensional SW system

To go from the $2D$ -dimensional harmonic oscillator Hamiltonian H^{osc} to some extended SW Hamiltonian H , let us first transform the original Cartesian coordinates X_μ , $\mu = 1, 2, \dots, 2D$, into some new ones x_μ , $\mu = 1, 2, \dots, 2D$, such that

$$\begin{aligned}
x_1 & = r \sin \phi_1 \sin \phi_2 \cdots \sin \phi_{D-1}, \\
x_\nu & = r \sin \phi_1 \sin \phi_2 \cdots \sin \phi_{D-\nu} \cos \phi_{D-\nu+1}, \quad \nu = 2, 3, \dots, D-1, \\
x_D & = r \cos \phi_1, \\
x_{D+\nu} & = \lambda_\nu, \quad \nu = 1, 2, \dots, D,
\end{aligned}$$

and

$$\begin{aligned}
R & = \sqrt{\omega} r, \quad \theta_\nu = \phi_\nu, \quad \nu = 1, 2, \dots, D-1, \\
0 \leq r & < \infty, \quad 0 \leq \phi_\nu < \frac{\pi}{2}, \quad \nu = 1, 2, \dots, D-1, \quad 0 \leq \lambda_\nu < 2\pi, \quad \nu = 1, 2, \dots, D.
\end{aligned}$$

Here $r, \phi_1, \phi_2, \dots, \phi_{D-1}$ are hyperspherical coordinates in the D -dimensional subspace (x_1, x_2, \dots, x_D) . The volume element in the transformed $2D$ -dimensional space is given by

$$dv = \prod_{\mu=1}^{2D} dx_\mu = r^{D-1} dr \left[\prod_{\nu=1}^{D-1} (\sin \phi_\nu)^{D-\nu-1} d\phi_\nu \right] \left(\prod_{\nu=1}^D d\lambda_\nu \right). \quad (3.1)$$

On making next the change of function

$$\Psi(r, \phi, \lambda) = \mathcal{O}^{1/2} \Psi^{\text{osc}}(R, \theta, \lambda), \quad (3.2)$$

with

$$\mathcal{O} = (\omega r)^D \prod_{\nu=1}^{D-1} (\sin \phi_\nu)^{D-\nu} \cos \phi_\nu, \quad (3.3)$$

the harmonic oscillator wavefunctions $\Psi^{\text{osc}}(R, \boldsymbol{\theta}, \boldsymbol{\lambda})$, living in a Hilbert space with measure dV given in (2.5), are mapped onto some functions $\Psi(r, \boldsymbol{\phi}, \boldsymbol{\lambda})$, living in a Hilbert space with measure dv defined in (3.1). As a consequence of (2.4), we obtain

$$\int dv |\Psi(r, \boldsymbol{\phi}, \boldsymbol{\lambda})|^2 = 1.$$

By this unitary transformation, the harmonic oscillator Hamiltonian H^{osc} is changed into

$$H/\omega = \mathcal{O}^{1/2} H^{\text{osc}} \mathcal{O}^{-1/2} \quad (3.4)$$

and similarly for other operators acting in the harmonic oscillator Hilbert space. A straightforward calculation leads to the result

$$\begin{aligned} H = & -\partial_r^2 - \frac{D-1}{r} \partial_r - \frac{1}{r^2} \left\{ \partial_{\phi_1}^2 + (D-2) \cot \phi_1 \partial_{\phi_1} \right. \\ & + \sum_{\nu=2}^{D-1} \frac{1}{\sin^2 \phi_1 \sin^2 \phi_2 \cdots \sin^2 \phi_{\nu-1}} [\partial_{\phi_\nu}^2 + (D-\nu-1) \cot \phi_\nu \partial_{\phi_\nu}] \\ & + \frac{1}{\sin^2 \phi_1 \sin^2 \phi_2 \cdots \sin^2 \phi_{D-1}} \left(\partial_{\lambda_1}^2 + \frac{1}{4} \right) \\ & \left. + \sum_{\nu=2}^{D-1} \frac{1}{\sin^2 \phi_1 \sin^2 \phi_2 \cdots \sin^2 \phi_{D-\nu} \cos^2 \phi_{D-\nu+1}} \left(\partial_{\lambda_\nu}^2 + \frac{1}{4} \right) + \frac{1}{\cos^2 \phi_1} \left(\partial_{\lambda_D}^2 + \frac{1}{4} \right) \right\} \\ & + \omega^2 r^2. \end{aligned} \quad (3.5)$$

The eigenvalues of H are directly obtained from (2.11) as

$$E_{n_r, j} = 2\omega(2n_r + 2j + D), \quad n_r = 0, 1, 2, \dots, \quad j = 0, \frac{1}{2}, 1, \frac{3}{2}, \dots \quad (3.6)$$

The corresponding wavefunctions can be derived from (2.3), (2.6), (2.7), (2.9), (2.12), (3.2), and (3.3) and read

$$\begin{aligned} \Psi_{n_r, \mathbf{n}\mathbf{p}}(r, \boldsymbol{\phi}, \boldsymbol{\lambda}) &= \mathcal{N}_{n_r, \mathbf{n}\mathbf{p}} \mathcal{Z}_{n_r}^{(j)}(z) \Phi_{\mathbf{n}}^{(\mathbf{p})}(\boldsymbol{\phi}) \left(\prod_{\nu=1}^D e^{ip_{D-\nu+1} \lambda_\nu} \right), \\ \mathcal{Z}_{n_r}^{(j)}(z) &= \left(\frac{z}{\omega} \right)^{j+\frac{D}{4}} L_{n_r}^{(2j+D-1)}(z) e^{-\frac{1}{2}z}, \quad z = \omega r^2, \\ \Phi_{\mathbf{n}}^{(\mathbf{p})}(\boldsymbol{\phi}) &= \prod_{\nu=1}^{D-1} \Phi_{n_\nu}^{(a_\nu, b_\nu)}(\phi_\nu) \\ &= \prod_{\nu=1}^{D-1} (\cos \phi_\nu)^{a_\nu} (\sin \phi_\nu)^{b_\nu + \frac{1}{2}(D-\nu-1)} P_{n_\nu}^{(a_\nu - \frac{1}{2}, b_\nu + D - \nu - \frac{3}{2})}(-\cos 2\phi_\nu), \\ \mathcal{N}_{n_r, \mathbf{n}\mathbf{p}} &= \omega^{j+\frac{D}{2}} \mathcal{N}_{n_r, \mathbf{n}\mathbf{p}}^{\text{osc}}, \end{aligned} \quad (3.7)$$

where $\mathbf{n} = n_1 n_2 \cdots n_{D-1}$, $\mathbf{p} = p_1 p_2 \cdots p_D$, $n_r, n_1, n_2, \dots, n_{D-1} \in \mathbb{N}$, $p_1, p_2, \dots, p_D \in \mathbb{Z}$, while j , and a_ν, b_ν are defined in (2.10) and (2.8), respectively.

In the subspace of functions $\Psi_{n_r, \mathbf{n}\mathbf{p}}(r, \boldsymbol{\phi}, \boldsymbol{\lambda})$ with fixed \mathbf{p} , the Hamiltonian H , defined in (3.5), has the same action as the D -dimensional Hamiltonian

$$H^{(\mathbf{k})} = -\partial_r^2 - \frac{D-1}{r} \partial_r - \frac{1}{r^2} \left\{ \partial_{\phi_1}^2 + (D-2) \cot \phi_1 \partial_{\phi_1} \right.$$

$$\begin{aligned}
& + \left. \sum_{\nu=2}^{D-1} \frac{1}{\sin^2 \phi_1 \sin^2 \phi_2 \cdots \sin^2 \phi_{\nu-1}} [\partial_{\phi_\nu}^2 + (D - \nu - 1) \cot \phi_\nu \partial_{\phi_\nu}] \right\} \\
& + \frac{k_1^2}{r^2 \sin^2 \phi_1 \sin^2 \phi_2 \cdots \sin^2 \phi_{D-1}} + \sum_{\nu=2}^{D-1} \frac{k_\nu^2}{r^2 \sin^2 \phi_1 \sin^2 \phi_2 \cdots \sin^2 \phi_{D-\nu} \cos^2 \phi_{D-\nu+1}} \\
& + \frac{k_D^2}{r^2 \cos^2 \phi_1} + \omega^2 r^2,
\end{aligned}$$

where we have defined $\mathbf{k} = k_1 k_2 \cdots k_D$ and

$$k_\nu = \sqrt{p_{D-\nu+1}^2 - \frac{1}{4}}, \quad \nu = 1, 2, \dots, D. \quad (3.8)$$

The latter Hamiltonian is but the SW one (1.1), expressed in hyperspherical coordinates $r, \phi_1, \phi_2, \dots, \phi_{D-1}$. We conclude that H is an extension of $H^{(\mathbf{k})}$, resulting from the introduction of D auxiliary continuous variables $\lambda_\nu = x_{D+\nu}$, $\nu = 1, 2, \dots, D$, and that, conversely, $H^{(\mathbf{k})}$ is obtained from H by projecting it down into the D -dimensional subspace (x_1, x_2, \dots, x_D) .³

As a by-product of this reduction process, we have determined the wavefunctions $\Psi_{n_r, \mathbf{n}}^{(\mathbf{k})}(r, \phi)$ of $H^{(\mathbf{k})}$ in hyperspherical coordinates. Equation (3.7) may indeed be rewritten as

$$\begin{aligned}
\Psi_{n_r, \mathbf{n}\mathbf{p}}(r, \phi, \boldsymbol{\lambda}) &= \Psi_{n_r, \mathbf{n}}^{(\mathbf{k})}(r, \phi) (2\pi)^{-D/2} \prod_{\nu=1}^D e^{ip_{D-\nu+1} \lambda_\nu}, \\
\Psi_{n_r, \mathbf{n}}^{(\mathbf{k})}(r, \phi) &= \mathcal{N}_{n_r, \mathbf{n}}^{(\mathbf{k})} \mathcal{Z}_{n_r}^{(j)}(z) \Phi_{\mathbf{n}}^{(\mathbf{p})}(\phi), \quad \mathcal{N}_{n_r, \mathbf{n}}^{(\mathbf{k})} = (2\pi)^{D/2} \mathcal{N}_{n_r, \mathbf{n}\mathbf{p}},
\end{aligned} \quad (3.9)$$

with \mathbf{k} and \mathbf{p} related as in (3.8).

By a transformation similar to (3.4), the generators $\bar{E}_{\mu\nu}$ of the harmonic oscillator symmetry algebra $\mathfrak{su}(2D)$ are changed into some operators acting on $\Psi_{n_r, \mathbf{n}\mathbf{p}}(r, \phi, \boldsymbol{\lambda})$. Since the latter may change n_r and j separately (provided their sum $n_r + j = N/2$ is preserved), this means in particular (see equation (2.10)) that the p_ν 's (hence the k_ν 's) may change too. The transformed $\mathfrak{su}(2D)$ algebra may therefore connect among themselves some wavefunctions of H belonging to the same energy eigenvalue (3.6), but associated with different reduced Hamiltonians $H^{(\mathbf{k})}$. We conclude that it provides us with a potential algebra for the SW system. Similarly, the transformed $\mathfrak{w}(2D) \oplus_s \mathfrak{sp}(4D, \mathbb{R})$ algebra will be a dynamical potential algebra for the same.

As a final point, it is worth observing that in the harmonic oscillator wavefunctions (2.3), the quantum numbers p_ν , $\nu = 1, 2, \dots, D$, run over \mathbb{Z} . On the other hand, in (1.1), the parameters k_ν , $\nu = 1, 2, \dots, D$, have been assumed real and positive. From equation (3.8), however, it is clear that $p_{D-\nu+1} = 0$ would lead to an imaginary value of k_ν and to unphysical wavefunctions (3.9), while $|p_{D-\nu+1}|$ and $-|p_{D-\nu+1}|$ with $|p_{D-\nu+1}| \geq 1$ would give rise to the same k_ν , hence to some replicas of physical wavefunctions (3.9). The correspondence between the harmonic oscillator wavefunctions and the extended SW ones is therefore not one-to-one. This lack of bijectiveness is a known aspect of potential algebraic approaches (see [69] where this phenomenon was first pointed out).

4 The two-dimensional case

4.1 Harmonic oscillator symmetry and dynamical algebras

To deal in detail with the two-dimensional case, it is appropriate to rewrite the four-dimensional harmonic oscillator wavefunctions $\Psi_{n_r, n, p_1, p_2}^{\text{osc}}(R, \theta, \lambda_1, \lambda_2)$ (with $j = n + \frac{1}{2}(|p_1| + |p_2|)$) in an

³Strictly speaking, this is true only for those k_ν 's that can be written in the form (3.8) with integer $p_{D-\nu+1}$.

equivalent form $\bar{\Psi}_{n_r, j, m, m'}^{\text{osc}}(R, \theta, \lambda_1, \lambda_2)$ using either hyperspherical harmonics $Y_{2j, m, m'}(\alpha, \beta, \gamma)$ or (complex conjugate) rotation matrix elements $D_{m, -m'}^{j*}(\alpha, \beta, \gamma)$ expressed in terms of Euler angles α, β, γ [85],

$$Y_{2j, m, m'}(\alpha, \beta, \gamma) = (-1)^{j-m'} \left(\frac{2j+1}{2\pi^2} \right)^{1/2} D_{m, -m'}^{j*}(\alpha, \beta, \gamma),$$

$$D_{m, -m'}^{j*}(\alpha, \beta, \gamma) = e^{im\alpha} d_{m, -m'}^j(\beta) e^{-im'\gamma}.$$

Here j runs over $0, \frac{1}{2}, 1, \frac{3}{2}, \dots$, while m and m' take values in the set $\{j, j-1, \dots, -j\}$. On setting

$$\theta = \frac{1}{2}\beta, \quad \lambda_1 = \frac{1}{2}(\gamma - \alpha), \quad \lambda_2 = \frac{1}{2}(\gamma + \alpha), \quad p_1 = m - m', \quad p_2 = -m - m',$$

or, conversely,

$$\alpha = \lambda_2 - \lambda_1, \quad \beta = 2\theta, \quad \gamma = \lambda_2 + \lambda_1, \quad m = \frac{1}{2}(p_1 - p_2), \quad m' = -\frac{1}{2}(p_1 + p_2),$$

and on using the relation between rotation functions $d_{m, -m'}^j(\beta)$ and Jacobi polynomials [85], we indeed get

$$\bar{\Psi}_{n_r, n, p_1, p_2}^{\text{osc}}(R, \theta, \lambda_1, \lambda_2) = (-1)^{\frac{1}{2}(|p_1|+p_1)+|p_2|} \bar{\Psi}_{n_r, j, m, m'}^{\text{osc}}(R, \theta, \lambda_1, \lambda_2) \quad (4.1)$$

with

$$\bar{\Psi}_{n_r, j, m, m'}^{\text{osc}}(R, \theta, \lambda_1, \lambda_2) = (-1)^{j-m'} \left(\frac{(2j+1)n_r!}{\pi^2(n_r+2j+1)!} \right)^{1/2} \mathcal{L}_{n_r}^{(j)}(z)$$

$$\times d_{m, -m'}^j(2\theta) e^{-i(m+m')\lambda_1} e^{i(m-m')\lambda_2},$$

$$\mathcal{L}_{n_r}^{(j)}(z) = z^j L_{n_r}^{(2j+1)}(z) e^{-\frac{1}{2}z}, \quad z = R^2. \quad (4.2)$$

The advantage of this new form is that the wavefunctions $\Psi_{n_r, n, p_1, p_2}^{\text{osc}}(R, \theta, \lambda_1, \lambda_2)$, which were classified according to

$$\begin{array}{ccc} \text{su}(4) & \supset & \text{so}(4) \\ [N] & & (2j) \end{array}$$

with $N = 2n_r + 2j$ and $2j = 2n + |p_1| + |p_2|$, now turn out to be explicitly reduced with respect to

$$\begin{array}{ccc} \text{su}(4) & \supset & \text{so}(4) \simeq \text{su}(2) \oplus \text{su}(2) & \supset & \text{u}(1) \oplus \text{u}(1) \\ [N] & & (2j) \simeq [j] \oplus [j] & & [m] \oplus [m'] \end{array} \quad (4.3)$$

This will allow us to use the full machinery of angular momentum theory for determining the explicit action of the symmetry and dynamical algebra generators on wavefunctions.

The two $\text{su}(2)$ algebras appearing in chain (4.3) are generated by J_i and K_i , $i = 1, 2, 3$, defined in terms of $L_{\mu\nu}$, $\mu, \nu = 1, 2, 3, 4$, (see equation (2.16)) by

$$J_i = \frac{1}{2} \left(\frac{1}{2} \epsilon_{ijk} L_{jk} - L_{i4} \right), \quad K_i = \frac{1}{2} \left(\frac{1}{2} \epsilon_{ijk} L_{jk} + L_{i4} \right), \quad (4.4)$$

where i, j, k run over 1, 2, 3 and ϵ_{ijk} is the antisymmetric tensor. The operators J_i and K_i satisfy the relations

$$[J_i, J_j] = i\epsilon_{ijk} J_k, \quad [K_i, K_j] = i\epsilon_{ijk} K_k, \quad [J_i, K_j] = 0,$$

$$J_i^\dagger = J_i, \quad K_i^\dagger = K_i.$$

Instead of the Cartesian components of \mathbf{J} and \mathbf{K} , we may use alternatively $J_0 = J_3$, $J_{\pm} = J_1 \pm iJ_2$, $K_0 = K_3$, $K_{\pm} = K_1 \pm iK_2$, with J_0 and K_0 generating the two $u(1)$ subalgebras in (4.3).

The differential operator form of J_0 , J_{\pm} , K_0 , and K_{\pm} can be obtained by combining equations (2.1), (2.13), (2.14), (2.16), (2.19), and (4.4) and is given by

$$\begin{aligned} J_0 &= \frac{1}{2}(\partial_{\lambda_1} - \partial_{\lambda_2}), & J_{\pm} &= \frac{1}{2}e^{\mp i(\lambda_1 - \lambda_2)}[\pm \partial_{\theta} - i(\cot \theta \partial_{\lambda_1} + \tan \theta \partial_{\lambda_2})], \\ K_0 &= \frac{1}{2}(\partial_{\lambda_1} + \partial_{\lambda_2}), & K_{\pm} &= \frac{1}{2}e^{\mp i(\lambda_1 + \lambda_2)}[\mp \partial_{\theta} + i(\cot \theta \partial_{\lambda_1} - \tan \theta \partial_{\lambda_2})]. \end{aligned}$$

From some differential equation relations satisfied by rotation functions $d_{m,-m'}^j(2\theta)$ [86], it is then easy to check that

$$\begin{aligned} J_0 \bar{\Psi}_{n_r, j, m, m'}^{\text{osc}} &= m \bar{\Psi}_{n_r, j, m, m'}^{\text{osc}}, & J_{\pm} \bar{\Psi}_{n_r, j, m, m'}^{\text{osc}} &= [(j \mp m)(j \pm m + 1)]^{1/2} \bar{\Psi}_{n_r, j, m \pm 1, m'}^{\text{osc}}, \\ K_0 \bar{\Psi}_{n_r, j, m, m'}^{\text{osc}} &= m' \bar{\Psi}_{n_r, j, m, m'}^{\text{osc}}, & K_{\pm} \bar{\Psi}_{n_r, j, m, m'}^{\text{osc}} &= [(j \mp m')(j \pm m' + 1)]^{1/2} \bar{\Psi}_{n_r, j, m, m' \pm 1}^{\text{osc}}, \end{aligned}$$

which proves the above-mentioned result.

It is now convenient to rewrite all operators of physical interest as components $T_{\sigma, \tau}^{(s, t)}$, $\sigma = s, s-1, \dots, -s$, $\tau = t, t-1, \dots, -t$, of irreducible tensors of rank (s, t) with respect to $\mathfrak{su}(2) \oplus \mathfrak{su}(2)$. These must satisfy commutation relations of the type

$$\begin{aligned} [J_0, T_{\sigma, \tau}^{(s, t)}] &= \sigma T_{\sigma, \tau}^{(s, t)}, & [J_{\pm}, T_{\sigma, \tau}^{(s, t)}] &= [(s \mp \sigma)(s \pm \sigma + 1)]^{1/2} T_{\sigma \pm 1, \tau}^{(s, t)}, \\ [K_0, T_{\sigma, \tau}^{(s, t)}] &= \tau T_{\sigma, \tau}^{(s, t)}, & [K_{\pm}, T_{\sigma, \tau}^{(s, t)}] &= [(t \mp \tau)(t \pm \tau + 1)]^{1/2} T_{\sigma, \tau \pm 1}^{(s, t)}. \end{aligned}$$

Since the bosonic creation and annihilation operators serve as building blocks for the construction of other operators, let us start with them. The creation operators can be written as components $\mathcal{A}_{\sigma, \tau}^{\dagger}$, $\sigma, \tau = \frac{1}{2}, -\frac{1}{2}$, of an irreducible tensor of rank $(\frac{1}{2}, \frac{1}{2})$,

$$\mathcal{A}_{\pm \frac{1}{2}, \pm \frac{1}{2}}^{\dagger} = \mp \frac{1}{\sqrt{2}}(\alpha_1^{\dagger} \pm i\alpha_2^{\dagger}), \quad \mathcal{A}_{\pm \frac{1}{2}, \mp \frac{1}{2}}^{\dagger} = \frac{1}{\sqrt{2}}(\alpha_3^{\dagger} \mp i\alpha_4^{\dagger}). \quad (4.5)$$

The same is true for the annihilation operators, the corresponding components being given by

$$\mathcal{A}_{\sigma, \tau} = (-1)^{1-\sigma-\tau}(\mathcal{A}_{-\sigma, -\tau}^{\dagger})^{\dagger}, \quad \sigma, \tau = \frac{1}{2}, -\frac{1}{2}. \quad (4.6)$$

On coupling an operator \mathcal{A}^{\dagger} with an operator \mathcal{A} according to

$$[\mathcal{A}^{\dagger} \times \mathcal{A}]_{\sigma, \tau}^{s, t} = \sum_{\sigma', \tau'} \langle \frac{1}{2} \sigma', \frac{1}{2} \sigma - \sigma' | s \sigma \rangle \langle \frac{1}{2} \tau', \frac{1}{2} \tau - \tau' | t \tau \rangle \mathcal{A}_{\sigma', \tau'}^{\dagger} \mathcal{A}_{\sigma - \sigma', \tau - \tau'},$$

where $\langle, | \rangle$ denotes an $\mathfrak{su}(2)$ Wigner coefficient [85], we obtain the $\mathfrak{su}(4)$ symmetry algebra generators classified with respect to chain (4.3). These include

$$J_{\sigma} = [\mathcal{A}^{\dagger} \times \mathcal{A}]_{\sigma, 0}^{1, 0}, \quad K_{\tau} = [\mathcal{A}^{\dagger} \times \mathcal{A}]_{0, \tau}^{0, 1}, \quad \sigma, \tau = +1, 0, -1,$$

with $J_{\pm 1} = \mp J_{\pm} / \sqrt{2}$ and $K_{\pm 1} = \mp K_{\pm} / \sqrt{2}$, as well as the nine components of an irreducible tensor of rank $(1, 1)$,

$$\mathcal{T}_{\sigma, \tau} = [\mathcal{A}^{\dagger} \times \mathcal{A}]_{\sigma, \tau}^{1, 1}, \quad \sigma, \tau = +1, 0, -1. \quad (4.7)$$

The latter may be written as

$$\begin{aligned} \mathcal{T}_{\pm 1, \pm 1} &= -\frac{1}{4}(T_{11} \pm 2iT_{12} - T_{22}), & \mathcal{T}_{\pm 1, 0} &= \frac{1}{2\sqrt{2}}(\pm T_{13} - iT_{14} + iT_{23} \pm T_{24}), \\ \mathcal{T}_{\pm 1, \mp 1} &= -\frac{1}{4}(T_{33} \mp 2iT_{34} - T_{44}), & \mathcal{T}_{0, \pm 1} &= \frac{1}{2\sqrt{2}}(\pm T_{13} + iT_{14} + iT_{23} \mp T_{24}), \end{aligned}$$

$$\mathcal{T}_{0,0} = \frac{1}{2}(T_{11} + T_{22}) = -\frac{1}{2}(T_{33} + T_{44})$$

in terms of the operators $T_{\mu\nu}$, defined in (2.17). Observe that the $u(4)$ first-order Casimir operator (2.15) is, up to some constants, the scalar that can be obtained in such a coupling procedure,

$$[\mathcal{A}^\dagger \times \mathcal{A}]_{0,0}^{0,0} = \frac{1}{2}(\mathcal{E} - 2).$$

Similarly, the coupling of two operators \mathcal{A}^\dagger provides us with the raising operators belonging to $\mathfrak{sp}(8, \mathbb{R})$,

$$\mathcal{D}^\dagger = \mathcal{A}^\dagger \cdot \mathcal{A}^\dagger = -2[\mathcal{A}^\dagger \times \mathcal{A}^\dagger]_{0,0}^{0,0}, \quad \mathcal{D}_{\sigma,\tau}^\dagger = [\mathcal{A}^\dagger \times \mathcal{A}^\dagger]_{\sigma,\tau}^{1,1}, \quad \sigma, \tau = +1, 0, -1,$$

or, in detail,

$$\mathcal{D}^\dagger = D_{11}^\dagger + D_{22}^\dagger + D_{33}^\dagger + D_{44}^\dagger$$

and

$$\begin{aligned} \mathcal{D}_{\pm 1, \pm 1}^\dagger &= \frac{1}{2}(D_{11}^\dagger \pm 2iD_{12}^\dagger - D_{22}^\dagger), & \mathcal{D}_{\pm 1, 0}^\dagger &= -\frac{1}{\sqrt{2}}(\pm D_{13}^\dagger - iD_{14}^\dagger + iD_{23}^\dagger \pm D_{24}^\dagger), \\ \mathcal{D}_{\pm 1, \mp 1}^\dagger &= \frac{1}{2}(D_{33}^\dagger \mp 2iD_{34}^\dagger - D_{44}^\dagger), & \mathcal{D}_{0, \pm 1}^\dagger &= -\frac{1}{\sqrt{2}}(\pm D_{13}^\dagger + iD_{14}^\dagger + iD_{23}^\dagger \mp D_{24}^\dagger), \\ \mathcal{D}_{0, 0}^\dagger &= \frac{1}{2}(-D_{11}^\dagger - D_{22}^\dagger + D_{33}^\dagger + D_{44}^\dagger) \end{aligned}$$

in terms of $D_{\mu\nu}^\dagger$ defined in (2.18). The corresponding lowering operators are then

$$\mathcal{D} = (\mathcal{D}^\dagger)^\dagger, \quad \mathcal{D}_{\sigma,\tau} = (-1)^{\sigma+\tau}(\mathcal{D}_{-\sigma,-\tau}^\dagger)^\dagger, \quad \sigma, \tau = +1, 0, -1.$$

It is now straightforward to determine the action of $\mathcal{A}_{\sigma,\tau}^\dagger$ on the wavefunctions $\bar{\Psi}_{n_r, j, m, m'}^{\text{osc}}(R, \theta, \lambda_1, \lambda_2)$. Application of the Wigner–Eckart theorem with respect to $\mathfrak{su}(2) \oplus \mathfrak{su}(2)$ [85] indeed leads to the relation

$$\begin{aligned} \mathcal{A}_{\sigma,\tau}^\dagger \bar{\Psi}_{n_r, j, m, m'}^{\text{osc}} &= \sum_{n'_r, j'} \langle n'_r, j' | \mathcal{A}^\dagger | n_r, j \rangle \langle j, m, \frac{1}{2} \sigma | j', m + \sigma \rangle \langle j, m', \frac{1}{2} \tau | j', m' + \tau \rangle \\ &\times \bar{\Psi}_{n'_r, j', m + \sigma, m' + \tau}^{\text{osc}}, \end{aligned} \quad (4.8)$$

where $\langle n'_r, j' | \mathcal{A}^\dagger | n_r, j \rangle$ denotes a reduced matrix element, the summation over j' runs over $j + \frac{1}{2}, j - \frac{1}{2}$, and n'_r is determined by the selection rule $n'_r + j' = n_r + j + \frac{1}{2}$ implying that $n'_r = n_r, n_r + 1$, respectively. To calculate the two independent reduced matrix elements, it is enough to consider equation (4.8) for the special case $m = m' = j$ and to use the differential operator form of $\mathcal{A}_{\pm \frac{1}{2}, \pm \frac{1}{2}}^\dagger$,

$$\mathcal{A}_{\pm \frac{1}{2}, \pm \frac{1}{2}}^\dagger = \frac{1}{2} e^{\mp i \lambda_1} \left[i \left(\sin \theta \partial_R + \frac{1}{R} \cos \theta \partial_\theta \right) \pm \frac{1}{R} \csc \theta \partial_{\lambda_1} - i R \sin \theta \right],$$

following from (2.1), (2.13), (2.19), and (4.5). Simple properties of the rotation function $d_{m, -m'}^j(2\theta)$ and of the Laguerre polynomial $L_{n_r}^{(2j+1)}(z)$ then lead to the results

$$\langle n_r, j + \frac{1}{2} | \mathcal{A}^\dagger | n_r, j \rangle = i \left(\frac{(2j+1)(n_r+2j+2)}{2j+2} \right)^{1/2},$$

$$\langle n_r + 1, j - \frac{1}{2} \| \mathcal{A}^\dagger \| n_r, j \rangle = -i \left(\frac{(2j+1)(n_r+1)}{2j} \right)^{1/2}. \quad (4.9)$$

The operators $\mathcal{A}_{\sigma,\tau}$ satisfy an equation similar to (4.8) with $\langle n'_r, j' \| \mathcal{A}^\dagger \| n_r, j \rangle$ replaced by $\langle n'_r, j' \| \mathcal{A} \| n_r, j \rangle$ and $n'_r = n_r - 1, n_r$ for $j' = j + \frac{1}{2}, j - \frac{1}{2}$, respectively. The corresponding reduced matrix elements can be directly calculated from the relation

$$\langle n'_r, j' \| \mathcal{A} \| n_r, j \rangle = \frac{2j+1}{2j'+1} \langle n_r, j \| \mathcal{A}^\dagger \| n'_r, j' \rangle^*, \quad (4.10)$$

which is a direct consequence of (4.6).

For the $\mathfrak{su}(4)$ generators that do not belong to $\mathfrak{so}(4)$, we get the equation

$$\begin{aligned} \mathcal{T}_{\sigma,\tau} \bar{\Psi}_{n_r, j, m, m'}^{\text{osc}} &= \sum_{n'_r, j'} \langle n'_r, j' \| \mathcal{T} \| n_r, j \rangle \langle j' m, 1 \sigma | j' m + \sigma \rangle \langle j m', 1 \tau | j' m' + \tau \rangle \\ &\times \bar{\Psi}_{n'_r, j', m + \sigma, m' + \tau}^{\text{osc}}, \end{aligned} \quad (4.11)$$

where $j' = j+1, j, j-1$ and $n'_r = n_r - 1, n_r, n_r + 1$, respectively. Equation (4.7) and the coupling law for reduced matrix elements [85] enable us to determine

$$\begin{aligned} \langle n_r - 1, j + 1 \| \mathcal{T} \| n_r, j \rangle &= - \left(\frac{(2j+1)n_r(n_r+2j+2)}{2j+3} \right)^{1/2}, \\ \langle n_r, j \| \mathcal{T} \| n_r, j \rangle &= n_r + j + 1, \\ \langle n_r + 1, j - 1 \| \mathcal{T} \| n_r, j \rangle &= - \left(\frac{(2j+1)(n_r+1)(n_r+2j+1)}{2j-1} \right)^{1/2} \end{aligned}$$

from (4.9) and (4.10).

The operators $\mathcal{D}_{\sigma,\tau}^\dagger$ and $\mathcal{D}_{\sigma,\tau}$ satisfy a relation similar to (4.11) with

$$\begin{aligned} \langle n_r, j + 1 \| \mathcal{D}^\dagger \| n_r, j \rangle &= - \left(\frac{(2j+1)(n_r+2j+2)(n_r+2j+3)}{2j+3} \right)^{1/2}, \\ \langle n_r + 1, j \| \mathcal{D}^\dagger \| n_r, j \rangle &= [(n_r+1)(n_r+2j+2)]^{1/2}, \\ \langle n_r + 2, j - 1 \| \mathcal{D}^\dagger \| n_r, j \rangle &= - \left(\frac{(2j+1)(n_r+1)(n_r+2)}{2j-1} \right)^{1/2}, \end{aligned}$$

and $\langle n'_r, j' \| \mathcal{D} \| n_r, j \rangle$ obtained from these as in (4.10).

Finally, with the equations

$$\begin{aligned} \mathcal{D}^\dagger \bar{\Psi}_{n_r, j, m, m'}^{\text{osc}} &= -2[(n_r+1)(n_r+2j+2)]^{1/2} \bar{\Psi}_{n_r+1, j, m, m'}^{\text{osc}}, \\ \mathcal{D} \bar{\Psi}_{n_r, j, m, m'}^{\text{osc}} &= -2[n_r(n_r+2j+1)]^{1/2} \bar{\Psi}_{n_r-1, j, m, m'}^{\text{osc}}, \end{aligned}$$

the action of the harmonic oscillator symmetry and dynamical algebra generators on $\bar{\Psi}_{n_r, j, m, m'}^{\text{osc}}(R, \theta, \lambda_1, \lambda_2)$ is completely determined.

4.2 SW system potential and dynamical potential algebras

In two dimensions, equations (2.8) and (3.8) simply lead to

$$k_1^2 = b(b-1), \quad k_2^2 = a(a-1).$$

In the following, it will prove convenient to use a and b instead of k_1 and k_2 . Up to the same phase factor as that occurring in (4.1), the extended SW Hamiltonian wavefunctions (3.9) can then be rewritten as⁴

$$\begin{aligned}\bar{\Psi}_{n_r, n, a, b}(r, \phi, \lambda_1, \lambda_2) &= \bar{\Psi}_{n_r, n}^{(a, b)}(r, \phi) (2\pi)^{-1} e^{i(b-\frac{1}{2})\lambda_1} e^{i(a-\frac{1}{2})\lambda_2}, \\ \bar{\Psi}_{n_r, n}^{(a, b)}(r, \phi) &= \mathcal{N}_{n_r, n}^{(a, b)} \mathcal{Z}_{n_r}^{(j)}(z) \Phi_n^{(a, b)}(\phi), \quad \mathcal{Z}_{n_r}^{(j)}(z) = \left(\frac{z}{\omega}\right)^{n+\frac{1}{2}(a+b)} L_{n_r}^{(2n+a+b)}(z) e^{-\frac{1}{2}z}, \\ \Phi_n^{(a, b)}(\phi) &= \cos^a \phi \sin^b \phi P_n^{(a-\frac{1}{2}, b-\frac{1}{2})}(-\cos 2\phi), \\ \mathcal{N}_{n_r, n}^{(a, b)} &= (-1)^{a+b-1} 2 \left(\frac{\omega^{2n+a+b+1} n_r! n! (2n+a+b)(n+a+b-1)!}{(n_r+2n+a+b)! (n+a-\frac{1}{2})! (n+b-\frac{1}{2})!} \right)^{1/2},\end{aligned}$$

where n , a , b are related to j , m , m' used in (4.2) through the relations⁵

$$j = n + \frac{1}{2}(a+b-1), \quad m = \frac{1}{2}(a-b), \quad m' = -\frac{1}{2}(a+b-1), \quad (4.12)$$

or, conversely,

$$a = m - m' + \frac{1}{2}, \quad b = -m - m' + \frac{1}{2}, \quad n = j + m'.$$

The generators of the $\mathfrak{su}(4)$ potential algebra, as well as those of the $\mathfrak{w}(4) \oplus_{\mathfrak{s}} \mathfrak{sp}(8, \mathbb{R})$ dynamical potential algebra, can be directly obtained by performing transformation (3.4) on the operators of Section 4.1. We get for instance⁶

$$\begin{aligned}J_0 &= \frac{i}{2}(\partial_{\lambda_1} - \partial_{\lambda_2}), \quad K_0 = \frac{i}{2}(\partial_{\lambda_1} + \partial_{\lambda_2}), \\ J_{\pm} &= \frac{1}{2} e^{\mp i(\lambda_1 - \lambda_2)} \left[\pm \partial_{\phi} - \cot \phi \left(i \partial_{\lambda_1} \pm \frac{1}{2} \right) - \tan \phi \left(i \partial_{\lambda_2} \mp \frac{1}{2} \right) \right], \\ K_{\pm} &= \frac{1}{2} e^{\mp i(\lambda_1 + \lambda_2)} \left[\mp \partial_{\phi} + \cot \phi \left(i \partial_{\lambda_1} \pm \frac{1}{2} \right) - \tan \phi \left(i \partial_{\lambda_2} \pm \frac{1}{2} \right) \right], \\ \mathcal{T}_{+1, +1} &= \frac{1}{4\omega} e^{-2i\lambda_1} \left[-\sin^2 \phi \partial_r^2 - \frac{2}{r} \sin \phi \cos \phi \partial_r^2 + \frac{2i}{r} \partial_r^2 - \frac{1}{r^2} \cos^2 \phi \partial_{\phi}^2 \right. \\ &\quad \left. + \frac{2i}{r^2} \cot \phi \partial_{\phi \lambda_1}^2 + \frac{1}{r^2} \csc^2 \phi \partial_{\lambda_1}^2 + \frac{1}{r} (1 + \sin^2 \phi) \partial_r + \frac{2}{r^2} (\cot \phi + \sin \phi \cos \phi) \partial_{\phi} \right. \\ &\quad \left. - \frac{3i}{r^2} \csc^2 \phi \partial_{\lambda_1} - \frac{5}{4r^2} \csc^2 \phi + \omega^2 r^2 \sin^2 \phi \right], \\ \mathcal{A}_{\pm \frac{1}{2}, \pm \frac{1}{2}}^{\dagger} &= \frac{1}{2} e^{\mp i\lambda_1} \left[i \left(\sin \phi \partial_r + \frac{1}{r} \cos \phi \partial_{\phi} \right) \pm \frac{1}{r} \csc \phi \partial_{\lambda_1} - \frac{i}{2r} \csc \phi - i r \sin \phi \right], \\ \mathcal{D}_{+1, +1}^{\dagger} &= \frac{1}{4\omega} e^{-2i\lambda_1} \left\{ -\sin^2 \phi \partial_r^2 - \frac{2}{r} \sin \phi \cos \phi \partial_r^2 + \frac{2i}{r} \partial_r^2 - \frac{1}{r^2} \cos^2 \phi \partial_{\phi}^2 + \frac{2i}{r^2} \cot \phi \partial_{\phi \lambda_1}^2 \right. \\ &\quad \left. + \frac{1}{r^2} \csc^2 \phi \partial_{\lambda_1}^2 + \left[\frac{1}{r} (1 + \sin^2 \phi) + 2\omega r \sin^2 \phi \right] \partial_r + 2 \left[\frac{1}{r^2} (\cot \phi + \sin \phi \cos \phi) \right. \right. \\ &\quad \left. \left. + \omega \sin \phi \cos \phi \right] \partial_{\phi} - i \left(\frac{3}{r^2} \csc^2 \phi + 2\omega \right) \partial_{\lambda_1} - \frac{5}{4r^2} \csc^2 \phi - \omega^2 r^2 \sin^2 \phi - \omega \right\},\end{aligned}$$

⁴It is worth observing here that integer or half-integer values of j , m , and m' are related to integer values of n and half-integer ones of a and b . The results for matrix elements of potential and dynamical potential algebra generators are only valid for such a and b (see footnote 3), although those for wavefunctions are not restricted to these values provided factorials are replaced by gamma functions.

⁵Equation (4.12) is valid for positive p_1 and p_2 , corresponding to physical wavefunctions (see discussion at the end of Section 3).

⁶For simplicity's sake, we denote both types of operators by the same symbols.

$$\mathcal{D}^\dagger = \frac{1}{2\omega}(-H - 2\omega r \partial_r + 2\omega^2 r^2 - 2\omega).$$

It is also straightforward to derive their matrix elements from the results of Section 4.1 and equation (4.12). We list them below:

$$\begin{aligned} J_0 \bar{\Psi}_{n_r, n, a, b} &= \frac{1}{2}(a-b) \bar{\Psi}_{n_r, n, a, b}, & K_0 \bar{\Psi}_{n_r, n, a, b} &= -\frac{1}{2}(a+b-1) \bar{\Psi}_{n_r, n, a, b}, \\ J_+ \bar{\Psi}_{n_r, n, a, b} &= \left[\left(n+a+\frac{1}{2} \right) \left(n+b-\frac{1}{2} \right) \right]^{1/2} \bar{\Psi}_{n_r, n, a+1, b-1}, \\ J_- \bar{\Psi}_{n_r, n, a, b} &= \left[\left(n+a-\frac{1}{2} \right) \left(n+b+\frac{1}{2} \right) \right]^{1/2} \bar{\Psi}_{n_r, n, a-1, b+1}, \\ K_+ \bar{\Psi}_{n_r, n, a, b} &= \left[(n+1)(n+a+b-1) \right]^{1/2} \bar{\Psi}_{n_r, n+1, a-1, b-1}, \\ K_- \bar{\Psi}_{n_r, n, a, b} &= \left[n(n+a+b) \right]^{1/2} \bar{\Psi}_{n_r, n-1, a+1, b+1}, \\ \mathcal{T}_{\sigma, \tau} \bar{\Psi}_{n_r, n, a, b} &= \sum_{n'=n+\tau-1}^{n+\tau+1} t_{n'}(n_r, 2n+a+b) \\ &\times \langle n+\frac{1}{2}(a+b-1) \frac{1}{2}(a-b), 1 \sigma | n' - \tau + \frac{1}{2}(a+b-1) \frac{1}{2}(a-b) + \sigma \rangle \\ &\times \langle n+\frac{1}{2}(a+b-1) - \frac{1}{2}(a+b-1), 1 \tau | n' - \tau + \frac{1}{2}(a+b-1) - \frac{1}{2}(a+b-1) + \tau \rangle \\ &\times \bar{\Psi}_{n_r - (n' - n - \tau), n', a + \sigma - \tau, b - \sigma - \tau}, \\ \mathcal{A}_{\sigma, \tau}^\dagger \bar{\Psi}_{n_r, n, a, b} &= \sum_{n'=n+\tau-\frac{1}{2}}^{n+\tau+\frac{1}{2}} a_{n'}(n_r, 2n+a+b) \\ &\times \langle n+\frac{1}{2}(a+b-1) \frac{1}{2}(a-b), \frac{1}{2} \sigma | n' - \tau + \frac{1}{2}(a+b-1) \frac{1}{2}(a-b) + \sigma \rangle \\ &\times \langle n+\frac{1}{2}(a+b-1) - \frac{1}{2}(a+b-1), \frac{1}{2} \tau | n' - \tau + \frac{1}{2}(a+b-1) - \frac{1}{2}(a+b-1) + \tau \rangle \\ &\times \bar{\Psi}_{n_r - (n' - n - \tau) + \frac{1}{2}, n', a + \sigma - \tau, b - \sigma - \tau}, \\ \mathcal{D}_{\sigma, \tau}^\dagger \bar{\Psi}_{n_r, n, a, b} &= \sum_{n'=n+\tau-1}^{n+\tau+1} d_{n'}(n_r, 2n+a+b) \\ &\times \langle n+\frac{1}{2}(a+b-1) \frac{1}{2}(a-b), 1 \sigma | n' - \tau + \frac{1}{2}(a+b-1) \frac{1}{2}(a-b) + \sigma \rangle \\ &\times \langle n+\frac{1}{2}(a+b-1) - \frac{1}{2}(a+b-1), 1 \tau | n' - \tau + \frac{1}{2}(a+b-1) - \frac{1}{2}(a+b-1) + \tau \rangle \\ &\times \bar{\Psi}_{n_r - (n' - n - \tau) + 1, n', a + \sigma - \tau, b - \sigma - \tau}, \\ \mathcal{D}^\dagger \bar{\Psi}_{n_r, n, a, b} &= -2[(n_r+1)(n_r+2n+a+b+1)]^{1/2} \bar{\Psi}_{n_r+1, n, a, b}. \end{aligned}$$

Here

$$\begin{aligned} t_{n'}(n_r, 2n+a+b) &= \begin{cases} - \left(\frac{(2n+a+b)n_r(n_r+2n+a+b+1)}{2n+a+b+2} \right)^{1/2} & \text{if } n' = n + \tau + 1, \\ n_r + n + \frac{1}{2}(a+b+1) & \text{if } n' = n + \tau, \\ - \left(\frac{(2n+a+b)(n_r+1)(n_r+2n+a+b)}{2n+a+b-2} \right)^{1/2} & \text{if } n' = n + \tau - 1, \end{cases} \\ a_{n'}(n_r, 2n+a+b) &= \begin{cases} i \left(\frac{(2n+a+b)(n_r+2n+a+b+1)}{2n+a+b+1} \right)^{1/2} & \text{if } n' = n + \tau + \frac{1}{2}, \\ -i \left(\frac{(2n+a+b)(n_r+1)}{2n+a+b-1} \right)^{1/2} & \text{if } n' = n + \tau - \frac{1}{2}, \end{cases} \end{aligned}$$

and

$$d_{n'}(n_r, 2n+a+b) = \begin{cases} - \left(\frac{(2n+a+b)(n_r+2n+a+b+1)(n_r+2n+a+b+2)}{2n+a+b+2} \right)^{1/2} & \text{if } n' = n + \tau + 1, \\ [(n_r+1)(n_r+2n+a+b+1)]^{1/2} & \text{if } n' = n + \tau, \\ - \left(\frac{(2n+a+b)(n_r+1)(n_r+2)}{2n+a+b-2} \right)^{1/2} & \text{if } n' = n + \tau - 1. \end{cases}$$

From these results, we conclude that the potential algebra generators produce transitions between levels belonging to spectra of Hamiltonians characterized by parameters (a, b) , $(a \pm 1, b \mp 1)$, $(a \pm 1, b \pm 1)$, $(a \pm 2, b)$, and $(a, b \pm 2)$. For the dynamical potential algebra generators, the same Hamiltonians are involved together with those associated with $(a \pm 1, b)$ and $(a, b \pm 1)$.

5 Conclusion

In the present paper, we have re-examined the D -dimensional SW system, which may be considered as the archetype of D -dimensional superintegrable system. We have completed Evans previous algebraic study, wherein its symmetry and dynamical algebras had been determined, by constructing its potential and dynamical potential algebras.

In our approach based on the use of hyperspherical coordinates in the D -dimensional space and on the introduction of D auxiliary continuous variables, the SW system has been obtained by reducing a $2D$ -dimensional harmonic oscillator Hamiltonian. The $\mathfrak{su}(2D)$ symmetry and $\mathfrak{w}(2D) \oplus_s \mathfrak{sp}(4D, \mathbb{R})$ dynamical algebras of the latter have then been transformed into corresponding potential and dynamical potential algebras for the former. Finally, the two-dimensional case has been studied in the fullest detail.

Possible connections with other approaches currently used in connection with the SW system or, more generally, superintegrable systems, such as supersymmetry [52, 53], path integrals [87], coherent states [88], and deformations [89, 90], might be interesting topics for future investigation.

A Wavefunctions of the $2D$ -dimensional harmonic oscillator

The purpose of this appendix is to derive the explicit form of the harmonic oscillator wavefunctions (2.3).

On inserting (2.3) in the Schrödinger equation (2.2), the latter separates into $D - 1$ angular equations

$$\left\{ -d_{\theta_\nu}^2 - [(2D - 2\nu - 1) \cot \theta_\nu - \tan \theta_\nu] d_{\theta_\nu} + \frac{C_{\nu+1}}{\sin^2 \theta_\nu} + \frac{p_\nu^2}{\cos^2 \theta_\nu} - C_\nu \right\} \Theta_\nu(\theta_\nu) = 0, \\ \nu = 1, 2, \dots, D - 1, \quad (\text{A.1})$$

and a radial equation

$$\left(-d_R^2 - \frac{2D - 1}{R} d_R + \frac{C_1}{R^2} + R^2 - E^{\text{osc}} \right) \mathcal{L}(z) = 0. \quad (\text{A.2})$$

Here C_1, C_2, \dots, C_{D-1} are $D - 1$ separation constants, while C_D is defined by

$$C_D = p_D^2. \quad (\text{A.3})$$

In the following, we are going to show that there does exist a solution to the whole set of D equations (A.1) and (A.2) such that all the separation constants C_ν , $\nu = 1, 2, \dots, D - 1$, are nonnegative.

Let us start by solving the angular equation (A.1) corresponding to the variable θ_ν in terms of p_ν and $C_{\nu+1}$. The ansatz

$$\Theta_\nu(\theta_\nu) = (\cos \theta_\nu)^{a_\nu - \frac{1}{2}} (\sin \theta_\nu)^{b_\nu - \frac{1}{2}} F_\nu(u_\nu), \quad u_\nu = \cos^2 \theta_\nu,$$

transforms it into the hypergeometric differential equation [84]

$$\{ u_\nu(1 - u_\nu) d_{u_\nu}^2 + [\gamma - (\alpha + \beta + 1)u_\nu] d_{u_\nu} - \alpha\beta \} F_\nu(u_\nu) = 0 \quad (\text{A.4})$$

provided we choose the constants a_ν and b_ν in such a way that

$$(a_\nu - \frac{1}{2})^2 = p_\nu^2, \quad (b_\nu - \frac{1}{2})(b_\nu + 2D - 2\nu - \frac{5}{2}) = C_{\nu+1}. \quad (\text{A.5})$$

In (A.4), α , β , and γ are given by

$$\alpha = \frac{1}{2}(a_\nu + b_\nu + D - \nu - 1 + \Delta_\nu), \quad \beta = \frac{1}{2}(a_\nu + b_\nu + D - \nu - 1 - \Delta_\nu), \quad \gamma = a_\nu + \frac{1}{2},$$

where

$$\Delta_\nu = \sqrt{(D - \nu)^2 + C_\nu}. \quad (\text{A.6})$$

There are altogether four solutions to the two quadratic equations (A.5), which may be written as

$$a_\nu = \frac{1}{2} + \epsilon|p_\nu|, \quad b_\nu = -(D - \nu - \frac{3}{2}) + \epsilon'\Delta_{\nu+1}, \quad \epsilon, \epsilon' = \pm 1. \quad (\text{A.7})$$

Consequently, we get

$$\begin{aligned} \alpha &= \frac{1}{2}(1 + \epsilon|p_\nu| + \epsilon'\Delta_{\nu+1} + \Delta_\nu), & \beta &= \frac{1}{2}(1 + \epsilon|p_\nu| + \epsilon'\Delta_{\nu+1} - \Delta_\nu), \\ \gamma &= 1 + \epsilon|p_\nu|. \end{aligned} \quad (\text{A.8})$$

The general solution of the differential equation (A.4) may be written down as

$$F_\nu(u_\nu) = A {}_2F_1(\alpha, \beta; \gamma; u_\nu) + B u_\nu^{1-\gamma} {}_2F_1(\alpha - \gamma + 1, \beta - \gamma + 1; 2 - \gamma; u_\nu),$$

where A and B are two constants to be determined so that the angular function $\Theta_\nu(\theta_\nu)$ be physically acceptable, i.e., vanish for $\theta_\nu \rightarrow 0$ and $\theta_\nu \rightarrow \frac{\pi}{2}$. On considering the four possibilities for the pair (ϵ, ϵ') in (A.7) and (A.8) successively, we arrive at a single solution corresponding either to $\epsilon = +1$, $\epsilon' = +1$, $B = 0$, $\beta = -n_\nu$ ($n_\nu \in \mathbb{N}$) or to $\epsilon = -1$, $\epsilon' = +1$, $A = 0$, $\beta - \gamma + 1 = -n_\nu$ ($n_\nu \in \mathbb{N}$). It can be expressed in terms of a Jacobi polynomial [84] as in equation (2.7), where

$$a_\nu = |p_\nu| + \frac{1}{2}, \quad b_\nu = -(D - \nu - \frac{3}{2}) + \Delta_{\nu+1}, \quad n_\nu = 0, 1, 2, \dots, \quad (\text{A.9})$$

while the separation constant C_ν must satisfy the equation

$$\Delta_\nu = 2n_\nu + |p_\nu| + \Delta_{\nu+1} + 1 \quad (\text{A.10})$$

with Δ_ν defined in (A.6).

To obtain a solution to the whole set of $D - 1$ angular equations (A.1), as expressed in equation (2.6), it only remains to solve the recursion relation (A.10) for Δ_ν with the starting value $\Delta_D = |p_D|$ corresponding to (A.3). The results for Δ_ν and C_ν , $\nu = 1, 2, \dots, D - 1$, read

$$\Delta_\nu = 2n_\nu + 2n_{\nu+1} + \dots + 2n_{D-1} + |p_\nu| + |p_{\nu+1}| + \dots + |p_D| + D - \nu$$

and

$$\begin{aligned} C_\nu &= (2n_\nu + 2n_{\nu+1} + \dots + 2n_{D-1} + |p_\nu| + |p_{\nu+1}| + \dots + |p_D|) \\ &\quad \times (2n_\nu + 2n_{\nu+1} + \dots + 2n_{D-1} + |p_\nu| + |p_{\nu+1}| + \dots + |p_D| + 2D - 2\nu), \end{aligned} \quad (\text{A.11})$$

respectively. As a consequence, b_ν in (A.9) can be rewritten as in equation (2.8). This completes the proof of equations (2.6) to (2.8).

Turning now ourselves to the radial equation (A.2), we note from (A.11) that C_1 can be written as

$$C_1 = 4j(j + D - 1)$$

in terms of j defined in (2.10). Finally, it is straightforward to show that the physically acceptable solutions vanishing for r (or z) going to zero and infinity are given by (2.9) and correspond to the eigenvalues (2.11).

References

- [1] Goldstein H., Classical mechanics, 2nd ed., *Addison-Wesley Series in Physics*, Addison-Wesley Publishing Co., Reading, Mass., 1980.
- [2] Pauli W., Über das Wasserstoffspektrum von Standpunkt der neuen Quantenmechanik, *Z. Phys.* **36** (1926), 336–363.
- [3] Fock V., Zur Theorie des Wasserstoffatoms, *Z. Phys.* **98** (1935), 145–154.
- [4] Bargmann V., Zur Theorie des Wasserstoffatoms. Bemerkungen zur gleichnamigen Arbeit von V. Fock, *Z. Phys.* **99** (1936), 576–582.
- [5] Jauch J.M., Hill E.L., On the problem of degeneracy in quantum mechanics, *Phys. Rev.* **57** (1940), 641–645.
- [6] Moshinsky M., Smirnov Yu.F., The harmonic oscillator in modern physics, *Contemporary Concepts in Physics*, Vol. 9, Harwood, Amsterdam, 1996.
- [7] Fris I., Mandrosov V., Smorodinsky Ya.A., Uhlir M., Winternitz P., On higher symmetries in quantum mechanics, *Phys. Lett.* **16** (1965), 354–356.
- [8] Winternitz P., Smorodinsky Ya.A., Uhlir M., Fris I., Symmetry groups in classical and quantum mechanics, *Soviet J. Nuclear Phys.* **4** (1967), 444–450.
- [9] Makarov A.A., Smorodinsky Ya.A., Valiev Kh., Winternitz P., A systematic search for non-relativistic system with dynamical symmetries, *Nuovo Cim. A* **52** (1967), 1061–1084.
- [10] Evans N.W., Superintegrability in classical mechanics, *Phys. Rev. A* **41** (1990), 5666–5676.
- [11] Kalnins E.G., Kress J.M., Miller W. Jr., Second-order superintegrable systems in conformally flat spaces. I. Two-dimensional classical structure theory, *J. Math. Phys.* **46** (2005), 053509, 28 pages.
- [12] Kalnins E.G., Kress J.M., Miller W. Jr., Second order superintegrable systems in conformally flat spaces. II. The classical two-dimensional Stäckel transform, *J. Math. Phys.* **46** (2005), 053510, 15 pages.
- [13] Kalnins E.G., Kress J.M., Miller W. Jr., Second order superintegrable systems in conformally flat spaces. III. Three-dimensional classical structure, *J. Math. Phys.* **46** (2005), 103507, 28 pages.
- [14] Kalnins E.G., Kress J.M., Miller W. Jr., Second order superintegrable systems in conformally flat spaces. IV. The classical 3D Stäckel transform and 3D classification theory, *J. Math. Phys.* **47** (2006), 043514, 26 pages.
- [15] Kalnins E.G., Kress J.M., Miller W. Jr., Second-order superintegrable systems in conformally flat spaces. V. Two- and three-dimensional quantum systems, *J. Math. Phys.* **47** (2006), 093501, 25 pages.
- [16] Daskaloyannis C., Ypsilantis K., Unified treatment and classification of superintegrable systems with integrals quadratic in momenta on a two dimensional manifold, *J. Math. Phys.* **47** (2006), 042904, 38 pages, [math-ph/0412055](#).
- [17] Daskaloyannis C., Tanoudis Y., Quantum superintegrable systems with quadratic integrals on a two dimensional manifold, *J. Math. Phys.* **48** (2007), 072108, 22 pages, [math-ph/0607058](#).
- [18] Drach J., Sur l'intégration logique des équations de la dynamique à deux variables: Forces conservatrices. Intégrales cubiques. Mouvements dans le plan, *C. R. Séances Acad. Sci. III* **200** (1935), 22–26.
- [19] Drach J., Sur l'intégration logique et sur la transformation des équations de la dynamique à deux variables: Forces conservatrices. Intégrales cubiques, *C. R. Séances Acad. Sci. III* **200** (1935), 599–602.
- [20] Gravel S., Winternitz P., Superintegrability with third-order invariants in quantum and classical mechanics, *J. Math. Phys.* **43** (2002), 5902–5912, [math-ph/0206046](#).
- [21] Gravel S., Hamiltonians separable in Cartesian coordinates and third-order integrals of motion, *J. Math. Phys.* **45** (2004), 1003–1019, [math-ph/0302028](#).
- [22] Verrier P.E., Evans N.W., A new superintegrable Hamiltonian, *J. Math. Phys.* **49** (2008), 022902, 8 pages, [arXiv:0712.3677](#).
- [23] Evans N.W., Verrier P.E., Superintegrability of the caged anisotropic oscillator, *J. Math. Phys.* **49** (2008), 092902, 10 pages, [arXiv:0808.2146](#).
- [24] Rodríguez M.A., Tempesta P., Winternitz P., Reduction of superintegrable systems: the anisotropic harmonic oscillator, *Phys. Rev. E* **78** (2008), 046608, 6 pages, [arXiv:0809.3259](#).
- [25] Marquette I., Superintegrability with third order integrals of motion, cubic algebras, and supersymmetric quantum mechanics. I. Rational function potentials, *J. Math. Phys.* **50** (2009), 012101, 23 pages, [arXiv:0807.2858](#).

- [26] Marquette I., Superintegrability with third order integrals of motion, cubic algebras, and supersymmetric quantum mechanics. II. Painlevé transcendental potentials, *J. Math. Phys.* **50** (2009), 095202, 18 pages, [arXiv:0811.1568](#).
- [27] Tremblay F., Turbiner A.V., Winternitz P., An infinite family of solvable and integrable quantum systems on a plane, *J. Phys. A: Math. Theor.* **42** (2009), 242001, 10 pages, [arXiv:0904.0738](#).
- [28] Quesne C., Superintegrability of the Tremblay–Turbiner–Winternitz quantum Hamiltonians on a plane for odd k , *J. Phys. A: Math. Theor.* **43** (2010), 082001, 10 pages, [arXiv:0911.4404](#).
- [29] Kalnins E.G., Kress J.M., Miller W. Jr., Superintegrability and higher order integrals for quantum systems, *J. Phys. A: Math. Theor.* **43** (2010), 265205, 21 pages.
- [30] Kalnins E.G., Kress J.M., Miller W. Jr., Tools for verifying classical and quantum superintegrability, *SIGMA* **6** (2010), 066, 23 pages, [arXiv:1006.0864](#).
- [31] Post S., Winternitz P., An infinite family of superintegrable deformations of the Coulomb potential, *J. Phys. A: Math. Theor.* **43** (2010), 222001, 11 pages, [arXiv:1003.5230](#).
- [32] Kalnins E.G., Kress J.M., Miller W. Jr., A recurrence relation approach to higher order quantum superintegrability, *SIGMA* **7** (2011), 031, 24 pages, [arXiv:1011.6548](#).
- [33] Evans N.W., Super-integrability of the Winternitz system, *Phys. Lett. A* **147** (1990), 483–486.
- [34] Evans N.W., Group theory of the Smorodinsky–Winternitz system, *J. Math. Phys.* **32** (1991), 3369–3375.
- [35] Granovskii Ya.I., Lutzenko I.M., Zhedanov A.S., Mutual integrability, quadratic algebras, and dynamical symmetry, *Ann. Physics* **217** (1992), 1–20.
- [36] Granovskii Ya.I., Zhedanov A.S., Lutsenko I.M., Quadratic algebras and dynamics in curved space. I. Oscillator, *Theoret. and Math. Phys.* **91** (1992), 474–480.
- [37] Granovskii Ya.I., Zhedanov A.S., Lutsenko I.M., Quadratic algebras and dynamics in curved spaces. II. The Kepler problem, *Theoret. and Math. Phys.* **91** (1992), 604–612.
- [38] Daskaloyannis C., Quadratic Poisson algebras of two-dimensional classical superintegrable systems and quadratic associative algebras of quantum superintegrable systems, *J. Math. Phys.* **42** (2001), 1100–1119, [math-ph/0003017](#).
- [39] Daskaloyannis C., Generalized deformed oscillator and nonlinear algebras, *J. Phys. A: Math. Gen.* **24** (1991), L789–L794.
- [40] Quesne C., Generalized deformed parafermions, nonlinear deformations of $so(3)$ and exactly solvable potentials, *Phys. Lett. A* **193** (1994), 245–250.
- [41] Cooper F., Khare A., Sukhatme U., Supersymmetry and quantum mechanics, *Phys. Rep.* **251** (1995), 267–385, [hep-th/9405029](#).
- [42] Junker G., Supersymmetric methods in quantum and statistical physics, *Texts and Monographs in Physics*, Springer-Verlag, Berlin, 1996.
- [43] Andrianov A.A., Ioffe M.V., Cannata F., Dedonder J.-P., Second order derivative supersymmetry, q deformations and the scattering problem, *Internat. J. Modern Phys. A* **10** (1995), 2683–2702, [hep-th/9404061](#).
- [44] Andrianov A.A., Ioffe M.V., Nishnianidze D.N., Polynomial supersymmetry and dynamical symmetries in quantum mechanics, *Theoret. and Math. Phys.* **104** (1995), 1129–1140.
- [45] Andrianov A.A., Ioffe M.V., Nishnianidze D.N., Polynomial SUSY in quantum mechanics and second derivative Darboux transformations, *Phys. Lett. A* **201** (1995), 103–110, [hep-th/9404120](#).
- [46] Samsonov B.F., New features in supersymmetry breakdown in quantum mechanics, *Modern Phys. Lett. A* **11** (1996), 1563–1567, [quant-ph/9611012](#).
- [47] Bagchi B., Ganguly A., Bhaumik D., Mitra A., Higher derivative supersymmetry, a modified Crum–Darboux transformation and coherent state, *Modern Phys. Lett. A* **14** (1999), 27–34.
- [48] Plyushchay M., Hidden nonlinear supersymmetries in pure parabosonic systems, *Internat. J. Modern Phys. A* **15** (2000), 3679–3698, [hep-th/9903102](#).
- [49] Klishevich S., Plyushchay M., Nonlinear supersymmetry, quantum anomaly and quasi-exactly solvable systems, *Nuclear Phys. B* **606** (2001), 583–612, [cond-mat/0007461](#).
- [50] Aoyama H., Sato M., Tanaka T., \mathcal{N} -fold supersymmetry in quantum mechanics: general formalism, *Nuclear Phys. B* **619** (2001), 105–127, [quant-ph/0106037](#).
- [51] Fernández C. D.J., Fernández-García N., Higher-order supersymmetric quantum mechanics, *AIP Conf. Proc.* **744** (2005), 236–273, [quant-ph/0502098](#).

- [52] Marquette I., Supersymmetry as a method of obtaining new superintegrable systems with higher order integrals of motion, *J. Math. Phys.* **50** (2009), 122102, 10 pages, [arXiv:0908.1246](https://arxiv.org/abs/0908.1246).
- [53] Marquette I., Superintegrability and higher order polynomial algebras, *J. Phys. A: Math. Theor.* **43** (2010), 135203, 15 pages, [arXiv:0908.4399](https://arxiv.org/abs/0908.4399).
- [54] Turbiner A.V., Quasi-exactly-solvable problems and $sl(2)$ algebra, *Comm. Math. Phys.* **118** (1988), 467–474.
- [55] Shifman M.A., Turbiner A.V., Quantal problems with partial algebraization of the spectrum, *Comm. Math. Phys.* **126** (1989), 347–365.
- [56] Ushveridze A.G., Quasi-exactly solvable models in quantum mechanics, *Soviet J. Particles and Nuclei* **20** (1989), 504–528.
- [57] Tempesta P., Turbiner A.V., Winternitz P., Exact solvability of superintegrable systems, *J. Math. Phys.* **42** (2001), 4248–4257, [hep-th/0011209](https://arxiv.org/abs/hep-th/0011209).
- [58] Kalnins E.G., Miller W. Jr., Pogosyan G.S., Exact and quasixact solvability of second-order superintegrable quantum systems. I. Euclidean space preliminaries, *J. Math. Phys.* **47** (2006), 033502, 30 pages.
- [59] Kalnins E.G., Miller W. Jr., Pogosyan G.S., Exact and quasixact solvability of second-order superintegrable quantum systems. II. Relation to separation of variables, *J. Math. Phys.* **48** (2007), 023503, 20 pages.
- [60] Demkov Yu.N., The definition of the symmetry group of a quantum system. The anisotropic oscillator, *Soviet Phys. JETP* **17** (1963), 1349–1351.
- [61] Dothan Y., Finite-dimensional spectrum-generating algebras, *Phys. Rev. D* **2** (1970), 2944–2954.
- [62] Bonatsos D., Kolokotronis P., Lenis D., Daskaloyannis C., Deformed $u(2)$ algebra as the symmetry algebra of the planar anisotropic quantum harmonic oscillator with rational ratio of frequencies, *Internat. J. Modern Phys. A* **12** (1997), 3335–3346.
- [63] Dothan Y., Gell-Mann M., Ne’eman Y., Series of hadron energy levels as representations of non-compact groups, *Phys. Lett.* **17** (1965), 148–151.
- [64] Mukunda M., O’Raifeartaigh L., Sudarshan E.C.G., Characteristic noninvariance groups of dynamical systems, *Phys. Rev. Lett.* **15** (1965), 1041–1044.
- [65] Barut A.O., Böhm A., Dynamical groups and mass formula, *Phys. Rev.* **139** (1965), B1107–B1112.
- [66] Alhassid Y., Gürsey F., Iachello F., Group theory approach to scattering, *Ann. Physics* **148** (1983), 346–380.
- [67] Alhassid Y., Gürsey F., Iachello F., Group theory approach to scattering. II. The Euclidean connection, *Ann. Physics* **167** (1986), 181–200.
- [68] Frank A., Wolf K.B., Lie algebras for potential scattering, *Phys. Rev. Lett.* **52** (1984), 1737–1739.
- [69] Quesne C., An $sl(4, \mathbb{R})$ Lie algebraic treatment of the first family of Pöschl–Teller potentials, *J. Phys. A: Math. Gen.* **21** (1988), 4487–4500.
- [70] Quesne C., $so(3,1)$ versus $sp(4, \mathbb{R})$ as dynamical potential algebra of the symmetrical Pöschl–Teller potentials, *J. Phys. A: Math. Gen.* **21** (1988), 4501–4511.
- [71] Quesne C., An $sl(4, \mathbb{R})$ Lie algebraic approach to the Bargmann functions and its application to the second Pöschl–Teller equation, *J. Phys. A: Math. Gen.* **22** (1989), 3723–3730.
- [72] Kuru Ş., Negro J., Dynamical algebras for Pöschl–Teller Hamiltonian hierarchies, *Ann. Physics* **324** (2009), 2548–2560.
- [73] Correa F., Jakubský V., Plyushchay M.S., Aharonov–Bohm effect on AdS_2 and nonlinear supersymmetry of reflectionless Pöschl–Teller system, *Ann. Physics* **324** (2009), 1078–1094, [arXiv:0806.1614](https://arxiv.org/abs/0806.1614).
- [74] Del Sol Mesa A., Quesne C., Smirnov Yu.F., Generalized Morse potential: symmetry and satellite potentials, *J. Phys. A: Math. Gen.* **31** (1998), 321–335, [physics/9708004](https://arxiv.org/abs/physics/9708004).
- [75] Del Sol Mesa A., Quesne C., Connection between type A and E factorizations and construction of satellite algebras, *J. Phys. A: Math. Gen.* **33** (2000), 4059–4071, [math-ph/0004027](https://arxiv.org/abs/math-ph/0004027).
- [76] Kerimov G.A., Non-central potentials related to the Lie algebra $u(4)$, *Phys. Lett. A* **358** (2006), 176–180.
- [77] Kerimov G.A., Algebraic approach to non-central potentials, *J. Phys. A: Math. Gen.* **39** (2006), 1183–1189.
- [78] Kerimov G.A., Quantum scattering from the Coulomb potential plus an angle-dependent potential: a group-theoretical study, *J. Phys. A: Math. Theor.* **40** (2007), 7297–7308.
- [79] Kerimov G.A., Non-spherically symmetric transparent potentials for the three-dimensional Schrödinger equation, *J. Phys. A: Math. Theor.* **40** (2007), 11607–11615.

-
- [80] Kerimov G.A., Ventura A., Group-theoretical approach to a non-central extension of the Kepler–Coulomb problem, *J. Phys. A: Math. Theor.* **43** (2010), 255304, 10 pages, [arXiv:1005.1215](#).
- [81] Calzada J.A., Negro J., del Olmo M.A., Superintegrable quantum $u(3)$ systems and higher rank factorizations, *J. Math. Phys.* **47** (2006), 043511, 17 pages, [math-ph/0601067](#).
- [82] Calzada J.A., Kuru Ş., Negro J., del Olmo M.A., Intertwining symmetry algebras of quantum superintegrable systems on the hyperboloid, *J. Phys. A: Math. Theor.* **41** (2008), 255201, 11 pages, [arXiv:0803.2117](#).
- [83] Calzada J.A., Negro J., del Olmo M.A., Intertwining symmetry algebras of quantum superintegrable systems, *SIGMA* **5** (2009), 039, 23 pages, [arXiv:0904.0170](#).
- [84] Gradshteyn I.S., Ryzhik I.M., Table of integrals, series, and products, Academic Press, New York, 1980.
- [85] Biedenharn L.C., Louck J.D., Angular momentum in quantum physics. Theory and application, *Encyclopedia of Mathematics and its Applications*, Vol. 8, Addison-Wesley Publishing Co., Reading, Mass., 1981.
- [86] Schneider C.K.E., Wilson R., Ladder operators of group matrix elements, *J. Math. Phys.* **20** (1979), 2380–2390.
- [87] Grosche C., Pogosyan G.S., Sissakian A.N., Path integral discussion for Smorodinsky–Winternitz potentials. I. Two- and three-dimensional Euclidean space, *Fortschr. Phys.* **43** (1995), 453–521, [hep-th/9402121](#).
- [88] Ünal N., Coherent states for Smorodinsky–Winternitz potentials, *Cent. Eur. J. Phys.* **7** (2009), 774–785.
- [89] Herranz F.J., Ballesteros A., Superintegrability on three-dimensional Riemannian and relativistic spaces of constant curvature, *SIGMA* **2** (2006), 010, 22 pages, [math-ph/0512084](#).
- [90] Cariñena J.F., Rañada M.F., Santander M., A super-integrable two-dimensional non-linear oscillator with an exactly solvable quantum analog, *SIGMA* **3** (2007), 030, 23 pages, [math-ph/0702084](#).