

New Combinatorial Formulae for Nested Bethe Vectors

Maksim KOSMAKOV^a and Vitaly TARASOV^b

^{a)} *Department of Mathematical Sciences, University of Cincinnati,
P.O. Box 210025, Cincinnati, OH 45221, USA*
E-mail: kosmakmm@ucmail.uc.edu

^{b)} *Department of Mathematical Sciences, Indiana University Indianapolis,
402 North Blackford St, Indianapolis, IN 46202-3216, USA*
E-mail: vtarasov@iu.edu

Received January 08, 2025, in final form July 08, 2025; Published online July 22, 2025

<https://doi.org/10.3842/SIGMA.2025.060>

Abstract. We give new combinatorial formulae for vector-valued weight functions (off-shell nested Bethe vectors) for the evaluation modules over the Yangian $Y(\mathfrak{gl}_4)$. The case of $Y(\mathfrak{gl}_n)$ for an arbitrary n is considered in [Lett. Math. Phys. **115** (2025), 12, 20 pages, arXiv:2402.15717].

Key words: Bethe ansatz; Yangian; weight functions

2020 Mathematics Subject Classification: 17B37; 81R50; 82B23

1 Introduction

In this paper we will give new combinatorial formulae for vector-valued weight functions for evaluation modules over the Yangian $Y(\mathfrak{gl}_n)$. The weight functions, also known as (off-shell) nested Bethe vectors, play an important role in the theory of quantum integrable models and representation theory of Lie algebras and quantum groups. Initially, they appeared in the framework of the nested algebraic Bethe ansatz as a tool to find eigenvectors and eigenvalues of transfer matrices of lattice integrable models associated with higher rank Lie algebras [6, 7], see [21, 22] for a review of the algebraic Bethe ansatz. The results of [6] has been extended to higher transfer matrices in [11].

Furthermore, the vector-valued weight functions were used to construct hypergeometric solutions of the quantized (difference) Knizhnik–Zamolodchikov equations [10, 28]. They also showed up in several related problems [3, 12, 24, 26]. In a more recent development, the weight functions were connected to the stable envelopes for particular Nakajima quiver varieties, the cotangent bundles of partial flag varieties [15, 16, 17, 18, 27].

For various applications, it is important to have expressions for vector-valued weight functions for tensor products of evaluation modules over $Y(\mathfrak{gl}_n)$. Such expressions can be obtained in two steps. The first step is to consider weight functions for a single evaluation module, and the second step is to combine expressions for individual evaluation modules into an expression for the whole tensor product. In this paper, we will focus on the first step. The second step is fairly standard and is not specifically discussed here.

By definition, an evaluation $Y(\mathfrak{gl}_n)$ -module is a \mathfrak{gl}_n -module equipped with the action of $Y(\mathfrak{gl}_n)$ via the evaluation homomorphism $Y(\mathfrak{gl}_n) \rightarrow U(\mathfrak{gl}_n)$, see Section 2. The goal is to expand the vector-valued weight function for the evaluation $Y(\mathfrak{gl}_n)$ -module in a basis coming from the representation theory of \mathfrak{gl}_n and find expressions for the coordinates. For Verma modules

over \mathfrak{gl}_n , such kind of expressions are given in [25]. In this paper, we give a generalization of formulae from [25].

Combinatorial formulae for the vector-valued weight functions associated with the differential Knizhnik–Zamolodchikov equations were developed in [1, 8, 9, 14, 19, 20].

The expressions for weight functions in [25] are based on recursions induced by the standard embeddings of Lie algebras, $\mathfrak{gl}_1 \oplus \mathfrak{gl}_{n-1} \subset \mathfrak{gl}_n$ and $\mathfrak{gl}_{n-1} \oplus \mathfrak{gl}_1 \subset \mathfrak{gl}_n$. The recursions allow one to write down weight functions for $Y(\mathfrak{gl}_n)$ via weight functions for $Y(\mathfrak{gl}_{n-1})$. This results in formulae for coordinates of weight functions in bases of Verma \mathfrak{gl}_n -modules of the form

$$\left\{ \prod_{i>j} e_{ij}^{m_{ij}} v, m_{ij} \in \mathbb{Z}_{\geq 0} \right\}, \quad (1.1)$$

where e_{ij} are the standard generators of \mathfrak{gl}_n , see (2.7), v is the highest weight vector, and some ordering of noncommuting factors is imposed. The ordering is determined by the in-between part of the involved chain of embeddings $\mathfrak{gl}_1 \oplus \cdots \oplus \mathfrak{gl}_1 \subset \cdots \subset \mathfrak{gl}_n$. For instance, the chain

$$\mathfrak{gl}_1 \oplus \cdots \oplus \mathfrak{gl}_1 \subset \cdots \subset \mathfrak{gl}_{n-2} \oplus \mathfrak{gl}_1 \oplus \mathfrak{gl}_1 \subset \mathfrak{gl}_{n-1} \oplus \mathfrak{gl}_1 \subset \mathfrak{gl}_n$$

yields the ordering

$$e_{ij}^{\otimes} \text{ is to the left of } e_{kl}^{\otimes} \text{ if } i > k \text{ or } i = k, j > l, \quad (1.2)$$

while the chain

$$\mathfrak{gl}_1 \oplus \cdots \oplus \mathfrak{gl}_1 \subset \cdots \subset \mathfrak{gl}_1 \oplus \mathfrak{gl}_1 \oplus \mathfrak{gl}_{n-2} \subset \mathfrak{gl}_1 \oplus \mathfrak{gl}_{n-1} \subset \mathfrak{gl}_n$$

yields the ordering

$$e_{ij}^{\otimes} \text{ is to the left of } e_{kl}^{\otimes} \text{ if } j < l \text{ or } j = l, i < k. \quad (1.3)$$

For example for $n = 4$, the product $e_{43}e_{42}e_{41}e_{32}e_{31}e_{21}$ obeys ordering (1.2), while the product $e_{21}e_{31}e_{41}e_{32}e_{42}e_{43}$ obeys ordering (1.3).

However, some natural orderings of noncommuting factors in (1.1) important for applications do not show up in the formulae established in [25], see, for instance, [8]. The first nontrivial example occurs at $n = 4$ and is given by the basis

$$\{e_{32}^{m_{32}} e_{31}^{m_{31}} e_{42}^{m_{42}} e_{41}^{m_{41}} e_{21}^{m_{21}} e_{43}^{m_{43}} v, m_{ij} \in \mathbb{Z}_{\geq 0}\}. \quad (1.4)$$

To make the set of covered orderings wider, one can consider recursions based on more general embeddings

$$\mathfrak{gl}_m \oplus \mathfrak{gl}_{n-m} \subset \mathfrak{gl}_n \quad \text{with } 1 < m < n - 1. \quad (1.5)$$

For instance, the embedding $\mathfrak{gl}_2 \oplus \mathfrak{gl}_2 \subset \mathfrak{gl}_4$ yields example (1.4). In this paper, we will work out example (1.4) in detail with the main result given by Theorem 5.7. We consider the general case in [5].

We would like to present the \mathfrak{gl}_4 case separately in order to explain calculations more clearly without introducing too cumbersome notation and to make the exposition paper more accessible. For the same purpose, we show explicitly intermediate steps in the proofs that commonly might be tacit for the sake of making a paper shorter. In particular, we give in Appendix A a proof of Proposition 5.1. Although this statement has a long history, going back to [4], numerous applications, and is explained in several lecture courses, see [22], its straightforward proof is not easily available in the literature.

At the same time we point out that the proof of Theorem 5.7 in this paper extends almost in a straightforward way to the proof of [5, Theorem 5.5] in the general \mathfrak{gl}_n case. In particular, the proof of the key Proposition 6.3 in [5] is literally the same as the proof of Proposition 5.6 in this paper.

Unlike [25], we will consider only the case of weight functions for Yangian modules (the rational case). It turns out that dealing with weight functions for modules over the quantum loop algebra $U_q(\widetilde{\mathfrak{gl}_n})$, the trigonometric case, hits an obstacle of essential noncommutativity of q -analogues of the generators e_{ij} , $i > j$. This obstacle does not show up for the embeddings $\mathfrak{gl}_1 \oplus \mathfrak{gl}_{n-1} \subset \mathfrak{gl}_n$ and $\mathfrak{gl}_{n-1} \oplus \mathfrak{gl}_1 \subset \mathfrak{gl}_n$ explored in [25], but reveals itself for embeddings (1.5). For instance, the obstacle in example (1.4) comes from the relation

$$e_{42}e_{31} - e_{31}e_{42} = (q - q^{-1})e_{32}e_{41}$$

that holds in the trigonometric case.

There is an alternative approach to get explicit expressions for the vector-valued weight functions in the trigonometric case, see [2, 3, 13], based on considering composed currents and half-currents in the quantum affine algebra and their projections on two Borel subalgebras of different kind. This approach allows one to recover combinatorial expression for vector-valued weight functions in evaluation modules in the trigonometric case obtained in [25]. It is an interesting open question whether the composed currents approach can be helpful to obtain trigonometric analogues of new combinatorial expressions for vector-valued weight functions developed in this paper.

2 Notations

We will be using the standard superscript notation for embeddings of tensor factors into tensor products. For a tensor product of vector spaces $V_1 \otimes V_2 \otimes \cdots \otimes V_k$ and an operator $A \in \text{End}(V_i)$, denote

$$A^{(i)} = 1^{\otimes(i-1)} \otimes A \otimes 1^{\otimes(k-i)} \in \text{End}(V_1 \otimes V_2 \otimes \cdots \otimes V_k).$$

Also, if $B \in \text{End}(V_j)$, $i \neq j$, denote $(A \otimes B)^{(ij)} = A^{(i)}B^{(j)}$, etc.

Fix a positive integer n . All over the paper we identify elements of $\text{End } \mathbb{C}^n$ with $n \times n$ matrices using the standard basis of \mathbb{C}^n . That is, for $L \in \text{End } \mathbb{C}^n$ we have $L = (L_b^a)_{a,b=1}^n$, where L_b^a are the entries of L . Entries of matrices acting in the tensor products $(\mathbb{C}^n)^{\otimes k}$ are naturally labeled by multiindices. For instance, if $M \in \text{End}(\mathbb{C}^n \otimes \mathbb{C}^n)$, then $M = (M_{cd}^{ab})_{a,b,c,d=1}^n$.

The rational R -matrix is $R(u) \in \text{End}(\mathbb{C}^n \otimes \mathbb{C}^n)$,

$$R(u) = 1 + \frac{1}{u} \sum_{a,b=1}^n E_{ab} \otimes E_{ba}, \quad (2.1)$$

where $E_{ab} \in \text{End}(\mathbb{C}^n)$ is the matrix with the only nonzero entry equal to 1 at the intersection of the a -th row and b -th column. The entries of $R(u)$ are

$$R_{cd}^{ab}(u) = \delta_{ac}\delta_{bd} + \frac{1}{u}\delta_{ad}\delta_{bc}.$$

The R -matrix satisfies the Yang–Baxter equation

$$R^{(12)}(u-v)R^{(13)}(u)R^{(23)}(v) = R^{(23)}(v)R^{(13)}(u)R^{(12)}(u-v). \quad (2.2)$$

The Yangian $Y(\mathfrak{gl}_n)$ is a unital associative algebra with generators $(T_b^a)^{\{s\}}$, $a, b = 1, \dots, n$, and $s = 1, 2, \dots$. Organize them into generating series

$$T_b^a(u) = \delta_{ab} + \sum_{s=1}^{\infty} (T_b^a)^{\{s\}} u^{-s}, \quad a, b = 1, \dots, n. \quad (2.3)$$

The defining relations in $Y(\mathfrak{gl}_n)$ are

$$(u - v)[T_b^a(u), T_d^c(v)] = T_d^a(u)T_b^c(v) - T_d^a(v)T_b^c(u) \quad (2.4)$$

for all $a, b, c, d = 1, \dots, n$.

Combine series (2.3) into a matrix $T(u) = \sum_{a,b=1}^n E_{ab} \otimes T_b^a(u)$ with entries in $Y(\mathfrak{gl}_n)$. Then relations (2.4) amount to the following equality:

$$R^{(12)}(u - v)T^{(1)}(u)T^{(2)}(v) = T^{(2)}(v)T^{(1)}(u)R^{(12)}(u - v),$$

where $T^{(1)}(u) = \sum_{a,b=1}^n E_{ab} \otimes 1 \otimes T_b^a(u)$ and $T^{(2)}(v) = \sum_{a,b=1}^n 1 \otimes E_{ab} \otimes T_b^a(v)$.

The Yangian $Y(\mathfrak{gl}_n)$ is a Hopf algebra. In terms of generating series (2.3), the coproduct $\Delta: Y(\mathfrak{gl}_n) \rightarrow Y(\mathfrak{gl}_n) \otimes Y(\mathfrak{gl}_n)$ reads as follows:

$$\Delta(T_b^a(u)) = \sum_{c=1}^n T_b^c(u) \otimes T_c^a(u), \quad a, b = 1, \dots, n. \quad (2.5)$$

Denote by $\tilde{\Delta}: Y(\mathfrak{gl}_n) \rightarrow Y(\mathfrak{gl}_n) \otimes Y(\mathfrak{gl}_n)$ the opposite coproduct

$$\tilde{\Delta}(T_b^a(u)) = \sum_{c=1}^n T_c^a(u) \otimes T_b^c(u), \quad a, b = 1, \dots, n. \quad (2.6)$$

There is a one-parameter family of automorphisms $\rho_x: Y(\mathfrak{gl}_n) \rightarrow Y(\mathfrak{gl}_n)$ defined in terms of the series $T(u)$ by the rule $\rho_x T(u) = T(u - x)$, where in the right-hand side, each expression $(u - x)^{-s}$ has to be expanded as a power series in u^{-1} .

Denote by e_{ab} , $a, b = 1, \dots, n$, the standard generators of the Lie algebra \mathfrak{gl}_n ,

$$[e_{ab}, e_{cd}] = e_{ad}\delta_{bc} - e_{cb}\delta_{ad}. \quad (2.7)$$

A vector v in a \mathfrak{gl}_n -module is called singular of weight $(\Lambda^1, \dots, \Lambda^n)$ if $e_{ab}v = 0$ for all $a < b$ and $e_{aa}v = \Lambda^a v$ for all $a = 1, \dots, n$.

The Yangian $Y(\mathfrak{gl}_n)$ contains the universal enveloping algebra $U(\mathfrak{gl}_n)$ as a Hopf subalgebra. The embedding is given by the rule $e_{ab} \mapsto (T_a^b)^{\{1\}}$ for all $a, b = 1, \dots, n$. We identify $U(\mathfrak{gl}_n)$ with its image in $Y(\mathfrak{gl}_n)$ under this embedding.

The evaluation homomorphism $\epsilon: Y(\mathfrak{gl}_n) \rightarrow U(\mathfrak{gl}_n)$ is given by the rule $\epsilon: (T_b^a)(u) \mapsto \delta_{ab} + e_{ba}u^{-1}$ for all $a, b = 1, \dots, n$. Both the automorphisms ρ_x and the homomorphism ϵ restricted to the subalgebra $U(\mathfrak{gl}_n)$ are the identity maps.

For a \mathfrak{gl}_n -module V , denote by $V(x)$ the $Y(\mathfrak{gl}_n)$ -module induced from V by the homomorphism $\epsilon \circ \rho_x$. The module $V(x)$ is called an evaluation module over $Y(\mathfrak{gl}_n)$.

A vector v in a $Y(\mathfrak{gl}_n)$ -module is called singular with respect to the action of $Y(\mathfrak{gl}_n)$ if $T_b^a(u)v = 0$ for all $1 \leq b < a \leq n$. A singular vector v that is an eigenvector for the action of $T_1^1(u), \dots, T_n^n(u)$ is called a weight singular vector, and the respective eigenvalues are denoted by $\langle T_1^1(u)v \rangle, \dots, \langle T_n^n(u)v \rangle$.

Example. Let V be a \mathfrak{gl}_n -module and $v \in V$ be a \mathfrak{gl}_n -singular vector of weight $(\Lambda^1, \dots, \Lambda^n)$. Then v is a weight singular vector with respect to the action of $Y(\mathfrak{gl}_n)$ in the evaluation module $V(x)$ and $\langle T_a^a(u)v \rangle = 1 + \Lambda^a(u - x)^{-1}$, $a = 1, \dots, n$.

For $k < n$, we consider two embeddings of the algebra $Y(\mathfrak{gl}_k)$ into $Y(\mathfrak{gl}_n)$, called ϕ_k and ψ_k :

$$\phi_k(T^{(k)}(u))_b^a = (T^{(n)}(u))_b^a, \quad \psi_k(T^{(k)}(u))_b^a = (T^{(n)}(u))_{b+n-k}^{a+n-k}(u), \quad (2.8)$$

with $a, b = 1, \dots, k$. Here $(T^{(k)}(u))_b^a$ and $(T^{(n)}(u))_b^a$ are series $T_b^a(u)$ for the algebras $Y(\mathfrak{gl}_k)$ and $Y(\mathfrak{gl}_n)$, respectively.

3 Combinatorial formulae for rational weight functions

Fix a collection of nonnegative integers $\xi_1, \xi_2, \dots, \xi_{n-1}$. Set $\boldsymbol{\xi} = (\xi_1, \xi_2, \dots, \xi_{n-1})$ and $\xi^a = \xi_1 + \dots + \xi_a$, $a = 1, \dots, n-1$. Consider the variables t_i^a , $a = 1, \dots, n-1$, $i = 1, \dots, \xi_a$. We will also write

$$\mathbf{t}^a = (t_1^a, \dots, t_{\xi_a}^a), \quad \mathbf{t} = (\mathbf{t}^1, \dots, \mathbf{t}^{n-1}).$$

We will use the ordered product notation for any noncommuting factors X_1, \dots, X_k ,

$$\prod_{1 \leq i \leq k}^{\rightarrow} X_i = X_1 X_2 \cdots X_k, \quad \prod_{1 \leq i \leq k}^{\leftarrow} X_i = X_k X_{k-1} \cdots X_1.$$

Consider the vector space $(\mathbb{C}^n)^{\otimes \xi^{n-1}}$ and define

$$\begin{aligned} \mathbb{T}^{[j]}(\mathbf{t}^j) &= \prod_{1 \leq k \leq \xi_j}^{\rightarrow} T^{(\xi^{j-1}+k)}(t_k^j), \\ \mathbb{R}^{[k,j]}(\mathbf{t}^k, \mathbf{t}^j) &= \prod_{1 \leq i \leq \xi_k}^{\rightarrow} \left(\prod_{1 \leq l \leq \xi_j}^{\leftarrow} R^{(\xi^{k-1}+i, \xi^{j-1}+l)}(t_i^k - t_l^j) \right), \end{aligned}$$

where we consider $T^{(\xi^{j-1}+k)}(t_k^j)$ as a matrix with noncommuting entries belonging to $Y(\mathfrak{gl}_n)$.

For the expression

$$\widehat{\mathbb{T}}_{\boldsymbol{\xi}}(\mathbf{t}) = \mathbb{T}^{[1]}(\mathbf{t}^1) \cdots \mathbb{T}^{[n-1]}(\mathbf{t}^{n-1}) \prod_{1 \leq i \leq n-1}^{\leftarrow} \left(\prod_{1 \leq j < i}^{\leftarrow} \mathbb{R}^{[i,j]}(\mathbf{t}^i, \mathbf{t}^j) \right), \quad (3.1)$$

denote by $\mathbb{B}_{\boldsymbol{\xi}}(\mathbf{t})$ the following entry

$$\mathbb{B}_{\boldsymbol{\xi}}(\mathbf{t}) = \left(\widehat{\mathbb{T}}_{\boldsymbol{\xi}}(\mathbf{t}) \right)_{\mathbf{2}^{\xi_1}, \mathbf{3}^{\xi_2}, \dots, \mathbf{n}^{\xi_{n-1}}}^{\mathbf{1}^{\xi_1}, \mathbf{2}^{\xi_2}, \dots, \mathbf{n-1}^{\xi_{n-1}}},$$

where

$$\begin{aligned} \mathbf{1}^{\xi_1}, \mathbf{2}^{\xi_2}, \dots, (\mathbf{n-1})^{\xi_{n-1}} &= \underbrace{1, 1, \dots, 1}_{\xi_1}, \underbrace{2, 2, \dots, 2}_{\xi_2}, \dots, \underbrace{n-1, n-1, \dots, n-1}_{\xi_{n-1}}, \\ \mathbf{2}^{\xi_1}, \mathbf{3}^{\xi_2}, \dots, \mathbf{n}^{\xi_{n-1}} &= \underbrace{2, 2, \dots, 2}_{\xi_1}, \underbrace{3, 3, \dots, 3}_{\xi_2}, \dots, \underbrace{n, n, \dots, n}_{\xi_{n-1}}. \end{aligned}$$

To indicate the dependence on n , if necessary, we will write $\mathbb{B}_{\boldsymbol{\xi}}^{(n)}(\mathbf{t})$.

Example. Let $n = 2$ and $\boldsymbol{\xi} = (\xi_1)$. Then $\mathbb{B}_{\boldsymbol{\xi}}^{(2)}(\mathbf{t}) = T_2^1(t_1^1) \cdots T_2^1(t_{\xi_1}^1)$. Abusing notation, we will farther write $\mathbb{B}_{\xi_1}^{(2)}(\mathbf{t})$ instead of $\mathbb{B}_{\boldsymbol{\xi}}^{(2)}(\mathbf{t})$.

Example. Let $n = 4$ and $\xi = (1, 1, 1)$. Then

$$\begin{aligned} \mathbb{B}_{\xi}^{(4)}(t) &= T_2^1(t_1)T_3^2(t_1^2)T_4^3(t_1^3) \\ &\quad + \frac{1}{t_1^2 - t_1^1}T_3^1(t_1)T_2^2(t_1^2)T_4^3(t_1^3) + \frac{1}{t_1^3 - t_1^2}T_2^1(t_1)T_4^2(t_1^2)T_3^3(t_1^3) \\ &\quad + \frac{1}{(t_1^2 - t_1^1)(t_1^3 - t_1^2)}(T_4^1(t_1)T_2^2(t_1^2)T_3^3(t_1^3) + T_3^1(t_1)T_4^2(t_1^2)T_2^3(t_1^3)) \\ &\quad + \frac{(t_1^2 - t_1^1)(t_1^3 - t_1^2) + 1}{(t_1^2 - t_1^1)(t_1^3 - t_1^1)(t_1^3 - t_1^2)}T_4^1(t_1)T_3^2(t_1^2)T_2^3(t_1^3). \end{aligned}$$

For a weight singular vector v with respect to the action of $Y(\mathfrak{gl}_n)$, we call the expression $\mathbb{B}_{\xi}(t)v$ the (rational) vector-valued weight function of weight $(\xi_1, \xi_2 - \xi_1, \dots, \xi_{n-1} - \xi_{n-2}, -\xi_{n-1})$ associated with v .

From now on, we will consider only the case $n = 4$. We are interested in writing down the following expansion for a weight function in a evaluation module over the $Y(\mathfrak{gl}_4)$:

$$\mathbb{B}_{\xi}(t)v = \sum_{\vec{m} \in \mathbb{Z}_{\geq 0}^6} F_{\vec{m}}(t) \cdot e_{32}^{m_{32}} e_{31}^{m_{31}} e_{42}^{m_{42}} e_{41}^{m_{41}} e_{21}^{m_{21}} e_{43}^{m_{43}} v \quad (3.2)$$

with the functions $F_{\vec{m}}(t)$ given by explicit formulae. Various similar expansions for $\mathbb{B}_{\xi}(t)v$ were obtained in [25], however, expansion (3.2) is not covered there.

4 Splitting property of the weight functions

Let $T_{ab}^{(2)}(u)$ be series (2.3) for the algebra $Y(\mathfrak{gl}_2)$, and $R^{(2)}(u)$ be the corresponding rational R -matrix, see (2.1). Consider two $Y(\mathfrak{gl}_2)$ -module structures on the vector space \mathbb{C}^2 . The first one, called $L(x)$, is given by the rule

$$\pi(x): T^{(2)}(u) \mapsto R^{(2)}(u - x),$$

and the second one, called $\bar{L}(x)$, is given by the rule

$$\varpi(x): T^{(2)}(u) \mapsto ((R^{(2)}(x - u))^{(21)})^{t_2},$$

where the superscript t_2 stands for the matrix transposition in the second tensor factor.

Let $\mathbf{w}_1, \mathbf{w}_2$ be the standard basis of the space \mathbb{C}^2 . The module $L(x)$ is a highest weight evaluation module with \mathfrak{gl}_2 highest weight $(1, 0)$ and highest weight vector \mathbf{w}_1 . The module $\bar{L}(x)$ is a highest weight evaluation module with \mathfrak{gl}_2 highest weight $(0, -1)$ and highest weight vector \mathbf{w}_2 . For any $X \in \text{End}(\mathbb{C}^2)$, set $\nu(X) = X\mathbf{w}_1$ and $\bar{\nu}(X) = X\mathbf{w}_2$.

Recall the coproducts Δ and $\tilde{\Delta}$, see (2.5) and (2.6), and the embeddings $\psi_2: Y(\mathfrak{gl}_2) \rightarrow Y(\mathfrak{gl}_4)$ and $\phi_2: Y(\mathfrak{gl}_2) \rightarrow Y(\mathfrak{gl}_4)$ given by (2.8). For any k , denote by

$$\Delta^{(k)}: Y(\mathfrak{gl}_2) \rightarrow (Y(\mathfrak{gl}_2))^{\otimes(k+1)} \quad \text{and} \quad \widetilde{\Delta}^{(k)}: Y(\mathfrak{gl}_2) \rightarrow (Y(\mathfrak{gl}_2))^{\otimes(k+1)}$$

the iterated coproduct and opposite coproduct. Consider the maps

$$\begin{aligned} \psi_2(x_1, \dots, x_k): Y(\mathfrak{gl}_2) &\rightarrow (\mathbb{C}^2)^{\otimes k} \otimes Y(\mathfrak{gl}_4), \\ \psi_2(x_1, \dots, x_k) &= (\nu^{\otimes k} \otimes \text{id}) \circ (\pi(x_1) \otimes \dots \otimes \pi(x_k) \otimes \psi_2) \circ \Delta^{(k)}, \end{aligned}$$

and

$$\begin{aligned} \phi_2(x_1, \dots, x_k): Y(\mathfrak{gl}_2) &\rightarrow (\mathbb{C}^2)^{\otimes k} \otimes Y(\mathfrak{gl}_4), \\ \phi_2(x_1, \dots, x_k) &= (\bar{\nu}^{\otimes k} \otimes \text{id}) \circ (\varpi(x_1) \otimes \dots \otimes \varpi(x_k) \otimes \phi_2) \circ \tilde{\Delta}^{(k)}. \end{aligned}$$

For any element $g \in (\mathbb{C}^2)^{\otimes k} \otimes Y(\mathfrak{gl}_4)$, we define its components $g^{\mathbf{a}}$, $\mathbf{a} = (a_1, \dots, a_k)$, by the rule

$$g = \sum_{a_1, \dots, a_k=1}^2 \mathbf{w}_{a_1} \otimes \dots \otimes \mathbf{w}_{a_k} \otimes g^{\mathbf{a}}.$$

In the \mathfrak{gl}_4 case, we have $\boldsymbol{\xi} = (\xi_1, \xi_2, \xi_3)$, and formula (3.1) takes the form

$$\mathbb{B}_{\boldsymbol{\xi}}(\mathbf{t}) = \left(\mathbb{T}^{[1]}(\mathbf{t}^1) \mathbb{T}^{[2]}(\mathbf{t}^2) \mathbb{T}^{[3]}(\mathbf{t}^3) \mathbb{R}^{[32]}(\mathbf{t}^3, \mathbf{t}^2) \mathbb{R}^{[31]}(\mathbf{t}^3, \mathbf{t}^1) \mathbb{R}^{[21]}(\mathbf{t}^2, \mathbf{t}^1) \right)_{\mathbf{2}^{\xi_1}, \mathbf{3}^{\xi_2}, \mathbf{4}^{\xi_3}}^{\mathbf{1}^{\xi_1}, \mathbf{2}^{\xi_2}, \mathbf{3}^{\xi_3}}.$$

Proposition 4.1. *Let v be a $Y(\mathfrak{gl}_4)$ -singular vector, ξ_1, ξ_2, ξ_3 be nonnegative integers, and $\mathbf{t} = (t_1^1, \dots, t_{\xi_1}^1; t_1^2, \dots, t_{\xi_2}^2; t_1^3, \dots, t_{\xi_3}^3)$. Then*

$$\mathbb{B}_{\boldsymbol{\xi}}(\mathbf{t})v = \sum_{\mathbf{a}, \mathbf{b}} (\mathcal{T}(\mathbf{t}^2))_{\mathbf{b}}^{\mathbf{a}} (\phi_2(\mathbf{t}^2) (\mathbb{B}_{\xi_1}^{(2)}(\mathbf{t}^1)))^{\mathbf{a}} (\psi_2(\mathbf{t}^2) (\mathbb{B}_{\xi_3}^{(2)}(\mathbf{t}^3)))^{\mathbf{b}-\mathbf{2}} v, \quad (4.1)$$

where the sum is taken over all sequences $\mathbf{a} = (a_1, a_2, \dots, a_{\xi_2})$, $\mathbf{b} = (b_1, b_2, \dots, b_{\xi_2})$, such that $a_i \in \{1, 2\}$, $b_i \in \{3, 4\}$ for all $i = 1, \dots, \xi_2$, $\mathbf{b} - \mathbf{2} = (b_1 - 2, b_2 - 2, \dots, b_{\xi_2} - 2)$, and

$$(\mathcal{T}(\mathbf{t}^2))_{\mathbf{b}}^{\mathbf{a}} = T(t_1^2)_{b_1}^{a_1} T(t_2^2)_{b_2}^{a_2} \dots T(t_{\xi_2}^2)_{b_{\xi_2}}^{a_{\xi_2}}.$$

Proof. Formula (4.1) follows from the definition of the maps $\psi_2(\mathbf{t}^2)$ and $\phi_2(\mathbf{t}^2)$ and Lemma 4.2 below. ■

Lemma 4.2. *One has*

$$\mathbb{B}_{\boldsymbol{\xi}}(\mathbf{t})v = \sum_{\mathbf{a}, \mathbf{b}} (\mathcal{T}(\mathbf{t}^2))_{\mathbf{b}}^{\mathbf{a}} \left(\mathbb{R}^{[21]}(\mathbf{t}^2, \mathbf{t}^1) \mathbb{T}^{[1]}(\mathbf{t}^1) \right)_{\mathbf{2}^{\xi_1}, \mathbf{a}, \mathbf{3}^{\xi_3}}^{\mathbf{1}^{\xi_1}, \mathbf{2}^{\xi_2}, \mathbf{3}^{\xi_3}} \left(\mathbb{T}^{[3]}(\mathbf{t}^3) \mathbb{R}^{[32]}(\mathbf{t}^3, \mathbf{t}^2) \right)_{\mathbf{1}^{\xi_1}, \mathbf{3}^{\xi_2}, \mathbf{4}^{\xi_3}}^{\mathbf{1}^{\xi_1}, \mathbf{b}, \mathbf{3}^{\xi_3}} v,$$

where the sum over \mathbf{a}, \mathbf{b} is the same as in formula (4.1).

Proof. Using Yang–Baxter equation (2.2), we can write $\mathbb{B}(\mathbf{t})$ in the following form:

$$\mathbb{B}_{\boldsymbol{\xi}}(\mathbf{t})v = \left(\mathbb{R}^{[21]}(\mathbf{t}^2, \mathbf{t}^1) \mathbb{T}^{[2]}(\mathbf{t}^2) \mathbb{T}^{[1]}(\mathbf{t}^1) \mathbb{T}^{[3]}(\mathbf{t}^3) \mathbb{R}^{[31]}(\mathbf{t}^3, \mathbf{t}^1) \mathbb{R}^{[32]}(\mathbf{t}^3, \mathbf{t}^2) \right)_{\mathbf{2}^{\xi_1}, \mathbf{3}^{\xi_2}, \mathbf{4}^{\xi_3}}^{\mathbf{1}^{\xi_1}, \mathbf{2}^{\xi_2}, \mathbf{3}^{\xi_3}} v.$$

Therefore,

$$\mathbb{B}_{\boldsymbol{\xi}}(\mathbf{t})v = \sum_{\mathbf{p}, \mathbf{q}, \mathbf{r}, \mathbf{s}} \left(\mathbb{R}^{[21]} \mathbb{T}^{[2]} \mathbb{T}^{[1]} \right)_{\mathbf{p}, \mathbf{q}, \mathbf{3}^{\xi_3}}^{\mathbf{1}^{\xi_1}, \mathbf{2}^{\xi_2}, \mathbf{3}^{\xi_3}} \left(\mathbb{T}^{[3]} \right)_{\mathbf{p}, \mathbf{q}, \mathbf{r}}^{\mathbf{p}, \mathbf{q}, \mathbf{3}^{\xi_3}} \left(\mathbb{R}^{[31]} \right)_{\mathbf{2}^{\xi_1}, \mathbf{q}, \mathbf{s}}^{\mathbf{p}, \mathbf{q}, \mathbf{r}} \left(\mathbb{R}^{[32]} \right)_{\mathbf{2}^{\xi_1}, \mathbf{3}^{\xi_2}, \mathbf{4}^{\xi_3}}^{\mathbf{2}^{\xi_1}, \mathbf{q}, \mathbf{s}} v, \quad (4.2)$$

where $\mathbf{p} = (p_1, \dots, p_{\xi_1})$, $\mathbf{q} = (q_1, \dots, q_{\xi_2})$, $\mathbf{r} = (r_1, \dots, r_{\xi_3})$, $\mathbf{s} = (s_1, \dots, s_{\xi_3})$. In (4.2), we omitted the arguments $\mathbf{t}^1, \mathbf{t}^2, \mathbf{t}^3$ since they can be restored from the context.

We say that $\mathbf{r} \geq \mathbf{3}^{\xi_3}$ if $r_i \geq 3$ for all $i = 1, \dots, \xi_3$. Observe that by the definition of a singular vector and the commutation relations

$$T_b^3(w) T_d^3(u) = \frac{w - u - 1}{w - u} T_d^3(u) T_b^3(w) + \frac{1}{w - u} T_d^3(w) T_b^3(u),$$

we have $\mathbb{T}^{[3]}(\mathbf{t}^3)_{\mathbf{r}}^{\mathbf{3}^{\xi_3}} v = 0$ unless $\mathbf{r} \geq \mathbf{3}^{\xi_3}$.

Furthermore, for $\mathbf{r} \geq \mathbf{3}^{\xi_3}$, we have by induction on ξ_3 that

$$\left(\mathbb{R}^{[31]} \right)_{\mathbf{2}^{\xi_1}, \mathbf{q}, \mathbf{s}}^{\mathbf{p}, \mathbf{q}, \mathbf{r}} = \delta_{\mathbf{p}, \mathbf{2}^{\xi_1}} \delta_{\mathbf{r}, \mathbf{s}}.$$

Indeed, for $\xi_3 = 0$, the statement is true. Assume that

$$\left(\begin{smallmatrix} [31] \\ \mathbb{R} \end{smallmatrix}\right)_{\mathbf{2}^{\xi_1}, \mathbf{q}, \mathbf{s}}^{\mathbf{p}, \mathbf{q}, \mathbf{r}} = \delta_{\mathbf{p}, \mathbf{2}^{\xi_1}} \delta_{\mathbf{r}, \mathbf{s}}$$

for $\mathbf{r} \geq \mathbf{3}^{\xi_3}$ if $\xi_3 = n - 1$, and consider the case $\xi_3 = n$. Let $\mathbf{r} = (r_1, \dots, r_n)$, $\mathbf{s} = (s_1, \dots, s_n)$, $\tilde{\mathbf{r}} = (r_1, \dots, r_{n-1})$, $\tilde{\mathbf{s}} = (s_1, \dots, s_{n-1})$, then we have

$$\left(\begin{smallmatrix} [31] \\ \mathbb{R} \end{smallmatrix}\right)_{\mathbf{2}^{\xi_1}, \mathbf{q}, \mathbf{s}}^{\mathbf{p}, \mathbf{q}, \mathbf{r}} = \sum_{\mathbf{x}} \left(\prod_{1 \leq i \leq n-1}^{\rightarrow} \left(\prod_{1 \leq j \leq \xi_1}^{\leftarrow} R^{(\xi^2+i, j)} \right) \right)_{\mathbf{x}, \mathbf{q}, \tilde{\mathbf{s}}, r_n}^{\mathbf{p}, \mathbf{q}, \tilde{\mathbf{r}}, r_n} \left(\prod_{1 \leq k \leq \xi_1}^{\leftarrow} R^{(\xi^3, k)} \right)_{\mathbf{2}^{\xi_1}, \mathbf{q}, \tilde{\mathbf{s}}, s_n}^{\mathbf{x}, \mathbf{q}, \tilde{\mathbf{s}}, r_n}. \quad (4.3)$$

Observe that the R -matrix entry R_{ik}^{jl} with $i \neq l$ is not zero if and only if $i = j$ and $k = l$, and $R_{ik}^{ik} = 1$. Because of that and since $r_n \geq 3$, the last factor $\left(\prod R^{(\xi^3, k)} \right)_{\mathbf{2}^{\xi_1}, \mathbf{q}, \tilde{\mathbf{s}}, s_n}^{\mathbf{x}, \mathbf{q}, \tilde{\mathbf{s}}, r_n}$ in (4.3) equals $\delta_{\mathbf{x}, \mathbf{2}^{\xi_1}} \delta_{r_n, s_n}$, and we get

$$\left(\begin{smallmatrix} [31] \\ \mathbb{R} \end{smallmatrix}\right)_{\mathbf{2}^{\xi_1}, \mathbf{q}, \mathbf{s}}^{\mathbf{p}, \mathbf{q}, \mathbf{r}} = \left(\prod_{1 \leq i \leq n-1}^{\rightarrow} \left(\prod_{1 \leq j \leq \xi_1}^{\leftarrow} R^{(\xi^2+i, j)} \right) \right)_{\mathbf{2}^{\xi_1}, \mathbf{q}, \tilde{\mathbf{s}}, r_n}^{\mathbf{p}, \mathbf{q}, \tilde{\mathbf{r}}, r_n} \delta_{r_n, s_n} = \delta_{\mathbf{p}, \mathbf{2}^{\xi_1}} \delta_{\tilde{\mathbf{r}}, \tilde{\mathbf{s}}} \delta_{r_n, s_n} = \delta_{\mathbf{p}, \mathbf{2}^{\xi_1}} \delta_{\mathbf{r}, \mathbf{s}},$$

by the induction assumption.

Since

$$\left(\begin{smallmatrix} [3] \\ \mathbb{T} \end{smallmatrix}\right)_{\mathbf{p}, \mathbf{q}, \mathbf{r}}^{\mathbf{p}, \mathbf{q}, \mathbf{3}^{\xi_3}} v = 0$$

unless $\mathbf{r} \geq \mathbf{3}^{\xi_3}$ and

$$\left(\begin{smallmatrix} [31] \\ \mathbb{R} \end{smallmatrix}\right)_{\mathbf{2}^{\xi_1}, \mathbf{q}, \mathbf{s}}^{\mathbf{p}, \mathbf{q}, \mathbf{r}} = \delta_{\mathbf{p}, \mathbf{2}^{\xi_1}} \delta_{\mathbf{r}, \mathbf{s}}$$

for $\mathbf{r} \geq 3$, formula (4.2) becomes

$$\mathbb{B}_{\xi}(\mathbf{t})v = \sum_{\mathbf{p}, \mathbf{q}, \mathbf{r}} \left(\begin{smallmatrix} [21] [2] [1] \\ \mathbb{R} \mathbb{T} \mathbb{T} \end{smallmatrix}\right)_{\mathbf{2}^{\xi_1}, \mathbf{q}, \mathbf{3}^{\xi_3}}^{\mathbf{1}^{\xi_1}, \mathbf{2}^{\xi_2}, \mathbf{3}^{\xi_3}} \left(\begin{smallmatrix} [3] \\ \mathbb{T} \end{smallmatrix}\right)_{\mathbf{2}^{\xi_1}, \mathbf{q}, \mathbf{r}}^{\mathbf{2}^{\xi_1}, \mathbf{q}, \mathbf{3}^{\xi_3}} \left(\begin{smallmatrix} [32] \\ \mathbb{R} \end{smallmatrix}\right)_{\mathbf{2}^{\xi_1}, \mathbf{3}^{\xi_2}, \mathbf{4}^{\xi_3}}^{\mathbf{2}^{\xi_1}, \mathbf{q}, \mathbf{r}} v,$$

and can be further transformed as

$$\mathbb{B}_{\xi}(\mathbf{t})v = \sum_{\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{r}} \left(\begin{smallmatrix} [2] \\ \mathbb{T} \end{smallmatrix}\right)_{\mathbf{c}, \mathbf{b}, \mathbf{3}^{\xi_3}}^{\mathbf{c}, \mathbf{a}, \mathbf{3}^{\xi_3}} \left(\begin{smallmatrix} [21] \\ \mathbb{R} \end{smallmatrix}\right)_{\mathbf{c}, \mathbf{a}, \mathbf{3}^{\xi_3}}^{\mathbf{1}^{\xi_1}, \mathbf{2}^{\xi_2}, \mathbf{3}^{\xi_3}} \left(\begin{smallmatrix} [1] \\ \mathbb{T} \end{smallmatrix}\right)_{\mathbf{2}^{\xi_1}, \mathbf{b}, \mathbf{3}^{\xi_1}}^{\mathbf{c}, \mathbf{b}, \mathbf{3}^{\xi_3}} \left(\begin{smallmatrix} [3] \\ \mathbb{T} \end{smallmatrix}\right)_{\mathbf{2}^{\xi_1}, \mathbf{b}, \mathbf{r}}^{\mathbf{2}^{\xi_1}, \mathbf{b}, \mathbf{3}^{\xi_3}} \left(\begin{smallmatrix} [32] \\ \mathbb{R} \end{smallmatrix}\right)_{\mathbf{2}^{\xi_1}, \mathbf{3}^{\xi_2}, \mathbf{4}^{\xi_3}}^{\mathbf{2}^{\xi_1}, \mathbf{b}, \mathbf{r}} v, \quad (4.4)$$

where the sum is over all sequences $\mathbf{a} = (a_1, \dots, a_{\xi_2})$, $\mathbf{b} = (b_1, \dots, b_{\xi_2})$, $\mathbf{c} = (c_1, \dots, c_{\xi_1})$, $\mathbf{r} = (r_1, \dots, r_{\xi_3})$ such that $a_i, b_i, c_i, r_i \in \{1, 2, 3, 4\}$. Since

$$\left(\begin{smallmatrix} [21] \\ \mathbb{R} \end{smallmatrix}\right)_{\mathbf{c}, \mathbf{a}, \mathbf{3}^{\xi_3}}^{\mathbf{1}^{\xi_1}, \mathbf{2}^{\xi_2}, \mathbf{3}^{\xi_3}} = 0$$

if $a_i \geq 3$ for some i , and

$$\left(\begin{smallmatrix} [32] \\ \mathbb{R} \end{smallmatrix}\right)_{\mathbf{2}^{\xi_1}, \mathbf{3}^{\xi_2}, \mathbf{4}^{\xi_3}}^{\mathbf{2}^{\xi_1}, \mathbf{b}, \mathbf{r}} = 0$$

if $b_i \leq 2$ for some i , terms in the sum in the right-hand side of (4.4) equal zero unless $a_i \in \{1, 2\}$ and $b_i \in \{3, 4\}$ for all i . Taking the sum over \mathbf{c} and \mathbf{r} in formula (4.4), we get the statement of Lemma 4.2. ■

Example. Here we illustrate the proof of the relation

$$\left(\begin{smallmatrix} [31] \\ \mathbb{R} \end{smallmatrix}\right)_{\mathbf{2}^{\xi_1}, \mathbf{q}, \mathbf{s}}^{\mathbf{p}, \mathbf{q}, \mathbf{r}} = \delta_{\mathbf{p}, \mathbf{2}^{\xi_1}} \delta_{\mathbf{r}, \mathbf{s}}$$

if $\mathbf{r} \geq \mathbf{3}^{\xi_3}$ for $\xi_1 = \xi_3 = 2$. In this case, $\mathbf{p} = (p_1, p_2)$, $\mathbf{r} = (r_1, r_2)$, $\mathbf{s} = (s_1, s_2)$, and

$$\left(\begin{smallmatrix} [31] \\ \mathbb{R} \end{smallmatrix}\right)_{\mathbf{2}^{\xi_1}, \mathbf{q}, \mathbf{s}}^{\mathbf{p}, \mathbf{q}, \mathbf{r}} = \sum_{a, b, c, d} R_{ab}^{p_2 r_1} (t_1^3 - t_2^1) R_{cs_1}^{p_1 b} (t_1^3 - t_1^1) R_{2d}^{ar_2} (t_2^3 - t_2^1) R_{2s_2}^{cd} (t_2^3 - t_1^1).$$

For $r_1 \geq 3$, $r_2 \geq 3$, we have $R_{2d}^{ar_2} (t_2^3 - t_2^1) = \delta_{a,2} \delta_{r_2, d}$, thus

$$\left(\begin{smallmatrix} [31] \\ \mathbb{R} \end{smallmatrix}\right)_{\mathbf{2}^{\xi_1}, \mathbf{q}, \mathbf{s}}^{\mathbf{p}, \mathbf{q}, \mathbf{r}} = \sum_{b, c} R_{2b}^{p_2 r_1} (t_1^3 - t_2^1) R_{cs_1}^{p_1 b} (t_1^3 - t_1^1) R_{2s_2}^{cr_2} (t_2^3 - t_1^1).$$

Then $R_{2s_2}^{cr_2} (t_2^3 - t_1^1) = \delta_{c,2} \delta_{r_2, s_2}$ and $R_{2b}^{p_2 r_1} (t_1^3 - t_2^1) = \delta_{p_2, 2} \delta_{r_1, b}$, so that

$$\left(\begin{smallmatrix} [31] \\ \mathbb{R} \end{smallmatrix}\right)_{\mathbf{2}^{\xi_1}, \mathbf{q}, \mathbf{s}}^{\mathbf{p}, \mathbf{q}, \mathbf{r}} = R_{2s_1}^{p_1 r_1} (t_1^3 - t_1^1) \delta_{p_2, 2} \delta_{r_2, s_2} = \delta_{p_2, 2} \delta_{p_1, 2} \delta_{r_1, s_1} \delta_{r_2, s_2} = \delta_{\mathbf{p}, \mathbf{2}^{\xi_1}} \delta_{\mathbf{r}, \mathbf{s}}.$$

5 Main theorem for the \mathfrak{gl}_4 case

The main result of this paper is Theorem 5.7 formulated at the end of this section. We will approach it in several steps.

For a nonnegative integer m , set

$$Q_m(t_1, \dots, t_m) = \prod_{1 \leq i < j \leq m} \frac{t_i - t_j - 1}{t_i - t_j}. \quad (5.1)$$

For an expression $f(t_1, \dots, t_m)$, define

$$\text{Sym}_{\mathbf{t}} f(t_1, \dots, t_m) = \sum_{\sigma \in S_m} f(t_{\sigma(1)}, \dots, t_{\sigma(m)}),$$

and

$$\overline{\text{Sym}}_{\mathbf{t}} f(t_1, \dots, t_m) = \overline{\text{Sym}}_{\mathbf{t}} (f(t_1, \dots, t_m) Q_m(t_1, \dots, t_m)). \quad (5.2)$$

To simplify notation, we will write $T_{ij}^{(2)}$ instead of $(T^{(2)})_j^i$.

Proposition 5.1. *Let ξ be a nonnegative integer and $\mathbf{t} = (t_1, \dots, t_\xi)$. Then*

$$\begin{aligned} \Delta(\mathbb{B}_\xi^{(2)}(\mathbf{t})) &= \sum_{\eta=0}^{\xi} \frac{1}{(\xi - \eta)! \eta!} \\ &\quad \times \overline{\text{Sym}}_{\mathbf{t}} \left((\mathbb{B}_\eta^{(2)}(t_1, \dots, t_\eta) \otimes \mathbb{B}_{\xi-\eta}^{(2)}(t_{\eta+1}, \dots, t_\xi)) \left(\prod_{i=\eta+1}^{\xi} T_{22}^{(2)}(t_i) \otimes \prod_{j=1}^{\eta} T_{11}^{(2)}(t_j) \right) \right). \end{aligned}$$

This proposition goes back to [4, 28]. For convenience, we give its proof in Appendix A.

Given a subset I of $\{1, 2, \dots, k\}$ denote by I^* the complement of I in $\{1, 2, \dots, k\}$. Define a vector $\mathbf{w}^I \in (\mathbb{C}^2)^{\otimes k}$ by the rule

$$\mathbf{w}^I = \mathbf{w}_{a_1} \otimes \mathbf{w}_{a_2} \otimes \dots \otimes \mathbf{w}_{a_k},$$

where $a_i = 2$ if $i \in I$, and $a_i = 1$ if $i \notin I$.

Fix a $Y(\mathfrak{gl}_2)$ -module V and a weight singular vector $v \in V$ with respect to the $Y(\mathfrak{gl}_2)$ -action,

$$T_{21}^{(2)}(u)v = 0, \quad T_{11}^{(2)}(u)v = \langle T_{11}^{(2)}(u)v \rangle v, \quad T_{22}^{(2)}(u)v = \langle T_{22}^{(2)}(u)v \rangle v.$$

Here $\langle T_{11}^{(2)}(u)v \rangle$ and $\langle T_{22}^{(2)}(u)v \rangle$ are the corresponding eigenvalues. Given complex numbers z_1, \dots, z_k , consider the $Y(\mathfrak{gl}_2)$ -module $L(z_1) \otimes \dots \otimes L(z_k) \otimes V$. Observe that $\mathbf{w}_1^{\otimes k} \otimes v$ is a weight singular vector with respect to the action of $Y(\mathfrak{gl}_2)$ in this module.

Proposition 5.2. *For the $Y(\mathfrak{gl}_2)$ -module $L(z_1) \otimes \dots \otimes L(z_k) \otimes V$, we have*

$$\begin{aligned} & \mathbb{B}_\xi^{(2)}(\mathbf{t})(\mathbf{w}_1^{\otimes k} \otimes v) \\ &= \sum_I \frac{1}{(\xi - |I|)!} \overline{\text{Sym}}_{\mathbf{t}} \left[F_I(\mathbf{t}, \mathbf{z}) \prod_{a=1}^{|I|} \langle T_{11}^{(2)}(t_a)v \rangle (\mathbf{w}^I \otimes \mathbb{B}_{\xi-|I|}^{(2)}(t_{|I|+1}, \dots, t_\xi)v) \right], \end{aligned} \quad (5.3)$$

where the sum is over all subsets $I \subset \{1, \dots, k\}$ such that $|I| \leq \xi$, and for a given $I = \{i_1 < i_2 < \dots < i_{|I|}\}$,

$$F_I(\mathbf{t}, \mathbf{z}) = \prod_{a=1}^{|I|} \left(\frac{1}{t_a - z_{i_a}} \prod_{m=i_a+1}^k \frac{t_a - z_m + 1}{t_a - z_m} \right). \quad (5.4)$$

Proof. Observe that for each $Y(\mathfrak{gl}_2)$ -module $L(z_i)$, $i = 1, \dots, k$, the corresponding vector $\mathbf{w}_1 \in L(z_i)$ is a weight singular vector,

$$T_{11}^{(2)}(u)\mathbf{w}_1 = (1 + (u - z_i)^{-1})\mathbf{w}_1, \quad T_{22}^{(2)}(u)\mathbf{w}_1 = \mathbf{w}_1, \quad T_{21}^{(2)}\mathbf{w}_1 = 0.$$

Moreover, $\mathbb{B}_1^{(2)}(u)\mathbf{w}_1 = T_{12}^{(2)}(u)\mathbf{w}_1 = (u - z_i)^{-1}\mathbf{w}_2$ and $\mathbb{B}_\zeta^{(2)}(u_1, \dots, u_\zeta)\mathbf{w}_1 = 0$ for $\zeta \geq 2$. Then formula (5.3) follows from Proposition 5.1 and identity (A.10) by induction on k . ■

Given complex numbers z_1, \dots, z_k , consider the $Y(\mathfrak{gl}_2)$ -module $V \otimes \bar{L}(z_k) \otimes \dots \otimes \bar{L}(z_1)$. Observe that $v \otimes \mathbf{w}_2^{\otimes k}$ is a weight singular vector with respect to the action of $Y(\mathfrak{gl}_2)$ in this module.

Proposition 5.3. *For the $Y(\mathfrak{gl}_2)$ module $V \otimes \bar{L}(z_k) \otimes \dots \otimes \bar{L}(z_1)$, we have*

$$\begin{aligned} & \mathbb{B}_\xi^{(2)}(\mathbf{t})(v \otimes \mathbf{w}_2^{\otimes k}) \\ &= \sum_I \frac{1}{(\xi - |I|)!} \overline{\text{Sym}}_{\mathbf{t}} \left[\tilde{F}_I(\mathbf{t}, \mathbf{z}) \prod_{i=1}^{|I|} \langle T_{22}^{(2)}(t_{\xi-|I|+i})v \rangle (\mathbb{B}_{\xi-|I|}^{(2)}(t_1, \dots, t_{\xi-|I|})v \otimes \mathbf{w}^{I*}) \right], \end{aligned} \quad (5.5)$$

where the sum is over all subsets $I \subset \{1, \dots, k\}$ such that $|I| \leq \xi$, and for a given $I = \{i_1 < i_2 < \dots < i_{|I|}\}$,

$$\tilde{F}_I(\mathbf{t}, \mathbf{z}) = \prod_{a=1}^{|I|} \left(\frac{1}{z_{i_a} - t_{\xi-a+1}} \prod_{m=i_a+1}^k \frac{z_m - t_{\xi-a+1} + 1}{z_m - t_{\xi-a+1}} \right). \quad (5.6)$$

Proof. Observe that for each $Y(\mathfrak{gl}_2)$ -module $\bar{L}(z_i)$, $i = 1, \dots, k$, the corresponding vector $\mathbf{w}_2 \in \bar{L}(z_i)$ is a weight singular vector,

$$T_{11}^{(2)}(u)\mathbf{w}_2 = \mathbf{w}_2, \quad T_{22}^{(2)}(u)\mathbf{w}_2 = (1 + (z_i - u)^{-1})\mathbf{w}_2, \quad T_{21}^{(2)}(u)\mathbf{w}_2 = 0.$$

Moreover, $\mathbb{B}_1^{(2)}(u)\mathbf{w}_2 = T_{12}^{(2)}(u)\mathbf{w}_2 = (z_i - u)^{-1}\mathbf{w}_1$, and $\mathbb{B}_\zeta^{(2)}(u_1, \dots, u_\zeta)\mathbf{w}_2 = 0$ for $\zeta \geq 2$. Then formula (5.5) follows from Proposition 5.1 and identity (A.10) by induction on k . ■

For $\mathbf{t} = (t_1, \dots, t_\xi)$, $\mathbf{z} = (z_1, \dots, z_k)$, $y \in \mathbb{C}$, and a subset $I = \{i_1 < i_2 < \dots < i_{|I|}\} \subset \{1, \dots, k\}$, define the functions

$$V_I(\mathbf{t}, \mathbf{z}, y) = \frac{1}{(\xi - |I|)!} \overline{\text{Sym}}_{\mathbf{t}} \left(F_I(\mathbf{t}, \mathbf{z}) \prod_{a=1}^{|I|} (t_a - y) \right) \quad (5.7)$$

and

$$\tilde{V}_I(\mathbf{t}, \mathbf{z}, y) = \frac{1}{(\xi - |I|)!} \overline{\text{Sym}}_{\mathbf{t}} \left(\tilde{F}_I(\mathbf{t}, \mathbf{z}) \prod_{a=1}^{|I|} (t_{\xi-a+1} - y) \right). \quad (5.8)$$

Consider the collection $\mathcal{S}_{p,q,r,k}$ of pairs of subsets of $\{1, \dots, k\}$ with given cardinalities of the subsets and their intersection. Namely,

$$\mathcal{S}_{p,q,r,k} = \{(I, J) \mid I, J \subset \{1, \dots, k\}, |I| = p, |J| = q, |I \cap J| = r\}.$$

For $I \subset \{1, \dots, k\}$, set $\tilde{I} = \{k - i + 1, i \in I\}$.

Theorem 5.4. *Let V be a \mathfrak{gl}_4 -module and $v \in V$ a \mathfrak{gl}_4 -singular vector of weight $(\Lambda^1, \Lambda^2, \Lambda^3, \Lambda^4)$. Let ξ_1, ξ_2, ξ_3 be nonnegative integers, $\mathbf{t}^1 = (t_1^1, \dots, t_{\xi_1}^1)$, $\mathbf{t}^2 = (t_1^2, \dots, t_{\xi_2}^2)$, $\mathbf{t}^3 = (t_1^3, \dots, t_{\xi_3}^3)$, and $\mathbf{t} = (\mathbf{t}^1, \mathbf{t}^2, \mathbf{t}^3)$. For every triple (p, q, r) , $p = 0, \dots, \min(\xi_2, \xi_3)$, $q = 0, \dots, \min(\xi_2, \xi_1)$, $r = \max(0, p + q - \xi_2), \dots, \min(p, q)$, fix a pair $(I_{p,q,r}, J_{p,q,r}) \in \mathcal{S}_{p,q,r,\xi_2}$. Then,*

(a) *In the evaluation $Y(\mathfrak{gl}_4)$ -module $V(x)$, one has*

$$\begin{aligned} \mathbb{B}_{\xi}(\mathbf{t})v &= \prod_{a=1}^3 \prod_{i=1}^{\xi_a} \frac{1}{t_i^a - x} \\ &\times \sum_{p=0}^{\min(\xi_2, \xi_3)} \sum_{q=0}^{\min(\xi_2, \xi_1)} \sum_{r=\max(0, p+q-\xi_2)}^{\min(p, q)} \left(\overline{\text{Sym}}_{\mathbf{t}^2} (V_{\tilde{I}_{p,q,r}}(\mathbf{t}^3, \mathbf{t}^2, x - \Lambda^3) \tilde{V}_{J_{p,q,r}}(\mathbf{t}^1, \check{\mathbf{t}}^2, x - \Lambda^2)) \right. \\ &\quad \left. \times \frac{e_{32}^{\xi_2-p-q+r} e_{31}^{q-r} e_{42}^{p-r} e_{41}^r e_{21}^{\xi_1-q} e_{43}^{\xi_3-p} v}{(p-r)!(q-r)!r!(\xi_2-p-q-r)!} \right), \end{aligned} \quad (5.9)$$

where $\check{\mathbf{t}}^2 = (t_{\xi_2}^2, \dots, t_1^2)$.

(b) *The function $\overline{\text{Sym}}_{\mathbf{t}^2} (V_{\tilde{I}_{p,q,r}}(\mathbf{t}^3, \mathbf{t}^2, x - \Lambda^3) \tilde{V}_{J_{p,q,r}}(\mathbf{t}^1, \check{\mathbf{t}}^2, x - \Lambda^2))$ in (5.9) does not depend on the choice of the pair $(I_{p,q,r}, J_{p,q,r})$.*

Proof. Item (a) follows from Propositions 5.5 and 5.6 given below. Item (b) is an immediate corollary of Proposition 5.6.

Propositions 5.5 and 5.6 are proved in Sections 6 and 7, respectively. ■

Proposition 5.5. *In the notation of Theorem 5.4, we have*

$$\begin{aligned} \mathbb{B}_{\xi}(\mathbf{t})v &= \prod_{a=1}^3 \prod_{i=1}^{\xi_a} \frac{1}{t_i^a - x} \\ &\times \sum_{p=0}^{\min(\xi_2, \xi_3)} \sum_{q=0}^{\min(\xi_2, \xi_1)} \sum_{r=\max(0, p+q-\xi_2)}^{\min(p, q)} \left(\sum_{(I, J) \in \mathcal{S}_{p,q,r,\xi_2}} \tilde{V}_J(\mathbf{t}^1, \mathbf{t}^2, x - \Lambda^2) V_I(\mathbf{t}^3, \mathbf{t}^2, x - \Lambda^3) \right. \\ &\quad \left. \times e_{32}^{\xi_2-p-q+r} e_{31}^{q-r} e_{42}^{p-r} e_{41}^r e_{21}^{\xi_1-q} e_{43}^{\xi_3-p} v \right). \end{aligned} \quad (5.10)$$

Proposition 5.6. *In the notation of Theorem 5.4, we have*

$$\begin{aligned} & \sum_{(I,J) \in \mathcal{S}_{p,q,r,\xi_2}} \tilde{V}_J(\mathbf{t}^1, \mathbf{t}^2, x - \Lambda^2) V_I(\mathbf{t}^3, \mathbf{t}^2, x - \Lambda^3) \\ &= \frac{\overline{\text{Sym}}_{\mathbf{t}^2}(V_{\tilde{I}_0}(\mathbf{t}^3, \mathbf{t}^2, x - \Lambda^3) \tilde{V}_{J_0}(\mathbf{t}^1, \mathbf{t}^2, x - \Lambda^2))}{(p-r)!(q-r)!r!(\xi_2 - p - q - r)!}, \end{aligned} \quad (5.11)$$

where (I_0, J_0) is any pair from $\mathcal{S}_{p,q,r,\xi_2}$.

Below we reformulate Theorem 5.4 in a more closed form.

Theorem 5.7. *Let V be a \mathfrak{gl}_4 -module and $v \in V$ a \mathfrak{gl}_4 -singular vector of weight $(\Lambda^1, \Lambda^2, \Lambda^3, \Lambda^4)$. Let ξ_1, ξ_2, ξ_3 be nonnegative integers, $\mathbf{t}^1 = (t_1^1, \dots, t_{\xi_1}^1)$, $\mathbf{t}^2 = (t_1^2, \dots, t_{\xi_2}^2)$, $\mathbf{t}^3 = (t_1^3, \dots, t_{\xi_3}^3)$, and $\mathbf{t} = (\mathbf{t}^1, \mathbf{t}^2, \mathbf{t}^3)$. For every triple (p, q, r) , $p = 0, \dots, \min(\xi_2, \xi_3)$, $q = 0, \dots, \min(\xi_2, \xi_1)$, $r = \max(0, p + q - \xi_2), \dots, \min(p, q)$, fix two sequences $\mathbf{i} = \{i_1 < \dots < i_p\}$ and $\mathbf{j} = \{j_1 < \dots < j_q\}$, such that $|\{i_1, \dots, i_p\} \cap \{j_1, \dots, j_q\}| = r$. Then,*

(a) *In the evaluation $Y(\mathfrak{gl}_4)$ -module $V(x)$, one has*

$$\begin{aligned} \mathbb{B}_{\xi}(\mathbf{t})v &= \prod_{a=1}^3 \prod_{i=1}^{\xi_a} \frac{1}{t_i^a - x} \\ &\times \sum_{p=0}^{\min(\xi_2, \xi_3)} \sum_{q=0}^{\min(\xi_2, \xi_1)} \sum_{r=\max(0, p+q-\xi_2)}^{\min(p, q)} \overline{\text{Sym}}_{\mathbf{t}^1} \overline{\text{Sym}}_{\mathbf{t}^2} \overline{\text{Sym}}_{\mathbf{t}^3} G_{\mathbf{i}, \mathbf{j}}(\mathbf{t}) \\ &\times \frac{e_{32}^{\xi_2 - p - q + r} e_{31}^{q-r} e_{42}^{p-r} e_{41}^r e_{21}^{\xi_1 - q} e_{43}^{\xi_3 - p} v}{(\xi_2 - p - q + r)!(q-r)!(p-r)!r!(\xi_1 - q)!(\xi_3 - p)!}, \end{aligned} \quad (5.12)$$

where

$$\begin{aligned} G_{\mathbf{i}, \mathbf{j}}(\mathbf{t}) &= \prod_{a=1}^p \left(\frac{t_a^3 - x + \Lambda^3}{t_a^3 - t_{i_a}^2} \prod_{m=i_a+1}^{\xi_2} \frac{t_a^3 - t_m^2 + 1}{t_a^3 - t_m^2} \right) \\ &\times \prod_{s=1}^q \left(\frac{t_{\xi_1 - q + s}^1 - x + \Lambda^2}{t_{j_s}^2 - t_{\xi_1 - q + s}^1} \prod_{l=1}^{\xi_2 - j_s} \frac{t_l^2 - t_{\xi_1 - q + s}^1 + 1}{t_l^2 - t_{\xi_1 - q + s}^1} \right). \end{aligned} \quad (5.13)$$

(b) *The function $\overline{\text{Sym}}_{\mathbf{t}^1} \overline{\text{Sym}}_{\mathbf{t}^2} \overline{\text{Sym}}_{\mathbf{t}^3} G_{\mathbf{i}, \mathbf{j}}(\mathbf{t})$ does not depend on the choice of the sequences \mathbf{i}, \mathbf{j} .*

Proof. Given the pair $(I_{p,q,r}, J_{p,q,r})$ from the formulation of Theorem 5.4, define the sequences $\mathbf{i} = \{i_1 < \dots < i_p\}$ and $\mathbf{j} = \{j_1 < \dots < j_q\}$ by the rule

$$\begin{aligned} I_{p,q,r} &= \{\xi_2 - i_1 + 1, \xi_2 - i_2 + 1, \dots, \xi_2 - i_p + 1\}, \\ J_{p,q,r} &= \{\xi_2 - j_1 + 1, \xi_2 - j_2 + 1, \dots, \xi_2 - j_q + 1\}. \end{aligned}$$

Notice that $\{i_1, \dots, i_p\} = \tilde{I}_{p,q,r}$, $\{j_1, \dots, j_q\} = \tilde{J}_{p,q,r}$, and

$$|\{i_1, \dots, i_p\} \cap \{j_1, \dots, j_q\}| = |I_{p,q,r} \cap J_{p,q,r}| = r.$$

Then combining formulae (5.4) and (5.6)–(5.8), we obtain that

$$V_{\tilde{I}_{p,q,r}}(\mathbf{t}^3, \mathbf{t}^2, x - \Lambda^3) = \frac{1}{(\xi_3 - p)!} \overline{\text{Sym}}_{\mathbf{t}^3} \prod_{a=1}^p \left(\frac{t_a^3 - x + \Lambda^3}{t_a^3 - t_{i_a}^2} \prod_{m=i_a+1}^{\xi_2} \frac{t_a^3 - t_m^2 + 1}{t_a^3 - t_m^2} \right) \quad (5.14)$$

and

$$\tilde{V}_{J_{p,q,r}}(\mathbf{t}^1, \mathbf{t}^2, x - \Lambda^2) = \frac{1}{(\xi_1 - q)!} \overline{\text{Sym}}_{\mathbf{t}^1} \prod_{b=1}^q \left(\frac{t_{\xi_1-b+1}^1 - x + \Lambda^2}{t_{j_{q-b+1}}^2 - t_{\xi_1-b+1}^1} \prod_{l=1}^{\xi_2-j_{q-b+1}} \frac{t_l^2 - t_{\xi_1-b+1}^1 + 1}{t_l^2 - t_{\xi_1-b+1}^1} \right).$$

After substituting $b = q + 1 - s$, the last formula becomes

$$\begin{aligned} & \tilde{V}_{J_{p,q,r}}(\mathbf{t}^1, \mathbf{t}^2, x - \Lambda^2) \\ &= \frac{1}{(\xi_1 - q)!} \overline{\text{Sym}}_{\mathbf{t}^1} \prod_{s=1}^q \left(\frac{t_{\xi_1-q+s}^1 - x + \Lambda^2}{t_{j_s}^2 - t_{\xi_1-q+s}^1} \prod_{l=1}^{\xi_2-j_s} \frac{t_l^2 - t_{\xi_1-q+s}^1 + 1}{t_l^2 - t_{\xi_1-q+s}^1} \right). \end{aligned} \quad (5.15)$$

Plugging (5.14) and (5.15) into formula (5.9), we obtain formulae (5.12) and (5.13).

Item (b) of Theorem 5.7 is a reformulation of item (b) of Theorem 5.4. ■

Example. Below we give two examples of natural choices of the sequences \mathbf{i}, \mathbf{j} in Theorem 5.7 and write down the corresponding expressions for the function $G_{\mathbf{i}, \mathbf{j}}(\mathbf{t})$, see formula (5.13).

(a) $\mathbf{i} = \mathbf{i}_1 = \{1 < \dots < p\}$, $\mathbf{j} = \mathbf{j}_1 = \{p + 1 - r < \dots < p + q - r\}$. Then

$$\begin{aligned} G_{\mathbf{i}_1, \mathbf{j}_1}(\mathbf{t}) &= \prod_{a=1}^p \left(\frac{t_a^3 - \Lambda^3}{t_a^3 - t_a^2} \prod_{m=a+1}^{\xi_2} \frac{t_a^3 - t_m^2 + 1}{t_a^3 - t_m^2} \right) \\ &\times \prod_{c=1}^q \left(\frac{t_{\xi_1-q+c}^1 - \Lambda^2}{t_{p-r+c}^2 - t_{\xi_1-q+c}^1} \prod_{l=1}^{\xi_2-p+r-c} \frac{t_l^2 - t_{\xi_1-q+c}^1 + 1}{t_l^2 - t_{\xi_1-q+c}^1} \right). \end{aligned}$$

(b) $\mathbf{i} = \mathbf{i}_2 = \{q + 1 - r < \dots < q + p - r\}$, $\mathbf{j} = \mathbf{j}_2 = \{1 < \dots < q\}$. Then

$$\begin{aligned} G_{\mathbf{i}_2, \mathbf{j}_2}(\mathbf{t}) &= \prod_{a=1}^p \left(\frac{t_a^3 - \Lambda^3}{t_a^3 - t_{q-r+a}^2} \prod_{m=q-r+a+1}^{\xi_2} \frac{t_a^3 - t_m^2 + 1}{t_a^3 - t_m^2} \right) \\ &\times \prod_{b=1}^q \left(\frac{t_{\xi_1-b+1}^1 - \Lambda^2}{t_b^2 - t_{\xi_1-b+1}^1} \prod_{l=1}^{\xi_2-b} \frac{t_l^2 - t_{\xi_1-b+1}^1 + 1}{t_l^2 - t_{\xi_1-b+1}^1} \right). \end{aligned}$$

Notice that the equality

$$\overline{\text{Sym}}_{\mathbf{t}^1} \overline{\text{Sym}}_{\mathbf{t}^2} \overline{\text{Sym}}_{\mathbf{t}^3} G_{\mathbf{i}_1, \mathbf{j}_1}(\mathbf{t}) = \overline{\text{Sym}}_{\mathbf{t}^1} \overline{\text{Sym}}_{\mathbf{t}^2} \overline{\text{Sym}}_{\mathbf{t}^3} G_{\mathbf{i}_2, \mathbf{j}_2}(\mathbf{t}),$$

stated in item (b) of Theorem 5.7, is not obvious.

6 Proof of Proposition 5.5

Let V be a \mathfrak{gl}_4 -module and $v \in V$ a \mathfrak{gl}_4 -singular vector of weight $(\Lambda^1, \Lambda^2, \Lambda^3, \Lambda^4)$. Let ξ_1, ξ_2, ξ_3 be nonnegative integers, $\mathbf{t}^1 = (t_1^1, \dots, t_{\xi_1}^1)$, $\mathbf{t}^2 = (t_1^2, \dots, t_{\xi_2}^2)$, $\mathbf{t}^3 = (t_1^3, \dots, t_{\xi_3}^3)$ and $\mathbf{t} = (\mathbf{t}^1, \mathbf{t}^2, \mathbf{t}^3)$. Recall that in the evaluation $Y(\mathfrak{gl}_4)$ -module $V(x)$, we have $T_b^a(u) = \delta_{ab} + e_{ba}(u - x)^{-1}$, thus

$$T_a^a(u)v = \frac{u - x + \Lambda^a}{u - x}v.$$

By Proposition 4.1,

$$\mathbb{B}_{\boldsymbol{\xi}}(\mathbf{t})v = \sum_{\mathbf{a}, \mathbf{b}} \left(\mathbb{T}^{[2]}(\mathbf{t}^2) \right)_{\mathbf{b}}^{\mathbf{a}} (\phi_2(\mathbf{t}^2) (\mathbb{B}_{\xi_1}^{(2)}(\mathbf{t}^1)))^{\mathbf{a}} (\psi_2(\mathbf{t}^2) (\mathbb{B}_{\xi_3}^{(2)}(\mathbf{t}^3)))^{\mathbf{b}-\mathbf{2}} v, \quad (6.1)$$

where the sum is taken over all sequences $\mathbf{a} = (a_1, a_2, \dots, a_{\xi_2})$, $\mathbf{b} = (b_1, b_2, \dots, b_{\xi_2})$, such that $a_i \in \{1, 2\}$, and $b_i \in \{3, 4\}$ for all $i = 1, \dots, \xi_2$.

Let ${}^\psi V(x)$ be the $Y(\mathfrak{gl}_2)$ -module obtained by pulling back the module $V(x)$ through the embedding ψ_2 . To compute $(\psi_2(\mathbf{t}^2)(\mathbb{B}_{\xi_3}^{(2)}(\mathbf{t}^3)))^{\mathbf{b}} v$, we take the weight function $\mathbb{B}_{\xi_3}^{(2)}(\mathbf{t}^3)(\mathbf{w}_1^{\otimes \xi_2} \otimes v)$ in the $Y(\mathfrak{gl}_2)$ -module $L(\mathbf{t}_1^2) \otimes \dots \otimes L(\mathbf{t}_{\xi_2}^2) \otimes {}^\psi V(x)$ and apply Proposition 5.2 for $k = \xi_2$. Then we obtain

$$\begin{aligned} & (\psi_2(\mathbf{t}^2)(\mathbb{B}_{\xi_3}^{(2)}(\mathbf{t}^3)))^{\mathbf{b}-2} v \\ &= \frac{1}{(\xi_3 - |I|)!} \overline{\text{Sym}}_{\mathbf{t}^3} \left[F_I(\mathbf{t}^3, \mathbf{t}^2) \prod_{m=1}^{|I|} \frac{t_m^3 - x + \Lambda^3}{t_m^3 - x} \prod_{r=|I|+1}^{\xi_3} \frac{1}{t_r^3 - x} \right] e_{43}^{\xi_3 - |I|} v, \end{aligned} \quad (6.2)$$

where the subset $I \subset \{1, \dots, \xi_2\}$ and the sequence $\mathbf{b} = (b_1, \dots, b_{\xi_2})$ are related as follows: $b_j = 3$ if $j \notin I$ and $b_j = 4$ if $j \in I$. Therefore, by formula (5.8) we have

$$(\psi_2(\mathbf{t}^2)(\mathbb{B}_{\xi_3}^{(2)}(\mathbf{t}^3)))^{\mathbf{b}-2} v = \prod_{r=1}^{\xi_3} \frac{1}{t_r^3 - x} V_I(\mathbf{t}^3, \mathbf{t}^2, x - \Lambda^3) e_{43}^{\xi_3 - |I|} v.$$

The next step is to compute $(\phi_2(\mathbf{t}^2)(\mathbb{B}_{\xi_1}^{(2)}(\mathbf{t}^1)))^{\mathbf{a}} e_{43}^{\xi_3 - |I|} v$. Notice that for any nonnegative integer m , we have

$$T_1^2(u) e_{43}^m v = 0, \quad T_2^2(u) e_{43}^m v = \frac{u - x + \Lambda^2}{u - x} e_{43}^m v, \quad T_1^1(u) e_{43}^m v = \frac{u - x + \Lambda^1}{u - x} e_{43}^m v.$$

Let ${}^\phi V(x)$ the $Y(\mathfrak{gl}_2)$ -module obtained by pulling back $V(x)$ through the embedding ϕ_2 . To compute the $(\phi_2(\mathbf{t}^2)(\mathbb{B}_{\xi_1}^{(2)}(\mathbf{t}^1)))^{\mathbf{a}} e_{43}^{\xi_3 - |I|} v$, we take the weight function $\mathbb{B}_{\xi_1}^{(2)}(\mathbf{t}^1)(e_{43}^{\xi_3 - |I|} v \otimes \mathbf{w}_2^{\otimes \xi_2})$ in the $Y(\mathfrak{gl}_2)$ -module ${}^\phi V(x) \otimes L(\mathbf{t}_{\xi_2}^2) \otimes \dots \otimes L(\mathbf{t}_1^2)$ and apply Proposition 5.3 for $k = \xi_2$. Then we obtain

$$\begin{aligned} & (\phi_2(\mathbf{t}^2)(\mathbb{B}_{\xi_1}^{(2)}(\mathbf{t}^1)))^{\mathbf{a}} e_{43}^{\xi_3 - |I|} v \\ &= \frac{1}{(\xi_1 - |J|)!} \overline{\text{Sym}}_{\mathbf{t}^1} \left[\tilde{F}_J(\mathbf{t}^1, \mathbf{t}^2) \prod_{m=1}^{|J|} \frac{t_m^1 - x + \Lambda^2}{t_m^1 - x} \prod_{r=|J|+1}^{\xi_1} \frac{1}{t_r^1 - x} \right] e_{21}^{\xi_1 - |J|} e_{43}^{\xi_3 - |I|} v, \end{aligned} \quad (6.3)$$

where the subset $J \subset \{1, \dots, \xi_2\}$ and the sequence $\mathbf{a} = (a_1, \dots, a_{\xi_2})$ are related as follows: $a_j = 1$ if $j \in J$ and $a_j = 2$ if $j \notin J$. Therefore, by formula (5.7) we have

$$(\phi_2(\mathbf{t}^2)(\mathbb{B}_{\xi_1}^{(2)}(\mathbf{t}^1)))^{\mathbf{a}} e_{43}^{\xi_3 - |I|} v = \prod_{r=1}^{\xi_1} \frac{1}{t_r^1 - x} \tilde{V}_J(\mathbf{t}^1, \mathbf{t}^2, x - \Lambda^2) e_{21}^{\xi_1 - |J|} e_{43}^{\xi_3 - |I|} v.$$

Finally, for the sequences \mathbf{a}, \mathbf{b} that are related to the sets I, J as above, we have

$$\left(\mathbb{T}^{[2]}(\mathbf{t}^2) \right)_{\mathbf{b}}^{\mathbf{a}} = \prod_{r=1}^{\xi_2} \frac{1}{t_r^2 - x} e_{32}^{\xi_2 - p - q + s} e_{31}^{q-s} e_{42}^{p-s} e_{41}^s, \quad (6.4)$$

where $p = |I|$, $q = |J|$, $s = |I \cap J|$. Now formula (5.10) follows from formulae (6.1)–(6.4).

7 Proof of Proposition 5.6

Consider the algebra \mathcal{A} generated by two commuting copies of the symmetric group S_k and rational functions of z_1, \dots, z_k subject to relations (7.1) below. We denote the copies of S_k

in \mathcal{A} by \dot{S}_k and \ddot{S}_k , and mark elements of \dot{S}_k and \ddot{S}_k by the corresponding dots, keeping the notation S_k without dots for the abstract symmetric group.

Let $\mathbf{z} = (z_1, \dots, z_k)$ and $\mathbf{z}^\sigma = (z_{\sigma(1)}, \dots, z_{\sigma(k)})$. The additional relations in \mathcal{A} are

$$\dot{\sigma} f(\mathbf{z}) = f(\mathbf{z}^\sigma) \dot{\sigma}, \quad \ddot{\tau} f(\mathbf{z}) = f(\mathbf{z}) \ddot{\tau}. \quad (7.1)$$

For $a = 1, \dots, k-1$, let $s_a \in S_k$ be the transposition of a and $a+1$. Consider the elements $\hat{s}_1, \dots, \hat{s}_k$ of \mathcal{A} ,

$$\hat{s}_a = \left(\frac{z_a - z_{a+1}}{z_a - z_{a+1} - 1} \ddot{s}_a - \frac{1}{z_a - z_{a+1} - 1} \right) \dot{s}_a. \quad (7.2)$$

It is straightforward to check that they satisfy the following relations:

$$\hat{s}_a \hat{s}_{a+1} \hat{s}_a = \hat{s}_{a+1} \hat{s}_a \hat{s}_{a+1}, \quad \hat{s}_a^2 = 1.$$

Therefore, the assignment $s_a \mapsto \hat{s}_a$ defines an algebra homomorphism $\mathbb{C}S_k \rightarrow \mathcal{A}$. For any $\sigma \in S_k$, we denote by $\hat{\sigma}$ the corresponding element of \mathcal{A} . Every element $\hat{\sigma}$ can be written in the following form:

$$\hat{\sigma} = \sum_{\tau \in S_k} X_{\sigma, \tau}(\mathbf{z}) \ddot{\tau} \dot{\sigma}, \quad (7.3)$$

where $X_{\sigma, \tau}(\mathbf{z})$ are functions of z_1, \dots, z_k .

Let $|\sigma|$ denote the length of $\sigma \in S_k$.

Lemma 7.1. *The functions $X_{\sigma, \tau}(\mathbf{z})$ have the following properties:*

$$X_{\sigma, \tau}(\mathbf{z}) = 0 \quad \text{if } |\tau| > |\sigma|, \quad (7.4)$$

$$X_{\sigma, \tau}(\mathbf{z}) = \delta_{\sigma, \tau} X_{\sigma, \sigma}(\mathbf{z}) \quad \text{if } |\tau| = |\sigma|, \quad (7.5)$$

$$X_{\sigma, \sigma}(\mathbf{z}) = \prod_{a < b, \sigma^{-1}(a) > \sigma^{-1}(b)} \frac{z_a - z_b}{z_a - z_b - 1}. \quad (7.6)$$

Proof. Formulae (7.4) and (7.5) follow from formula (7.2) by inspection. Formula (7.6) can be shown by induction on $|\sigma|$. ■

Denote by σ_0 the longest element of S_k , $\sigma_0(i) = k - i + 1$, $i = 1, \dots, k$. Let

$$\Phi(\mathbf{z}) = \prod_{a < b} \frac{z_a - z_b - 1}{z_a - z_b}.$$

Notice that

$$\Phi(\mathbf{z}) = \frac{1}{X_{\sigma_0, \sigma_0}(\mathbf{z})}. \quad (7.7)$$

Lemma 7.2. *One has*

$$\sum_{\lambda \in S_k} X_{\lambda, \rho}(\mathbf{z}) \Phi(\mathbf{z}^{\lambda \sigma_0}) X_{\sigma_0 \lambda^{-1}, \sigma_0 \tau^{-1}}(\mathbf{z}^{\lambda \sigma_0}) = \delta_{\rho, \tau}. \quad (7.8)$$

Proof. Since $\widehat{\sigma \tau} = \hat{\sigma} \hat{\tau}$, by formula (7.3) we have

$$X_{\sigma \tau, \rho}(\mathbf{z}) = \sum_{\pi} X_{\sigma, \pi}(\mathbf{z}) X_{\tau, \pi^{-1} \rho}(\mathbf{z}^\sigma). \quad (7.9)$$

Taking here $\rho = \sigma_0$, and using Lemma 7.1 and formula (7.7), we get

$$\sum_{\pi} X_{\sigma,\pi}(z) X_{\tau,\pi^{-1}\sigma_0}(z^\sigma) = \delta_{\sigma\tau,\sigma_0} \frac{1}{\Phi(z)}. \quad (7.10)$$

Replacing now z by $z^{\sigma^{-1}}$ in formula (7.10) and taking there $\tau = \mu^{-1}\sigma_0$, we get

$$\sum_{\pi} X_{\sigma,\pi}(z^{\sigma^{-1}}) X_{\mu^{-1}\sigma_0,\pi^{-1}\sigma_0}(z) = \delta_{\sigma,\mu} \frac{1}{\Phi(z^{\sigma^{-1}})}. \quad (7.11)$$

Formula (7.11) can be understood as the matrix equality $AB = C$ for $k! \times k!$ matrices A, B, C with entries labeled by permutations:

$$A_{\sigma,\pi} = X_{\sigma,\pi}(z^{\sigma^{-1}}), \quad B_{\pi,\mu} = X_{\mu^{-1}\sigma_0,\pi^{-1}\sigma_0}(z), \quad C_{\sigma,\mu} = \delta_{\sigma,\mu} \frac{1}{\Phi(z^{\sigma^{-1}})}.$$

Therefore, the product $BC^{-1}A$ equals the identity matrix, which can be written as follows:

$$\sum_{\mu} X_{\mu^{-1}\sigma_0,\pi^{-1}\sigma_0}(z) \Phi(z^{\mu^{-1}}) X_{\mu,\sigma}(z^{\mu^{-1}}) = \delta_{\pi,\sigma}.$$

After the substitution $\lambda = \mu^{-1}\sigma_0$, $\rho = \pi^{-1}\sigma_0$, $\tau = \sigma^{-1}\sigma_0$, we get formula (7.8). ■

Lemma 7.3. *One has*

$$X_{\mu,\sigma}(z^{s_a}) = \frac{z_a - z_{a+1}}{z_a - z_{a+1} + 1} X_{s_a\mu,s_a\sigma}(z) + \frac{1}{z_a - z_{a+1} + 1} X_{s_a\mu,\sigma}(z). \quad (7.12)$$

Proof. By formulae (7.2) and (7.3), we have

$$X_{s_a,s_a}(z) = \frac{z_a - z_{a+1}}{z_a - z_{a+1} - 1}, \quad X_{s_a,\text{id}}(z) = \frac{-1}{z_a - z_{a+1} - 1},$$

and $X_{s_a,\tau}(z) = 0$, otherwise. Therefore, by formula (7.9) we obtain

$$X_{s_a\pi,\sigma}(z) = \frac{z_a - z_{a+1}}{z_a - z_{a+1} - 1} X_{\pi,s_a\sigma}(z^{s_a}) - \frac{1}{z_a - z_{a+1} - 1} X_{\pi,\sigma}(z^{s_a}).$$

Replacing here z by z^{s_a} and making the substitution $\pi = s_a\mu$, we get formula (7.12). ■

Lemma 7.4. *One has*

$$X_{\mu,\sigma}(z^{s_a\mu^{-1}}) = \frac{z_a - z_{a+1}}{z_a - z_{a+1} - 1} X_{\mu s_a,\sigma s_a}(z^{s_a\mu^{-1}}) - \frac{1}{z_a - z_{a+1} - 1} X_{\mu s_a,\sigma}(z^{s_a\mu^{-1}}). \quad (7.13)$$

Proof. By formula (7.9), we have

$$X_{\mu s_a,\sigma}(z) = \sum_{\pi} X_{\mu,\pi}(z) X_{s_a,\pi^{-1}\sigma}(z^\mu).$$

Thus

$$X_{\mu s_a,\sigma}(z) = \sum_{\pi} X_{\mu,\pi}(z) X_{s_a,\pi^{-1}\sigma}(z^\mu) = X_{\mu,\sigma}(z) X_{s_a,\text{id}}(z^\mu) + X_{\mu,\sigma s_a}(z) X_{s_a,s_a}(z^\mu).$$

Replacing here z by $z^{\mu^{-1}}$, we get

$$X_{\mu s_a,\sigma}(z^{\mu^{-1}}) = X_{\mu,\sigma}(z^{\mu^{-1}}) X_{s_a,\text{id}}(z) + X_{\mu,\sigma s_a}(z^{\mu^{-1}}) X_{s_a,s_a}(z).$$

Substituting now μ with μs_a , we obtain (7.13). ■

For $\sigma \in S_n$ and a subset $I = \{i_1, \dots, i_m\} \subset \{1, \dots, n\}$, denote

$$\sigma(I) = \{\sigma(i_1), \dots, \sigma(i_m)\}.$$

Recall the functions $V_I(\mathbf{t}, \mathbf{z}, y)$, $\tilde{V}_J(\mathbf{t}, \mathbf{z}, y)$, see formulae (5.4), (5.7) and (5.6), (5.8).

Lemma 7.5. *For each $a = 1, \dots, k-1$, we have*

$$V_I(\mathbf{t}, \mathbf{z}^{s_a}, y) = \frac{z_{a+1} - z_a}{z_{a+1} - z_a - 1} V_{s_a(I)}(\mathbf{t}, \mathbf{z}, y) - \frac{1}{z_{a+1} - z_a - 1} V_I(\mathbf{t}, \mathbf{z}, y), \quad (7.14)$$

$$\tilde{V}_I(\mathbf{t}, \mathbf{z}^{s_a}, y) = \frac{z_a - z_{a+1}}{z_a - z_{a+1} - 1} \tilde{V}_{s_a(I)}(\mathbf{t}, \mathbf{z}, y) - \frac{1}{z_a - z_{a+1} - 1} \tilde{V}_I(\mathbf{t}, \mathbf{z}, y). \quad (7.15)$$

Proof. By the structure of formulae (5.4) and (5.7) for the function $V_I(\mathbf{t}, \mathbf{z}, y)$, it is enough to prove formula (7.14) for $k = 2$. In this case, the statement follows from the identities

$$\begin{aligned} 1 &= \frac{z' - z}{z' - z - 1} - \frac{1}{z' - z - 1}, \\ \frac{1}{t - z'} \cdot \frac{t - z + 1}{t - z} &= \frac{z' - z}{z' - z - 1} \cdot \frac{1}{t - z'} - \frac{1}{z' - z - 1} \cdot \frac{1}{t - z} \cdot \frac{t - z' + 1}{t - z'}, \\ (t' - z)(t - z' + 1)(t - t' - 1) - (t - z)(t' - z' + 1)(t' - t - 1) \\ &= (t' - z')(t - z + 1)(t - t' - 1) - (t - z')(t' - z + 1)(t' - t - 1). \end{aligned}$$

The proof of (7.15) is similar by using formulae (5.6) and (5.8) for functions $\tilde{V}_J(\mathbf{t}, \mathbf{z}, y)$. ■

The statement of Proposition 5.6 is given by formula (5.11). It can be written as follows:

$$\begin{aligned} &\sum_{(I, J) \in \mathcal{S}_{p, q, r, k}} V_I(\mathbf{t}^3, \mathbf{z}, x - \Lambda^3) \tilde{V}_J(\mathbf{t}^1, \mathbf{z}, x - \Lambda^2) \\ &= \frac{1}{C_{p, q, r, k}} \sum_{\sigma \in S_k} V_{\sigma_0(I_0)}(\mathbf{t}^3, \mathbf{z}^\sigma, x - \Lambda^3) \tilde{V}_{J_0}(\mathbf{t}^1, \mathbf{z}^{\sigma \sigma_0}, x - \Lambda^2) \Phi(\mathbf{z}^\sigma), \end{aligned} \quad (7.16)$$

where $\mathcal{S}_{p, q, r, k}$ in the left-hand side is the set of all pairs of subsets I, J of $\{1, \dots, k\}$, such that $|I| = p$, $|J| = q$, $|I \cap J| = r$, and we use $k = \xi_2$, $\mathbf{z} = \mathbf{t}^2$. In the right-hand side,

$$C_{p, q, r, k} = (p - r)!(q - r)!r!(k - p - q - r)!$$

and (I_0, J_0) is any fixed pair from $\mathcal{S}_{p, q, r, k}$. We also expanded $\overline{\text{Sym}}_{\mathbf{t}^2}$ according to formulae (5.1) and (5.2), and observed that

$$\check{I}_0 = \sigma_0(I_0), \quad \check{\mathbf{t}}^2 = \mathbf{z}^{\sigma_0}.$$

In the rest of the proof, we will suppress the arguments $\mathbf{t}^1, \mathbf{t}^3, x - \Lambda^2, x - \Lambda^3$ because they are the same in both sides of formula (7.16) and will never be changed in the reasoning.

Notice that every pair $(I, J) \in \mathcal{S}_{p, q, r, k}$ can be obtained from an arbitrary fixed pair $(I_0, J_0) \in \mathcal{S}_{p, q, r, k}$ by the action of the symmetric group S_k . Therefore, the left-hand side of the formula (7.16) can be written in the following way:

$$\sum_{(I, J) \in \mathcal{S}_{p, q, r, k}} \tilde{V}_J(\mathbf{z}) V_I(\mathbf{z}) = \frac{1}{C_{p, q, r, k}} \sum_{\sigma \in S_k} \tilde{V}_{\sigma(J_0)}(\mathbf{z}) V_{\sigma(I_0)}(\mathbf{z}). \quad (7.17)$$

Using Lemma 7.2, we get

$$\begin{aligned} & \sum_{\sigma \in S_k} \tilde{V}_{\sigma(J_0)}(\mathbf{z}) V_{\sigma(I_0)}(\mathbf{z}) \\ &= \sum_{\sigma, \pi, \tau \in S_k} \tilde{V}_{\sigma(J_0)}(\mathbf{z}) X_{\pi, \sigma}(\mathbf{z}) \Phi(\mathbf{z}^{\pi \sigma_0}) X_{\sigma_0 \pi^{-1}, \sigma_0 \tau^{-1}}(\mathbf{z}^{\pi \sigma_0}) V_{\tau(I_0)}(\mathbf{z}), \end{aligned} \quad (7.18)$$

since from formula (7.8)

$$\sum_{\pi \in S_k} X_{\pi, \sigma}(\mathbf{z}) \Phi(\mathbf{z}^{\pi \sigma_0}) X_{\sigma_0 \pi^{-1}, \sigma_0 \tau^{-1}}(\mathbf{z}^{\pi \sigma_0}) = \delta_{\sigma, \tau}.$$

Lemma 7.6. *We have*

$$\sum_{\sigma \in S_k} \tilde{V}_{\sigma(J_0)}(\mathbf{z}) X_{\pi, \sigma}(\mathbf{z}) = \tilde{V}_{J_0}(\mathbf{z}^\pi). \quad (7.19)$$

Proof. We will use induction on the length of the permutation π . For $\pi = \text{id}$, formula (7.19) is clear, and for $\pi = s_a$ with some $a = 1, \dots, k-1$, formula (7.19) coincides with formula (7.15). For the induction step, we find a such that $|s_a \pi| = |\pi| - 1$, and denote $\rho = s_a \pi$. Then by the induction assumption

$$\sum_{\sigma} \tilde{V}_{\sigma(J_0)}(\mathbf{z}) X_{\rho, \sigma}(\mathbf{z}) = \tilde{V}_{J_0}(\mathbf{z}^\rho).$$

Replacing here \mathbf{z} by \mathbf{z}^{s_a} , we get

$$\sum_{\sigma} \tilde{V}_{\sigma(J_0)}(\mathbf{z}^{s_a}) X_{\rho, \sigma}(\mathbf{z}^{s_a}) = \tilde{V}_{J_0}(\mathbf{z}^{s_a \rho}) = \tilde{V}_{J_0}(\mathbf{z}^\pi). \quad (7.20)$$

Using formulae (7.12) and (7.15), the left-hand side of (7.20) becomes

$$\begin{aligned} & \sum_{\sigma} \frac{(z_a - z_{a+1})^2}{(z_a - z_{a+1})^2 - 1} \tilde{V}_{s_a \sigma(J_0)}(\mathbf{z}) X_{s_a \rho, s_a \sigma}(\mathbf{z}) \\ &+ \sum_{\sigma} \frac{z_a - z_{a+1}}{(z_a - z_{a+1})^2 - 1} \tilde{V}_{s_a \sigma(J_0)}(\mathbf{z}) X_{s_a \rho, \sigma}(\mathbf{z}) \\ &- \sum_{\sigma} \frac{z_a - z_{a+1}}{(z_a - z_{a+1})^2 - 1} \tilde{V}_{\sigma(J_0)}(\mathbf{z}) X_{s_a \rho, s_a \sigma}(\mathbf{z}) \\ &- \sum_{\sigma} \frac{1}{(z_a - z_{a+1})^2 - 1} \tilde{V}_{\sigma(J_0)}(\mathbf{z}) X_{s_a \rho, \sigma}(\mathbf{z}). \end{aligned}$$

Changing the summation index in the first and second sums from σ to $s_a \sigma$, we observe that the second and third sums cancel each other, while the first and forth sums combine together and simplify to the expression

$$\sum_{\sigma} \tilde{V}_{\sigma(J_0)}(\mathbf{z}) X_{s_a \rho, \sigma}(\mathbf{z}) = \sum_{\sigma} \tilde{V}_{\sigma(J_0)}(\mathbf{z}) X_{\pi, \sigma}(\mathbf{z}),$$

which appears in the left-hand side of formula (7.19). ■

Lemma 7.7. *We have*

$$\sum_{\tau} V_{\tau(I_0)}(\mathbf{z}) X_{\sigma_0 \pi^{-1}, \sigma_0 \tau^{-1}}(\mathbf{z}^{\pi \sigma_0}) = V_{\sigma_0(I_0)}(\mathbf{z}^{\pi \sigma_0}). \quad (7.21)$$

Proof. Recall the notation $\sigma_0(I_0) = \check{I}_0$. Transform formula (7.21) by making the substitutions $\mu = \sigma_0\pi^{-1}$, $\sigma = \sigma_0\tau^{-1}$,

$$\sum_{\sigma} V_{\sigma^{-1}(\check{I}_0)}(z) X_{\mu,\sigma}(z^{\mu^{-1}}) = V_{\check{I}_0}(z^{\mu^{-1}}). \quad (7.22)$$

The rest of the proof is analogous to that of Lemma 7.6.

To prove formula (7.22), we will use induction on the length of μ . For $\mu = \text{id}$, formula (7.22) is clear, and for $\mu = s_a$ with some $a = 1, \dots, k-1$, formula (7.22) coincides with formula (7.14). For the induction step, we find a such that $|\mu s_a| = |\mu| - 1$, and denote $\rho = \mu s_a$. Then by the induction assumption,

$$\sum_{\sigma} V_{\sigma^{-1}(\check{I}_0)}(z) X_{\rho,\sigma}(z^{\rho^{-1}}) = V_{\check{I}_0}(z^{\rho^{-1}}),$$

and replacing here z by z^{s_a} , we get

$$\sum_{\sigma} V_{\sigma^{-1}(\check{I}_0)}(z^{s_a}) X_{\rho,\sigma}(z^{s_a\rho^{-1}}) = V_{\check{I}_0}(z^{\mu^{-1}}) = V_{\check{I}_0}(z^{s_a\rho^{-1}}). \quad (7.23)$$

Using formulae (7.13), (7.14), the left-hand of (7.23) side becomes

$$\begin{aligned} & \sum_{\sigma} \frac{(z_{a+1} - z_a)^2}{(z_{a+1} - z_a)^2 - 1} V_{s_a\sigma^{-1}(\check{I}_0)}(z) X_{\rho s_a, \sigma s_a}(z^{s_a\rho^{-1}}) \\ & - \sum_{\sigma} \frac{z_a - z_{a+1}}{(z_a - z_{a+1})^2 - 1} V_{s_a\sigma^{-1}(\check{I}_0)}(z) X_{\rho s_a, \sigma}(z^{s_a\rho^{-1}}) \\ & + \sum_{\sigma} \frac{z_a - z_{a+1}}{(z_a - z_{a+1})^2 - 1} V_{\sigma^{-1}(\check{I}_0)}(z) X_{\rho s_a, \sigma s_a}(z^{s_a\rho^{-1}}) \\ & - \sum_{\sigma} \frac{1}{(z_{a+1} - z_a)^2 - 1} V_{\sigma^{-1}(\check{I}_0)}(z) X_{\rho s_a, \sigma}(z^{s_a\rho^{-1}}). \end{aligned}$$

Changing the summation index in the first and second sums from σ to σs_a , we observe that the second and third sums cancel each other, while the first and the forth sums combine together and simplify to the expression

$$\sum_{\sigma} V_{\sigma^{-1}(\check{I}_0)}(z) X_{\rho s_a, \sigma}(z^{s_a\rho^{-1}}) = \sum_{\sigma} V_{\sigma^{-1}(\check{I}_0)}(z) X_{\mu, \sigma}(z^{\mu^{-1}}),$$

which appears in the left-hand side of formula (7.21). ■

Using Lemmas 7.6 and 7.7, we evaluate the sums over σ and τ in the right-hand side of the formula (7.18) and get the equality

$$\sum_{\sigma \in S_k} \tilde{V}_{\sigma(J_0)}(z) V_{\sigma(I_0)}(z) = \sum_{\pi} V_{\sigma_0(I_0)}(z^{\pi\sigma_0}) \tilde{V}_{J_0}(z^{\pi}) \Phi(z^{\pi\sigma_0}). \quad (7.24)$$

Using formula (7.17) in the left-hand side and making the substitution $\pi = \sigma\sigma_0$ in the right-hand side, we obtain that (7.24) can be written as

$$\sum_{(I,J) \in \mathcal{S}_{p,q,r,k}} \tilde{V}_J(z) V_I(z) = \frac{1}{C_{p,q,r,k}} \sum_{\sigma} V_{\sigma_0(I_0)}(z^{\sigma}) \tilde{V}_{J_0}(z^{\sigma\sigma_0}) \Phi(z^{\sigma}),$$

which is formula (7.16). Proposition 5.6 is proved.

A Proof of Proposition 5.1

In this appendix, we will consider only the algebra $Y(\mathfrak{gl}_2)$ and, for convenience, we will not write the superscript $\langle 2 \rangle$. We will use the commutation relations

$$\begin{aligned} T_{11}(u)T_{11}(t) &= T_{11}(t)T_{11}(u), & T_{12}(u)T_{12}(t) &= T_{12}(t)T_{12}(u), \\ T_{22}(u)T_{22}(t) &= T_{22}(t)T_{22}(u), \end{aligned} \quad (\text{A.1})$$

$$T_{11}(u)T_{12}(t) = \frac{u-t-1}{u-t}T_{12}(t)T_{11}(u) + \frac{1}{u-t}T_{12}(u)T_{11}(t), \quad (\text{A.2})$$

$$T_{22}(u)T_{12}(t) = \frac{u-t+1}{u-t}T_{12}(t)T_{22}(u) - \frac{1}{u-t}T_{12}(u)T_{22}(t), \quad (\text{A.3})$$

following from the defining relations in $Y(\mathfrak{gl}_2)$, see (2.4). We will also use the next statement.

Proposition A.1. *One has*

$$\begin{aligned} T_{11}(u)T_{12}(t_1) \cdots T_{12}(t_k) &= \prod_{i=1}^k \frac{u-t_i-1}{u-t_i} T_{12}(t_1) \cdots T_{12}(t_k) T_{11}(u) \\ &+ \sum_{l=1}^k \frac{1}{u-t_l} \prod_{\substack{m=1 \\ m \neq l}}^k \frac{t_l-t_m-1}{t_l-t_m} T_{12}(t_1) \cdots T_{12}(t_{l-1}) T_{12}(t_{l+1}) \cdots T_{12}(t_k) T_{12}(u) T_{11}(t_l), \end{aligned} \quad (\text{A.4})$$

$$\begin{aligned} T_{22}(u)T_{12}(t_1) \cdots T_{12}(t_k) &= \prod_{i=1}^k \frac{u-t_i+1}{u-t_i} T_{12}(t_1) \cdots T_{12}(t_k) T_{22}(u) \\ &- \sum_{l=1}^k \frac{1}{u-t_l} \prod_{\substack{m=1 \\ m \neq l}}^k \frac{t_l-t_m+1}{t_l-t_m} T_{12}(t_1) \cdots T_{12}(t_{l-1}) T_{12}(t_{l+1}) \cdots T_{12}(t_k) T_{12}(u) T_{22}(t_l). \end{aligned} \quad (\text{A.5})$$

Proof. The statement goes back to [23]. We will prove it by induction on k . Consider formula (A.4). The statement for $k=1$ is given by formula (A.2). We use the induction assumption to move $T_{11}(u)$ through the product $T_{12}(t_1) \cdots T_{12}(t_{k-1})$:

$$\begin{aligned} T_{11}(u)T_{12}(t_1) \cdots T_{12}(t_{k-1})T_{12}(t_k) &= \prod_{i=1}^{k-1} \frac{u-t_i-1}{u-t_i} T_{12}(t_1) \cdots T_{12}(t_{k-1}) T_{11}(u) T_{12}(t_k) \\ &+ \sum_{l=1}^{k-1} \frac{1}{u-t_l} \prod_{\substack{m=1 \\ m \neq l}}^{k-1} \frac{t_l-t_m-1}{t_l-t_m} T_{12}(t_1) \cdots T_{12}(t_{l-1}) T_{12}(t_{l+1}) \cdots T_{12}(t_{k-1}) T_{12}(u) T_{11}(t_l) T_{12}(t_k). \end{aligned} \quad (\text{A.6})$$

Then we apply (A.2) to the product $T_{11}(u)T_{12}(t_k)$ and $T_{11}(t_l)T_{12}(t_k)$ and the right-hand side of (A.6) becomes

$$\begin{aligned} &\prod_{i=1}^k \frac{u-t_i-1}{u-t_i} T_{12}(t_1) \cdots T_{12}(t_{k-1}) T_{12}(t_k) T_{11}(u) \\ &+ \frac{1}{u-t_k} \prod_{i=1}^{k-1} \frac{u-t_i-1}{u-t_i} T_{12}(t_1) \cdots T_{12}(t_{k-1}) T_{12}(u) T_{11}(t_k) \\ &+ \sum_{l=1}^{k-1} \frac{1}{u-t_l} \prod_{\substack{m=1 \\ m \neq l}}^k \frac{t_l-t_m-1}{t_l-t_m} \\ &\quad \times T_{12}(t_1) \cdots T_{12}(t_{l-1}) T_{12}(t_{l+1}) \cdots T_{12}(t_{k-1}) T_{12}(u) T_{12}(t_k) T_{11}(t_l) \end{aligned}$$

$$\begin{aligned}
& + \sum_{l=1}^{k-1} \frac{1}{u - t_l} \frac{1}{t_l - t_k} \prod_{\substack{m=1 \\ m \neq l}}^{k-1} \frac{t_l - t_m - 1}{t_l - t_m} \\
& \quad \times T_{12}(t_1) \cdots T_{12}(t_{l-1}) T_{12}(t_{l+1}) \cdots T_{12}(t_{k-1}) T_{12}(u) T_{12}(t_l) T_{11}(t_k).
\end{aligned}$$

The first term here coincide with the first term in the right-hand side of formula (A.4). The third term here is the second term of (A.4) without $l = k$ summand. We also used that $T_{12}(u)$ and $T_{12}(t_k)$ commute, see (A.1). The second and forth summands combine into the product

$$\frac{1}{u - t_k} \prod_{m=1}^{k-1} \frac{t_k - t_m - 1}{t_k - t_m} T_{12}(t_1) \cdots T_{12}(t_{l-1}) T_{12}(t_{l+1}) \cdots T_{12}(t_{k-1}) T_{12}(u) T_{11}(t_l), \quad (\text{A.7})$$

using the following identity:

$$\frac{1}{u - t_k} \prod_{i=1}^{k-1} \frac{u - t_i - 1}{u - t_i} + \sum_{l=1}^{k-1} \frac{1}{(u - t_l)(t_l - t_k)} \prod_{\substack{m=1 \\ m \neq l}}^{k-1} \frac{t_l - t_m - 1}{t_l - t_m} = \frac{1}{u - t_k} \prod_{m=1}^{k-1} \frac{t_k - t_m - 1}{t_k - t_m}.$$

The product (A.7) is exactly the summand with $l = k$ of the second term in (A.4). Formula (A.4) is proved.

The proof of formula (A.5) is similar to that of formula (A.4) with relation (A.3) used instead of (A.2). \blacksquare

Recall that for the \mathfrak{gl}_2 case we have

$$\mathbb{B}_\xi(\mathbf{t}) = T_{12}(t_1) \cdots T_{12}(t_\xi),$$

and thus Proposition 5.1 can be rewritten as follows.

Proposition A.2. *Let ξ be a nonnegative integer and $\mathbf{t} = (t_1, \dots, t_\xi)$. Then*

$$\begin{aligned}
& \Delta(T_{12}(t_1) \cdots T_{12}(t_\xi)) \\
& = \sum_{\eta=0}^{\xi} \frac{1}{(\xi - \eta)! \eta!} \overline{\text{Sym}}_{\mathbf{t}} \left[\left(\prod_{i=1}^{\eta} T_{12}(t_i) \otimes \prod_{j=\eta+1}^{\xi} T_{12}(t_j) \right) \left(\prod_{k=\eta+1}^{\xi} T_{22}(t_k) \otimes \prod_{l=1}^{\eta} T_{11}(t_l) \right) \right].
\end{aligned} \quad (\text{A.8})$$

Remark. Notice that according to (A.1), the factors in each of the large products commute among themselves, so the order of the factors is irrelevant. (A.8).

Proof. Consider the summand from the right-hand side of (A.8) with a given η ,

$$F_{\eta, \xi - \eta}(\mathbf{t}) = \overline{\text{Sym}}_{\mathbf{t}} \left[\left(\prod_{i=1}^{\eta} T_{12}(t_i) \otimes \prod_{j=\eta+1}^{\xi} T_{12}(t_j) \right) \left(\prod_{k=\eta+1}^{\xi} T_{22}(t_k) \otimes \prod_{l=1}^{\eta} T_{11}(t_l) \right) \right]. \quad (\text{A.9})$$

Let

$$\begin{aligned}
P_{\eta, \xi - \eta}(\mathbf{t}) &= \prod_{1 \leq i \leq \eta < j \leq \xi} \frac{t_i - t_j - 1}{t_i - t_j} \\
&\quad \times \left(\prod_{i=1}^{\eta} T_{12}(t_i) \otimes \prod_{j=\eta+1}^{\xi} T_{12}(t_j) \right) \left(\prod_{k=\eta+1}^{\xi} T_{22}(t_k) \otimes \prod_{l=1}^{\eta} T_{11}(t_l) \right), \\
U_{\eta, \xi - \eta}(\mathbf{t}) &= \prod_{1 \leq i < j \leq \eta} \frac{t_i - t_j - 1}{t_i - t_j} \prod_{\eta+1 \leq i < j \leq \xi} \frac{t_i - t_j - 1}{t_i - t_j}, \quad \mathbf{t}^\sigma = (t_{\sigma(1)}, \dots, t_{\sigma(\xi)}).
\end{aligned}$$

Using this notation, formula (A.9) can be written as

$$F_{\eta, \xi-\eta}(\mathbf{t}) = \sum_{\sigma \in S_\xi} U_{\eta, \xi-\eta}(\mathbf{t}^\sigma) P_{\eta, \xi-\eta}(\mathbf{t}^\sigma).$$

Observe that $F_{\eta, \xi-\eta}(\mathbf{t})$ is symmetric in t_1, \dots, t_ξ . Denote by $S_\eta \times S_{\xi-\eta}$ the subgroup of S_ξ stabilizing the subsets $\{1, \dots, \eta\}$ and $\{\eta+1, \dots, \xi\}$. We have

$$\begin{aligned} F_{\eta, \xi-\eta}(\mathbf{t}) &= \frac{1}{\eta!(\xi-\eta)!} \sum_{\tau \in S_\eta \times S_{\xi-\eta}} F_{\eta, \xi-\eta}(\mathbf{t}^\tau) \\ &= \frac{1}{\eta!(\xi-\eta)!} \sum_{\tau \in S_\eta \times S_{\xi-\eta}} \sum_{\sigma \in S_\xi} U_{\eta, \xi-\eta}(\mathbf{t}^{\tau\sigma}) P_{\eta, \xi-\eta}(\mathbf{t}^{\tau\sigma}). \end{aligned}$$

Changing the summation variable in the inner sum, $\sigma = \tau^{-1}\rho\tau$, and using the fact that $P_{\eta, \xi-\eta}(\mathbf{t}^{\rho\tau}) = P_{\eta, \xi-\eta}(\mathbf{t}^\rho)$ for all $\tau \in S_\eta \times S_{\xi-\eta}$, we get

$$\begin{aligned} F_{\eta, \xi-\eta}(\mathbf{t}) &= \frac{1}{\eta!(\xi-\eta)!} \sum_{\tau \in S_\eta \times S_{\xi-\eta}} \sum_{\rho \in S_\xi} U_{\eta, \xi-\eta}(\mathbf{t}^{\rho\tau}) P_{\eta, \xi-\eta}(\mathbf{t}^{\rho\tau}) \\ &= \frac{1}{\eta!(\xi-\eta)!} \sum_{\rho \in S_\xi} P_{\eta, \xi-\eta}(\mathbf{t}^\rho) \sum_{\tau \in S_\eta \times S_{\xi-\eta}} U_{\eta, \xi-\eta}(\mathbf{t}^{\rho\tau}). \end{aligned}$$

Furthermore, using the identity

$$\sum_{\tau \in S_n} \prod_{1 \leq i < j \leq n} \frac{x_{\tau(i)} - x_{\tau(j)} - 1}{x_{\tau(i)} - x_{\tau(j)}} = n!, \quad (\text{A.10})$$

we obtain that $\sum_{\tau \in S_\eta \times S_{\xi-\eta}} U_{\eta, \xi-\eta}(\mathbf{t}^{\rho\tau}) = \eta!(\xi-\eta)!$ and

$$F_{\eta, \xi-\eta}(\mathbf{t}) = \sum_{\rho \in S_\xi} P_{\eta, \xi-\eta}(\mathbf{t}^\rho). \quad (\text{A.11})$$

Using formula (A.11), the statement of Proposition A.2 can be formulated as follows:

$$\Delta(T_{12}(t_1) \cdots T_{12}(t_\xi)) = \sum_{\eta=0}^{\xi} \frac{1}{(\xi-\eta)!\eta!} \sum_{\rho \in S_\xi} P_{\eta, \xi-\eta}(\mathbf{t}^\rho). \quad (\text{A.12})$$

We will prove this formula using the induction on ξ . The base of induction at $\xi = 1$ is given by formula (2.5):

$$\Delta(T_{12}(t_1)) = T_{12}(t_1) \otimes T_{11}(t_1) + T_{22}(t_1) \otimes T_{12}(t_1). \quad (\text{A.13})$$

To make the induction step, we use that

$$\Delta(\mathbb{B}_\xi(\mathbf{t})) = \Delta(T_{12}(t_1)) \Delta(T_{12}(t_2) \cdots T_{12}(t_\xi)), \quad (\text{A.14})$$

expand the first factor according to (A.13), and apply the induction assumption to expand the second factor. Denote by $S'_{\xi-1} \subset S_\xi$ the subgroup of permutations ρ , such that $\rho(1) = 1$. Then the right-hand side of formula (A.14) becomes

$$\begin{aligned} T_{12}(t_1) \otimes T_{11}(t_1) &\sum_{\eta=1}^{\xi} \frac{1}{(\xi-\eta)!(\eta-1)!} \sum_{\tau \in S'_{\xi-1}} P_{\eta-1, \xi-\eta}(t_{\tau(2)}, \dots, t_{\tau(\xi)}) \\ &+ T_{22}(t_1) \otimes T_{12}(t_1) \sum_{\eta=0}^{\xi-1} \frac{1}{(\xi-1-\eta)!\eta!} \sum_{\rho \in S'_{\xi-1}} P_{\eta, \xi-\eta-1}(t_{\rho(2)}, \dots, t_{\rho(\xi)}), \end{aligned} \quad (\text{A.15})$$

where in the first term we shifted the summation variable of the exterior sum. Using the definition of $P_{\eta, \xi-1-\eta}(\mathbf{t})$ and $P_{\eta-1, \xi-\eta}(\mathbf{t})$, we further expand expression (A.15):

$$\begin{aligned} & \sum_{\eta=1}^{\xi} \frac{1}{(\xi-\eta)!(\eta-1)!} \sum_{\tau \in S'_{\xi-1}} \prod_{1 < i \leq \eta < j \leq \xi} \frac{t_{\tau(i)} - t_{\tau(j)} - 1}{t_{\tau(i)} - t_{\tau(j)}} \\ & \quad \times \left(T_{12}(t_1) \prod_{i=2}^{\eta} T_{12}(t_{\tau(i)}) \otimes T_{11}(t_1) \prod_{j=\eta+1}^{\xi} T_{12}(t_{\tau(j)}) \right) \left(\prod_{k=\eta+1}^{\xi} T_{22}(t_{\tau(k)}) \otimes \prod_{l=2}^{\eta} T_{11}(t_{\tau(l)}) \right) \\ & + \sum_{\eta=0}^{\xi-1} \frac{1}{(\xi-\eta-1)!\eta!} \sum_{\rho \in S'_{\xi-1}} \prod_{1 < i \leq \eta+1 < j \leq \xi} \frac{t_{\rho(i)} - t_{\rho(j)} - 1}{t_{\rho(i)} - t_{\rho(j)}} \\ & \quad \times \left(T_{22}(t_1) \prod_{i=2}^{\eta+1} T_{12}(t_{\rho(i)}) \otimes T_{12}(t_1) \prod_{j=\eta+2}^{\xi} T_{12}(t_{\rho(j)}) \right) \left(\prod_{k=\eta+2}^{\xi} T_{22}(t_{\rho(k)}) \otimes \prod_{l=2}^{\eta+1} T_{11}(t_{\rho(l)}) \right). \end{aligned}$$

In the first term, we move $T_{11}(t_1)$ through the product $\prod_{j=\eta+1}^{\xi} T_{12}(t_{\tau(j)})$ using formula (A.4):

$$\begin{aligned} T_{11}(t_1) \prod_{j=\eta+1}^{\xi} T_{12}(t_{\tau(j)}) &= \left(\prod_{j=\eta+1}^{\xi} \frac{t_1 - t_{\tau(j)} - 1}{t_1 - t_j} T_{12}(t_{\tau(j)}) \right) T_{11}(t_1) \\ &+ \sum_{p=\eta+1}^{\xi} \frac{1}{t_1 - t_{\tau(p)}} \left(\prod_{\substack{j=\eta+1 \\ j \neq p}}^{\xi} \frac{t_{\tau(p)} - t_{\tau(j)} - 1}{t_{\tau(p)} - t_{\tau(j)}} T_{12}(t_{\tau(j)}) \right) T_{12}(t_1) T_{11}(t_{\tau(p)}). \end{aligned}$$

Similarly, in the second term we move $T_{22}(t_1)$ through the product $\prod_{m=2}^{\eta} T_{12}(t_{\rho(m)})$ using formula (A.5):

$$\begin{aligned} T_{22}(t_1) \prod_{i=2}^{\eta+1} T_{12}(t_{\rho(i)}) &= \left(\prod_{i=2}^{\eta+1} \frac{t_1 - t_{\rho(i)} + 1}{t_1 - t_{\rho(i)}} T_{12}(t_{\rho(m)}) \right) T_{22}(t_1) \\ &- \sum_{s=2}^{\eta+1} \frac{1}{t_1 - t_s} \left(\prod_{\substack{i=1 \\ i \neq s}}^{\eta+1} \frac{t_{\rho(s)} - t_{\rho(i)} + 1}{t_{\rho(s)} - t_{\rho(i)}} T_{12}(t_{\rho(i)}) \right) T_{12}(t_1) T_{22}(t_{\rho(s)}), \end{aligned}$$

After all, the right-hand side of (A.14) becomes a sum of four terms:

$$\Delta(\mathbb{B}_{\xi}(\mathbf{t})) = Y_1(\mathbf{t}) + Y_2(\mathbf{t}) + Y_3(\mathbf{t}) + Y_4(\mathbf{t}),$$

where

$$\begin{aligned} Y_1(\mathbf{t}) &= \sum_{\eta=1}^{\xi} \frac{1}{(\xi-\eta)!(\eta-1)!} \sum_{\tau \in S'_{\xi-1}} \left[\prod_{l=\eta+1}^{\xi} \frac{t_1 - t_{\tau(l)} - 1}{t_1 - t_{\tau(l)}} \prod_{1 < i \leq \eta < j \leq \xi} \frac{t_{\tau(i)} - t_{\tau(j)} - 1}{t_{\tau(i)} - t_{\tau(j)}} \right. \\ & \quad \times \left(T_{12}(t_1) \prod_{i=2}^{\eta} T_{12}(t_{\tau(i)}) \otimes \prod_{j=\eta+1}^{\xi} T_{12}(t_{\tau(j)}) \right) \\ & \quad \times \left. \left(\prod_{k=\eta+1}^{\xi} T_{22}(t_{\tau(k)}) \otimes T_{11}(t_1) \prod_{l=2}^{\eta} T_{11}(t_{\tau(l)}) \right) \right], \end{aligned}$$

$$\begin{aligned}
Y_2(\mathbf{t}) &= \sum_{\eta=1}^{\xi-1} \frac{1}{(\xi-\eta)!(\eta-1)!} \\
&\times \sum_{\tau \in S'_{\xi-1}} \sum_{l=\eta+1}^{\xi} \left[\frac{1}{t_1 - t_{\tau(l)}} \prod_{1 < i \leq \eta < j \leq \xi} \frac{t_{\tau(i)} - t_{\tau(j)} - 1}{t_{\tau(i)} - t_{\tau(j)}} \prod_{\substack{m=\eta+1 \\ m \neq l}}^{\xi} \frac{t_{\tau(l)} - t_{\tau(m)} - 1}{t_{\tau(l)} - t_{\tau(m)}} \right. \\
&\quad \times \left(T_{12}(t_1) \prod_{k=2}^{\eta} T_{12}(t_{\tau(k)}) \otimes T_{12}(t_1) \prod_{\substack{m=\eta+1 \\ m \neq l}}^{\xi} T_{12}(t_{\tau(m)}) \right) \\
&\quad \times \left. \left(\prod_{i=\eta+1}^{\xi} T_{22}(t_{\tau(i)}) \otimes T_{11}(t_{\tau(l)}) \prod_{j=2}^{\eta} T_{11}(t_{\tau(j)}) \right) \right]. \\
Y_3(\mathbf{t}) &= \sum_{\eta=0}^{\xi-1} \frac{1}{(\xi-\eta-1)!\eta!} \sum_{\rho \in S'_{\xi-1}} \left[\prod_{m=2}^{\eta+1} \frac{t_{\rho(m)} - t_1 - 1}{t_{\rho(m)} - t_1} \prod_{1 < i \leq \eta+1 < j \leq \xi} \frac{t_{\rho(i)} - t_{\rho(j)} - 1}{t_{\rho(i)} - t_{\rho(j)}} \right. \\
&\quad \times \left(\prod_{m=2}^{\eta+1} T_{12}(t_{\rho(m)}) \otimes T_{12}(t_1) \prod_{l=\eta+2}^{\xi} T_{12}(t_{\rho(l)}) \right) \\
&\quad \times \left. \left(T_{22}(t_1) \prod_{i=\eta+2}^{\xi} T_{22}(t_{\rho(i)}) \otimes \prod_{j=2}^{\eta+1} T_{11}(t_{\rho(j)}) \right) \right]. \\
Y_4(\mathbf{t}) &= - \sum_{\eta=1}^{\xi-1} \frac{1}{(\xi-\eta-1)!\eta!} \\
&\times \sum_{\rho \in S'_{\xi-1}} \left[\sum_{k=2}^{\eta+1} \frac{1}{t_1 - t_{\rho(k)}} \prod_{1 < i \leq \eta+1 < j \leq \xi} \frac{t_{\rho(i)} - t_{\rho(j)} - 1}{t_{\rho(i)} - t_{\rho(j)}} \prod_{\substack{m=2 \\ m \neq k}}^{\eta+1} \frac{t_{\rho(k)} - t_{\rho(m)} + 1}{t_{\rho(k)} - t_{\rho(m)}} \right. \\
&\quad \times \left(T_{12}(t_1) \prod_{\substack{m=2 \\ m \neq k}}^{\eta+1} T_{12}(t_{\rho(m)}) \otimes T_{12}(t_1) \prod_{l=\eta+2}^{\xi} T_{12}(t_{\rho(l)}) \right) \\
&\quad \times \left. \left(T_{22}(t_{\rho(k)}) \prod_{i=\eta+2}^{\xi} T_{22}(t_{\rho(i)}) \otimes \prod_{j=2}^{\eta+1} T_{11}(t_{\rho(j)}) \right) \right].
\end{aligned}$$

To complete the proof, we will show that

$$Y_1(\mathbf{t}) + Y_3(\mathbf{t}) = \sum_{\eta=0}^{\xi} \frac{1}{(\xi-\eta)!\eta!} \sum_{\rho \in S_{\xi}} P_{\eta, \xi-\eta}(\mathbf{t}^{\rho}) \quad \text{and} \quad Y_2(\mathbf{t}) + Y_4(\mathbf{t}) = 0. \quad (\text{A.16})$$

We will start with the first equality in (A.16).

Observe that

$$Y_1(\mathbf{t}) = \sum_{\eta=1}^{\xi} \frac{1}{(\xi-\eta)!(\eta-1)!} \sum_{\substack{\sigma \in S_{\xi} \\ \sigma(1)=1}} P_{\eta, \xi-\eta}(\mathbf{t}^{\sigma}),$$

and

$$Y_3(\mathbf{t}) = \sum_{\eta=0}^{\xi-1} \frac{1}{(\xi-\eta-1)!\eta!} \sum_{\substack{\sigma \in S_{\xi} \\ \sigma(\eta+1)=1}} P_{\eta, \xi-\eta}(\mathbf{t}^{\sigma}).$$

On the other hand, we have

$$\begin{aligned} \sum_{\eta=0}^{\xi} \frac{1}{(\xi-\eta)! \eta!} \sum_{\sigma \in S_{\xi}} P_{\eta, \xi-\eta}(\mathbf{t}^{\sigma}) &= \sum_{\eta=1}^{\xi} \frac{1}{(\xi-\eta)! \eta!} \sum_{\substack{\sigma \in S_{\xi} \\ \sigma^{-1}(1) \in \{1, \dots, \eta\}}} P_{\eta, \xi-\eta}(\mathbf{t}^{\sigma}) \\ &+ \sum_{\eta=0}^{\xi-1} \frac{1}{(\xi-\eta)! \eta!} \sum_{\substack{\sigma \in S_{\xi} \\ \sigma^{-1}(1) \in \{\eta+1, \dots, \xi\}}} P_{\eta, \xi-\eta}(\mathbf{t}^{\sigma}). \end{aligned}$$

Denote by $s_{a,b} \in S_{\xi}$ the transposition of a and b . Then we have

$$\sum_{\substack{\sigma \in S_{\xi} \\ \sigma^{-1}(1) \in \{1, \dots, \eta\}}} P_{\eta, \xi-\eta}(\mathbf{t}^{\sigma}) = \sum_{l=1}^{\eta} \sum_{\substack{\tau \in S_{\xi}, \\ \tau(1)=1}} P_{\eta, \xi-\eta}(\mathbf{t}^{\tau s_{1,l}}) = \eta \sum_{\substack{\tau \in S_{\xi}, \\ \tau(1)=1}} P_{\eta, \xi-\eta}(\mathbf{t}^{\tau}).$$

For the first step, we used $l = \sigma^{-1}(1)$ and $\tau = \sigma s_{1, \sigma^{-1}(1)}$, so that $\tau(1) = 1$. For the second step, we used the equality $P_{\eta, \xi-\eta}(\mathbf{t}^{\tau s_{1,l}}) = P_{\eta, \xi-\eta}(\mathbf{t}^{\tau})$. Similarly,

$$\sum_{\substack{\sigma \in S_{\xi} \\ \sigma^{-1}(1) \in \{\eta+1, \dots, \xi\}}} P_{\eta, \xi-\eta}(\mathbf{t}^{\sigma}) = (\xi - \eta) \sum_{\substack{\rho \in S_{\xi}, \\ \rho(\eta+1)=1}} P_{\eta, \xi-\eta}(\mathbf{t}^{\rho}).$$

Therefore,

$$\begin{aligned} \sum_{\eta=0}^{\xi} \frac{1}{(\xi-\eta)! \eta!} \sum_{\sigma \in S_{\xi}} P_{\eta, \xi-\eta}(\mathbf{t}^{\sigma}) &= \sum_{\eta=1}^{\xi} \frac{1}{(\xi-\eta)! (\eta-1)!} \sum_{\substack{\sigma \in S_{\xi} \\ \sigma(1)=1}} P_{\eta, \xi-\eta}(\mathbf{t}^{\sigma}) \\ &+ \sum_{\eta=0}^{\xi-1} \frac{1}{(\xi-1-\eta)! \eta!} \sum_{\substack{\sigma \in S_{\xi} \\ \sigma(\eta+1)=1}} P_{\eta, \xi-\eta}(\mathbf{t}^{\sigma}) \\ &= Y_1(\mathbf{t}) + Y_3(\mathbf{t}). \end{aligned}$$

Finally, we show that $Y_2(\mathbf{t}) + Y_4(\mathbf{t}) = 0$. Observe that $Y_2(\mathbf{t})$ can be written as

$$\begin{aligned} Y_2(\mathbf{t}) &= \sum_{\eta=1}^{\xi-1} \frac{1}{(\xi-\eta)! (\eta-1)!} \\ &\times \sum_{l=\eta+1}^{\xi} \sum_{\tau \in S'_{\xi-1}} \left[\frac{1}{t_1 - t_{\tau(l)}} \prod_{1 < i \leq \eta < j \leq \xi} \frac{t_{\tau(i)} - t_{\tau(j)} - 1}{t_{\tau(i)} - t_{\tau(j)}} \prod_{\substack{m=\eta+1 \\ m \neq l}}^{\xi} \frac{t_{\tau(l)} - t_{\tau(m)} - 1}{t_{\tau(l)} - t_{\tau(m)}} \right. \\ &\times \left(T_{12}(t_1) \prod_{k=2}^{\eta} T_{12}(t_{\tau(k)}) \otimes T_{12}(t_1) \prod_{\substack{m=\eta+1 \\ m \neq l}}^{\xi} T_{12}(t_{\tau(m)}) \right) \\ &\times \left. \left(\prod_{i=\eta+1}^{\xi} T_{22}(t_{\tau(i)}) \otimes T_{11}(t_{\tau(l)}) \prod_{j=2}^{\eta} T_{11}(t_{\tau(j)}) \right) \right]. \end{aligned}$$

Changing the summation variable in the inner sum, $\tau = \sigma s_{l,\eta+1}$, we obtain that

$$\begin{aligned}
Y_2(\mathbf{t}) &= \sum_{\eta=1}^{\xi-1} \frac{1}{(\xi - \eta)!(\eta - 1)!} \\
&\times \sum_{l=\eta+1}^{\xi} \sum_{\sigma \in S'_{\xi-1}} \left[\frac{1}{t_1 - t_{\sigma(\eta+1)}} \prod_{m=\eta+2}^{\xi} \frac{t_{\sigma(\eta+1)} - t_{\sigma(m)} - 1}{t_{\sigma(\eta+1)} - t_{\sigma(m)}} \prod_{1 < i \leq \eta < j \leq \xi} \frac{t_{\sigma(i)} - t_{\sigma(j)} - 1}{t_{\sigma(i)} - t_{\sigma(j)}} \right. \\
&\quad \times \left(T_{12}(t_1) \prod_{k=2}^{\eta} T_{12}(t_{\sigma(k)}) \otimes T_{12}(t_1) \prod_{m=\eta+1}^{\xi} T_{12}(t_{\sigma(m)}) \right) \\
&\quad \left. \times \left(\prod_{i=\eta+2}^{\xi} T_{22}(t_{\sigma(i)}) \otimes \prod_{j=2}^{\eta+1} T_{11}(t_{\sigma(j)}) \right) \right].
\end{aligned}$$

The expression under the inner sum over σ does not depend on l and after a redistribution of factors, we get

$$\begin{aligned}
Y_2(\mathbf{t}) &= \sum_{\eta=1}^{\xi-1} \frac{1}{(\xi - \eta - 1)!(\eta - 1)!} \\
&\times \sum_{\sigma \in S'_{\xi-1}} \sum_{\sigma \in S'_{\xi-1}} \left[\frac{1}{t_1 - t_{\sigma(\eta+1)}} \prod_{m=\eta+2}^{\xi} \frac{t_{\sigma(\eta+1)} - t_{\sigma(m)} - 1}{t_{\sigma(\eta+1)} - t_{\sigma(m)}} \prod_{i=2}^{\eta} \frac{t_{\sigma(i)} - t_{\sigma(\eta+1)} - 1}{t_{\sigma(i)} - t_{\sigma(\eta+1)}} \right. \\
&\quad \times \prod_{1 < i < \eta+1 < j \leq \xi} \frac{t_{\sigma(i)} - t_{\sigma(j)} - 1}{t_{\sigma(i)} - t_{\sigma(j)}} \\
&\quad \times \left(T_{12}(t_1) \prod_{k=2}^{\eta} T_{12}(t_{\sigma(k)}) \otimes T_{12}(t_1) \prod_{m=\eta+1}^{\xi} T_{12}(t_{\sigma(m)}) \right) \\
&\quad \left. \times \left(\prod_{i=\eta+2}^{\xi} T_{22}(t_{\sigma(i)}) \otimes \prod_{j=2}^{\eta+1} T_{11}(t_{\sigma(j)}) \right) \right]. \tag{A.17}
\end{aligned}$$

Similarly, $Y_4(\mathbf{t})$ can be written as

$$\begin{aligned}
Y_4(\mathbf{t}) &= \sum_{\eta=1}^{\xi-1} \frac{1}{(\xi - 1 - \eta)! \eta!} \\
&\times \sum_{k=2}^{\eta+1} \sum_{\rho \in S'_{\xi-1}} \left[\frac{1}{t_1 - t_{\rho(k)}} \prod_{1 < i \leq \eta+1 < j \leq \xi} \frac{t_{\rho(i)} - t_{\rho(j)} - 1}{t_{\rho(i)} - t_{\rho(j)}} \prod_{\substack{m=2 \\ m \neq k}}^{\eta+1} \frac{t_{\rho(k)} - t_{\rho(m)} + 1}{t_{\rho(k)} - t_{\rho(m)}} \right. \\
&\quad \times \left(T_{12}(t_1) \prod_{\substack{m=2 \\ m \neq k}}^{\eta+1} T_{12}(t_{\rho(m)}) \otimes T_{12}(t_1) \prod_{l=\eta+2}^{\xi} T_{12}(t_{\rho(l)}) \right) \\
&\quad \left. \times \left(T_{22}(t_{\rho(k)}) \prod_{i=\eta+2}^{\xi} T_{22}(t_{\rho(i)}) \otimes \prod_{j=2}^{\eta+1} T_{11}(t_{\rho(j)}) \right) \right],
\end{aligned}$$

and changing the summation variable in the inner sum, $\rho = \sigma s_{k,\eta+1}$, we get

$$Y_4(\mathbf{t}) = - \sum_{\eta=1}^{\xi-1} \frac{1}{(\xi - 1 - \eta)! \eta!}$$

$$\begin{aligned}
& \times \sum_{k=2}^{\eta+1} \sum_{\sigma \in S'_{\xi-1}} \left[\frac{1}{t_1 - t_{\sigma(\eta+1)}} \prod_{m=2}^{\eta} \frac{t_{\sigma(\eta+1)} - t_{\sigma(m)} + 1}{t_{\sigma(\eta+1)} - t_{\sigma(m)}} \prod_{1 < i \leq \eta+1 < j \leq \xi} \frac{t_{\sigma(i)} - t_{\sigma(j)} - 1}{t_{\sigma(i)} - t_{\sigma(j)}} \right. \\
& \quad \times \left(T_{12}(t_1) \prod_{m=2}^{\eta} T_{12}(t_{\sigma(m)}) \otimes T_{12}(t_1) \prod_{l=\eta+2}^{\xi} T_{12}(t_{\sigma(l)}) \right) \\
& \quad \times \left. \left(\prod_{i=\eta+1}^{\xi} T_{22}(t_{\sigma(i)}) \otimes \prod_{j=2}^{\eta+1} T_{11}(t_{\sigma(j)}) \right) \right].
\end{aligned}$$

The expression under the inner sum over σ does not depend on k and after a redistribution of factors, we get

$$\begin{aligned}
Y_4(\mathbf{t}) &= - \sum_{\eta=1}^{\xi-1} \frac{1}{(\xi - \eta - 1)!(\eta - 1)!} \\
& \times \sum_{\sigma \in S'_{\xi-1}} \left[\frac{1}{t_1 - t_{\sigma(\eta+1)}} \prod_{m=2}^{\eta} \frac{t_{\sigma(\eta+1)} - t_{\sigma(m)} + 1}{t_{\sigma(\eta+1)} - t_{\sigma(m)}} \prod_{q=\eta+2}^{\xi} \frac{t_{\sigma(\eta+1)} - t_{\sigma(q)} - 1}{t_{\sigma(\eta+1)} - t_{\sigma(q)}} \right. \\
& \quad \times \prod_{1 < i < \eta+1 < j \leq \xi} \frac{t_{\sigma(i)} - t_{\sigma(j)} - 1}{t_{\sigma(i)} - t_{\sigma(j)}} \\
& \quad \times \left(T_{12}(t_1) \prod_{m=2}^{\eta} T_{12}(t_{\sigma(m)}) \otimes T_{12}(t_1) \prod_{l=\eta+2}^{\xi} T_{12}(t_{\sigma(l)}) \right) \\
& \quad \times \left. \left(\prod_{i=\eta+1}^{\xi} T_{22}(t_{\sigma(i)}) \otimes \prod_{j=2}^{\eta+1} T_{11}(t_{\sigma(j)}) \right) \right]. \tag{A.18}
\end{aligned}$$

Formulae (A.17) and (A.18) show that $Y_2(\mathbf{t}) + Y_4(\mathbf{t}) = 0$. This completes the proof of formula (A.12). Proposition A.2 is proved. \blacksquare

Acknowledgements

The authors thank the referees for very careful reading of this paper and their valuable suggestions. The second author is supported in part by Simons Foundation grants 430235, 852996.

References

- [1] Felder G., Rimányi R., Varchenko A., Poincaré–Birkhoff–Witt expansions of the canonical elliptic differential form, in *Quantum Groups, Contemp. Math.*, Vol. 433, [American Mathematical Society](#), Providence, RI, 2007, 191–208, [arXiv:math.RT/0502296](#).
- [2] Khoroshkin S., Pakuliak S., Generating series for nested Bethe vectors, *SIGMA* **4** (2008), 081, 23 pages, [arXiv:0810.3131](#).
- [3] Khoroshkin S., Pakuliak S., Tarasov V., Off-shell Bethe vectors and Drinfeld currents, *J. Geom. Phys.* **57** (2007), 1713–1732, [arXiv:math.QA/0610517](#).
- [4] Korepin V.E., Calculation of norms of Bethe wave functions, *Comm. Math. Phys.* **86** (1982), 391–418.
- [5] Kosmakov M., Tarasov V., New combinatorial formulae for nested Bethe vectors II, *Lett. Math. Phys.* **115** (2025), 12, 20 pages, [arXiv:2402.15717](#).
- [6] Kulish P.P., Reshetikhin N.Yu., Diagonalisation of $GL(N)$ invariant transfer matrices and quantum N -wave system (Lee model), *J. Phys. A* **16** (1983), L591–L596.
- [7] Kulish P.P., Reshetikhin N.Yu., GL_3 -invariant solutions of the Yang–Baxter equation and associated quantum systems, *J. Sov. Math.* **34** (1986), 1948–1971.

- [8] Markov Y., Varchenko A., Hypergeometric solutions of trigonometric Knizhnik–Zamolodchikov equations satisfy dynamical difference equations, *Adv. Math.* **166** (2002), 100–147.
- [9] Matsuo A., An application of Aomoto–Gel’fand hypergeometric functions to the $SU(n)$ Knizhnik–Zamolodchikov equation, *Comm. Math. Phys.* **134** (1990), 65–77.
- [10] Miwa T., Takeyama Y., Tarasov V., Determinant formula for solutions of the quantum Knizhnik–Zamolodchikov equation associated with $U_q(\mathfrak{sl}_n)$ at $|q| = 1$, *Publ. Res. Inst. Math. Sci.* **35** (1999), 871–892, [arXiv:math.QA/9812096](#).
- [11] Mukhin E., Tarasov V., Varchenko A., Bethe eigenvectors of higher transfer matrices, *J. Stat. Mech. Theory Exp.* **2006** (2006), P08002, 44 pages, [arXiv:math.QA/0605015](#).
- [12] Mukhin E., Tarasov V., Varchenko A., Spaces of quasi-exponentials and representations of the Yangian $Y(\mathfrak{gl}_N)$, *Transform. Groups* **19** (2014), 861–885, [arXiv:1303.1578](#).
- [13] Oskin A., Pakuliak S., Silantyev A., On the universal weight function for the quantum affine algebra $U_q(\widehat{\mathfrak{gl}}_N)$, *St. Petersburg Math. J.* **21** (2009), 651–680, [arXiv:0711.2821](#).
- [14] Rimányi R., Stevens L., Varchenko A., Combinatorics of rational functions and Poincaré–Birchoff–Witt expansions of the canonical $U(\mathfrak{n}_-)$ -valued differential form, *Ann. Comb.* **9** (2005), 57–74, [arXiv:math.CO/0407101](#).
- [15] Rimányi R., Tarasov V., Varchenko A., Cohomology classes of conormal bundles of Schubert varieties and Yangian weight functions, *Math. Z.* **277** (2014), 1085–1104, [arXiv:1204.4961](#).
- [16] Rimányi R., Tarasov V., Varchenko A., Partial flag varieties, stable envelopes, and weight functions, *Quantum Topol.* **6** (2015), 333–364, [arXiv:1212.6240](#).
- [17] Rimányi R., Tarasov V., Varchenko A., Trigonometric weight functions as K -theoretic stable envelope maps for the cotangent bundle of a flag variety, *J. Geom. Phys.* **94** (2015), 81–119, [arXiv:1411.0478](#).
- [18] Rimányi R., Tarasov V., Varchenko A., Elliptic and K -theoretic stable envelopes and Newton polytopes, *Selecta Math. (N.S.)* **25** (2019), 16, 43 pages, [arXiv:1705.09344](#).
- [19] Schechtman V., Varchenko A., Hypergeometric solutions of Knizhnik–Zamolodchikov equations, *Lett. Math. Phys.* **20** (1990), 279–283.
- [20] Schechtman V., Varchenko A., Arrangements of hyperplanes and Lie algebra homology, *Invent. Math.* **106** (1991), 139–194.
- [21] Slavnov N., Algebraic Bethe ansatz, [arXiv:1804.07350](#).
- [22] Slavnov N., Algebraic Bethe ansatz and correlation functions—an advanced course, *World Scientific Publishing*, Hackensack, NJ, 2022.
- [23] Takhtadzhian L.A., Faddeev L.D., The quantum method for the inverse problem and the XYZ Heisenberg model, *Russ. Math. Surv.* **34** (1979), 11–68.
- [24] Tarasov V., Varchenko A., Selberg-type integrals associated with \mathfrak{sl}_3 , *Lett. Math. Phys.* **65** (2003), 173–185, [arXiv:math.QA/0302148](#).
- [25] Tarasov V., Varchenko A., Combinatorial formulae for nested Bethe vectors, *SIGMA* **9** (2013), 048, 28 pages, [arXiv:math.QA/0702277](#).
- [26] Tarasov V., Varchenko A., Hypergeometric solutions of the quantum differential equation of the cotangent bundle of a partial flag variety, *Cent. Eur. J. Math.* **12** (2014), 694–710, [arXiv:1301.2705](#).
- [27] Tarasov V., Varchenko A., q -hypergeometric solutions of quantum differential equations, quantum Pieri rules, and gamma theorem, *J. Geom. Phys.* **142** (2019), 179–212, [arXiv:1710.03177](#).
- [28] Varchenko A., Tarasov V., Jackson integral representations for solutions of the Knizhnik–Zamolodchikov quantum equation, *St. Petersburg Math. J.* **6** (1994), 275–313, [arXiv:hep-th/9311040](#).