

## IS ANY KIND OF MIXING POSSIBLE IN $ToP$ $\mathbb{N}$ -ACTIONS?

TOMASZ DOWNAROWICZ AND XIANGDONG YE

**Definition.** A topological dynamical system  $(X, T)$ , where  $X$  a compact space without isolated points and  $T$  a continuous map from  $X$  to itself, is called  $ToP$  if every point  $x \in X$  is either (topologically) transitive or periodic and both types are present.

Note that transitivity of  $(X, T)$  (i.e., there exists a point with dense orbit) has different meaning depending on whether we consider the  $\mathbb{N}$ -action (positive iterates of a continuous map  $T$ ) or  $\mathbb{Z}$ -action (both positive and negative iterates of a homeomorphism  $T$ ). Thus the notion  $ToP$  has different meanings in these cases, too. In a forthcoming paper we have constructed  $ToP$  systems with positive entropy and for  $\mathbb{Z}$ -action we know how to make one which is topologically weakly mixing. For  $\mathbb{N}$ -action our examples are not totally transitive!

**Question:** *Do there exist totally transitive (topologically weakly mixing or mixing)  $ToP$   $\mathbb{N}$ -actions?*

INSTYTUT MATEMATYKI, POLITECHNIKI WROCLAWSKIEJ, WYBRZEZE WYSPIANSKIEGO 27,  
50-370 WROCLAW, POLAND

*E-mail address:* `downar@im.pwr.wroc.pl`

UNIVERSITY OF SCIENCE & TECHNOLOGY OF CHINA, P.R. CHINA

*E-mail address:* `yexd@ustc.edu.cn`

Typeset by  $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{T}\mathcal{E}\mathcal{X}$